

[Course](#)

[Progress](#)

[Dates](#)

[Discussion](#)

[Syllabus](#)

[Outline](#)

[laff routines](#)

[Community](#)

[🏠 Course](#) / [Week 7: More Gaussian Elimination and Matrix Inversi...](#) / [7.3 The Inverse Mat...](#)



< Previous	 	 	 		 	Next >
-------------------------------	---	---	---	---	---	---------------------------

7.3.4 More Advanced (But Still Simple) Examples

 Bookmark this page

Week 7 due Nov 20, 2023 01:42 IST Completed

7.3.4 More Advanced (But Still Simple) Examples

Video

Start of transcript. Skip to the end.

▶

0:00 / 0:00

▶ 2.0x

🔊

⌂

CC

“

Dr. Robert van de Geijn: So now we're going to look at slightly more complex examples. And this is all in preparation for being able to discuss the general case next week. So here's the idea. If we have a matrix A, can we multiply it

Video

📄 [Download video file](#)

Transcripts

📄 [Download SubRip \(.srt\) file](#)

📄 [Download Text \(.txt\) file](#)

Reading Assignment

0 points possible (ungraded)
Read Unit 7.3.4 of the notes. [\[LINK\]](#)

☒ Done

✓

Submit

Discussion

Topic: Week 7 / 7.3.4

Hide Discussion

Add a Post

Show all posts ▼

by recent activity ▼

There are no posts in this topic yet.

✕

Calculator

Homework 7.3.4.1

1/1 point (graded)

Find $\begin{pmatrix} -2 & 0 \\ 4 & 2 \end{pmatrix}^{-1} =$

☐ $\begin{pmatrix} -\frac{1}{2} & 1 \\ 0 & \frac{1}{2} \end{pmatrix}$

☐ $\begin{pmatrix} \frac{1}{2} & 0 \\ 1 & \frac{1}{2} \end{pmatrix}$

☒ $\begin{pmatrix} -\frac{1}{2} & 0 \\ 1 & \frac{1}{2} \end{pmatrix}$

☐ $\begin{pmatrix} -\frac{1}{2} & 0 \\ 1 & -\frac{1}{2} \end{pmatrix}$



Homework 7.3.4.1 Compute $\begin{pmatrix} -2 & 0 \\ 4 & 2 \end{pmatrix}^{-1} = \begin{pmatrix} -\frac{1}{2} & 0 \\ 1 & \frac{1}{2} \end{pmatrix}.$

Answer: Here is how you can find the answer First, solve

$$\begin{pmatrix} -2 & 0 \\ 4 & 2 \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

If you do forward substitution, you see that the solution is $\begin{pmatrix} -\frac{1}{2} \\ 1 \end{pmatrix}$, which becomes the first column of A^{-1} .

Next, solve

$$\begin{pmatrix} -2 & 0 \\ 4 & 2 \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

If you do forward substitution, you see that the solution is $\begin{pmatrix} 0 \\ \frac{1}{2} \end{pmatrix}$, which becomes the second column of A^{-1} .

Check:

$$\begin{pmatrix} -2 & 0 \\ 4 & 2 \end{pmatrix} \begin{pmatrix} -\frac{1}{2} & 0 \\ 1 & \frac{1}{2} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Video

[Start of transcript. Skip to the end.](#)



▶

0:00 / 0:00

▶ 2.0x

🔊

🔍

📺

💬

Dr. Robert van de Geijn: OK, so the way to compute this inverse

is to note that the lower triangular matrix times its inverse partition

by columns must be equal to the identity partition by its columns.

That's what we talked about in the last video.

If you then say, well, I want to solve with the lower triangular matrix here

Video

📄 [Download video file](#)

Transcripts

📄 [Download SubRip \(.srt\) file](#)

📄 [Download Text \(.txt\) file](#)

Homework 7.3.4.2

1/1 point (graded)

Find $\begin{pmatrix} 1 & -2 \\ 0 & 2 \end{pmatrix}^{-1} =$

- ☐ $\begin{pmatrix} 1 & -\frac{1}{2} \\ 0 & \frac{1}{2} \end{pmatrix}$
- ☐ $\begin{pmatrix} 1 & -\frac{1}{2} \\ 0 & 1 \end{pmatrix}$
- ☒ $\begin{pmatrix} 1 & 1 \\ 0 & \frac{1}{2} \end{pmatrix}$
- ☐ $\begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}$



Answer:

$$\begin{pmatrix} 1 & 1 \\ 0 & \frac{1}{2} \end{pmatrix}.$$

Here is how you can find this matrix: First, solve

$$\begin{pmatrix} 1 & -2 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

Calculator

$$\begin{pmatrix} 1 & -2 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} \chi_0 \\ \chi_1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

If you do back substitution, you see that the solution is $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$, which becomes the first column of A^{-1} .
Next, solve

$$\begin{pmatrix} 1 & -2 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} \chi_0 \\ \chi_1 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

If you do back substitution, you see that the solution is $\begin{pmatrix} 1 \\ \frac{1}{2} \end{pmatrix}$, which becomes the second column of A^{-1} .
Check:

$$\begin{pmatrix} 1 & -2 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & \frac{1}{2} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Submit

i Answers are displayed within the problem

Homework 7.3.4.3

1/1 point (graded)

Let $\alpha_{0,0} \neq 0$ and $\alpha_{1,1} \neq 0$. Then $\begin{pmatrix} \alpha_{0,0} & 0 \\ \alpha_{1,0} & \alpha_{1,1} \end{pmatrix}^{-1} = \begin{pmatrix} \frac{1}{\alpha_{0,0}} & 0 \\ -\frac{\alpha_{1,0}}{\alpha_{0,0}\alpha_{1,1}} & \frac{1}{\alpha_{1,1}} \end{pmatrix}$

True

✔ Answer: True

Answer: **True**
Here is how you can find the matrix: First, solve

$$\begin{pmatrix} \alpha_{0,0} & 0 \\ \alpha_{1,0} & \alpha_{1,1} \end{pmatrix} \begin{pmatrix} \chi_0 \\ \chi_1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

If you solve the lower triangular system, you see that the solution is $\begin{pmatrix} \frac{1}{\alpha_{0,0}} \\ -\frac{\alpha_{1,0}}{\alpha_{0,0}\alpha_{1,1}} \end{pmatrix}$, which becomes the first column of A^{-1} .
Next, solve

$$\begin{pmatrix} \alpha_{0,0} & 0 \\ \alpha_{1,0} & \alpha_{1,1} \end{pmatrix} \begin{pmatrix} \chi_0 \\ \chi_1 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

If you solve this system, you find the solution $\begin{pmatrix} 0 \\ \frac{1}{\alpha_{1,1}} \end{pmatrix}$, which becomes the second column of A^{-1} .
Check:

$$\begin{pmatrix} \alpha_{0,0} & 0 \\ \alpha_{1,0} & \alpha_{1,1} \end{pmatrix} \begin{pmatrix} \frac{1}{\alpha_{0,0}} & 0 \\ -\frac{\alpha_{1,0}}{\alpha_{0,0}\alpha_{1,1}} & \frac{1}{\alpha_{1,1}} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Submit

 Calculator

i Answers are displayed within the problem

Homework 7.3.4.4

1/1 point (graded)
Partition lower triangular matrix L as

$$L = \left(\begin{array}{c|c} L_{00} & 0 \\ \hline l_{10}^T & \lambda_{11} \end{array} \right)$$

Assume that L has no zeroes on its diagonal. Then

$$L^{-1} = \left(\begin{array}{c|c} L_{00}^{-1} & 0 \\ \hline -\frac{l_{10}^T L_{00}^{-1}}{\lambda_{11}} & \frac{1}{\lambda_{11}} \end{array} \right)$$

True

✓ Answer: True

Answer: **True** Stictly speaking, one needs to show that L_{00} has an inverse... This would require a proof by induction. We'll skip that part. Instead, we'll just multiply out:

$$\begin{aligned} \left(\begin{array}{c|c} L_{00} & 0 \\ \hline l_{10}^T & \lambda_{11} \end{array} \right) \left(\begin{array}{c|c} L_{00}^{-1} & 0 \\ \hline -\frac{l_{10}^T L_{00}^{-1}}{\lambda_{11}} & \frac{1}{\lambda_{11}} \end{array} \right) &= \left(\begin{array}{c|c} L_{00} L_{00}^{-1} & L_{00} 0 + 0 \frac{1}{\lambda_{11}} \\ \hline l_{10}^T L_{00}^{-1} - \lambda_{11} \frac{l_{10}^T L_{00}^{-1}}{\lambda_{11}} & l_{10}^T \times 0 + \lambda_{11} \frac{1}{\lambda_{11}} \end{array} \right) \\ &= \left(\begin{array}{c|c} I & 0 \\ \hline 0 & 1 \end{array} \right). \end{aligned}$$

Submit

i Answers are displayed within the problem

Video

[Start of transcript. Skip to the end.](#)



▶ 0:00 / 0:00

▶ 2.0x

🔊

⌂

CC

“

Dr. Robert van de Geijn: So now we get to a more general case.

And the exercise is to show that the inverse of this lower triangular matrix partitioned like that is given by this.

The way to verify that that is indeed the inverse

is a matter of just multiplying the two matrices together.

Video
📄 [Download video file](#)

Transcripts
📄 [Download SubRip \(.srt\) file](#)
📄 [Download Text \(.txt\) file](#)

Calculator

Homework 7.3.4.5

1/1 point (graded)

The inverse of a lower triangular matrix with no zeroes on its diagonal is a lower triangular matrix.

TRUE



✔ Answer: TRUE

Answer: TrueProof by induction on n , the size of the square matrix.Let L be the lower triangular matrix.**Base case:** $n = 1$. Then $L = (\lambda_{11})$, with $\lambda_{11} \neq 0$. Clearly, $L^{-1} = (1/\lambda_{11})$.**Inductive step:** Inductive Hypothesis: Assume that the inverse of any $n \times n$ lower triangular matrix with no zeroes on its diagonal is a lower triangular matrix.We need to show that the inverse of any $(n+1) \times (n+1)$ lower triangular matrix, L , with no zeroes on its diagonal is a lower triangular matrix.

Partition

$$L = \left(\begin{array}{c|c} L_{00} & 0 \\ \hline l_{10}^T & \lambda_{11} \end{array} \right)$$

We know then that L_{00} has no zeroes on its diagonal and $\lambda_{11} \neq 0$. We also saw that then

$$L^{-1} = \left(\begin{array}{c|c} L_{00}^{-1} & 0 \\ \hline -\frac{1}{\lambda_{11}} l_{10}^T L_{00}^{-1} & \frac{1}{\lambda_{11}} \end{array} \right)$$

Hence, the matrix has an inverse, and it is lower triangular.

By the **Principle of Mathematical Induction**, the result holds.

Submit

Answers are displayed within the problem

The answer to the last exercise suggests an algorithm for inverting a lower triangular matrix. See if you can implement it!

Homework 7.3.4.7

1/1 point (graded)

Find $\begin{pmatrix} 1 & 2 \\ 1 & 1 \end{pmatrix}^{-1} =$

☐ $\begin{pmatrix} 1 & 2 \\ 1 & 1 \end{pmatrix}$

☒ $\begin{pmatrix} -1 & 2 \\ 1 & -1 \end{pmatrix}$

☐ $\begin{pmatrix} -1 & \frac{1}{2} \\ 1 & -1 \end{pmatrix}$

Calculator

☐ $\begin{pmatrix} 1 & \frac{1}{2} \\ 1 & 1 \end{pmatrix}$



Answer:

$$\begin{pmatrix} -1 & 2 \\ 1 & -1 \end{pmatrix}.$$

Here is how you can find this matrix: First, you compute the LU factorization. Since there is only one step, this is easy:

$$\begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ 0 & -1 \end{pmatrix}.$$

Thus,

$$\begin{pmatrix} 1 & 2 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 0 & -1 \end{pmatrix}.$$

Next, you solve

$$\begin{pmatrix} 1 & 2 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} \chi_0 \\ \chi_1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

by solving $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} \zeta_0 \\ \zeta_1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ followed by $\begin{pmatrix} 1 & 2 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} \chi_0 \\ \chi_1 \end{pmatrix} = \begin{pmatrix} \zeta_0 \\ \zeta_1 \end{pmatrix}$. If you do this right, you get $\begin{pmatrix} -1 \\ 1 \end{pmatrix}$, which becomes the first column of the inverse.

You solve

$$\begin{pmatrix} 1 & 2 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} \chi_0 \\ \chi_1 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

in a similar manner, yielding the second column of the inverse, $\begin{pmatrix} 2 \\ -1 \end{pmatrix}$.

Check:

$$\begin{pmatrix} 1 & 2 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} -1 & 2 \\ 1 & -1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Submit

i Answers are displayed within the problem

Homework 7.3.4.8

1/1 point (graded)

If $\alpha_{0,0}\alpha_{1,1} - \alpha_{1,0}\alpha_{0,1} \neq 0$ then

$$\begin{pmatrix} \alpha_{0,0} & \alpha_{0,1} \end{pmatrix}^{-1} = \frac{1}{\alpha_{0,0}\alpha_{1,1} - \alpha_{1,0}\alpha_{0,1}} \begin{pmatrix} \alpha_{1,1} & -\alpha_{0,1} \\ \alpha_{1,0} & -\alpha_{0,0} \end{pmatrix}$$

Calculator

$$\begin{pmatrix} \alpha_{1,0} & \alpha_{1,1} \end{pmatrix}^{-1} = \frac{1}{\alpha_{0,0}\alpha_{1,1} - \alpha_{1,0}\alpha_{0,1}} \begin{pmatrix} -\alpha_{1,0} & \alpha_{0,0} \end{pmatrix}$$

True

✔ Answer: True

Answer: True
Check:

$$\begin{pmatrix} \alpha_{0,0} & \alpha_{0,1} \\ \alpha_{1,0} & \alpha_{1,1} \end{pmatrix} \frac{1}{\alpha_{0,0}\alpha_{1,1} - \alpha_{1,0}\alpha_{0,1}} \begin{pmatrix} \alpha_{1,1} & -\alpha_{0,1} \\ -\alpha_{1,0} & \alpha_{0,0} \end{pmatrix}$$
$$= \frac{1}{\alpha_{0,0}\alpha_{1,1} - \alpha_{1,0}\alpha_{0,1}} \begin{pmatrix} \alpha_{0,0} & \alpha_{0,1} \\ \alpha_{1,0} & \alpha_{1,1} \end{pmatrix} \begin{pmatrix} \alpha_{1,1} & -\alpha_{0,1} \\ -\alpha_{1,0} & \alpha_{0,0} \end{pmatrix}$$
$$= \frac{1}{\alpha_{0,0}\alpha_{1,1} - \alpha_{1,0}\alpha_{0,1}} \begin{pmatrix} \alpha_{0,0}\alpha_{1,1} - \alpha_{0,1}\alpha_{1,0} & -\alpha_{0,0}\alpha_{0,1} + \alpha_{0,1}\alpha_{0,0} \\ \alpha_{1,0}\alpha_{1,1} - \alpha_{1,1}\alpha_{1,0} & -\alpha_{1,0}\alpha_{0,1} + \alpha_{1,1}\alpha_{0,0} \end{pmatrix}$$
$$= \frac{1}{\alpha_{0,0}\alpha_{1,1} - \alpha_{1,0}\alpha_{0,1}} \begin{pmatrix} \alpha_{0,0}\alpha_{1,1} - \alpha_{0,1}\alpha_{1,0} & 0 \\ 0 & \alpha_{1,1}\alpha_{0,0} - \alpha_{1,0}\alpha_{0,1} \end{pmatrix}$$
$$= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Submit

Answers are displayed within the problem

Homework 7.3.4.9

1/1 point (graded)

The 2×2 matrix $A = \begin{pmatrix} \alpha_{0,0} & \alpha_{0,1} \\ \alpha_{1,0} & \alpha_{1,1} \end{pmatrix}$ has an inverse if and only if $\alpha_{0,0}\alpha_{1,1} - \alpha_{1,0}\alpha_{0,1} \neq 0$.

True

✔ Answer: True

Answer: True
This is an immediate consequence of the last exercise.

Submit


Answers are displayed within the problem

In the below video, towards the end, the instructor misspeaks and says that the inverse exists if and only if the determinant is zero. The correct statement is that the inverse exists if and only if the determinant is **not** zero.

Video

Start of transcript. Skip to the end.

Calculator



is given by what's on the right, right here.

And notice that that formula can only be computed if this right here



Next >

[News](#)

Your Privacy Choices

Media Kit





© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)