

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Unit 5 Reinforcement Learning \(2 weeks\)](#)

[Lecture 17. Reinforcement Learning](#)
> [1](#)


> 6. Bellman Equations

6. Bellman Equations

Bellman Equations

[Start of transcript. Skip to the end.](#)

So now we will start introducing a tiny bit more notation.



Just three.

I promise to discuss Bellman equations.

So the first annotation that I will introduce,

as I already alluded, is called V . V star.

This is the value of state.

What does it mean?

It tells you the value of the expected reward



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

In this problem, we work through a numerical example to understand the Bellman equations:

Let there be 4 possible actions, a_1, a_2, a_3, a_4 from a given state s and assume while answering the following problems that the Q^* values are given as below:

$$Q^*(s, a_1) = 10Q^*(s, a_2) = -1Q^*(s, a_3) = 0Q^*(s, a_4) = 11$$

Recall from the lecture that,

$$V^*(s) = \max_a Q^*(s, a)$$

$$Q^*(s, a) = \sum_{s'} T(s, a, s') (R(s, a, s') + \gamma V^*(s'))$$

Value function in terms of Q function

1/1 point (graded)

Enter the value of $V^*(s)$ below:

✓ Answer: 11

Solution:

Note that $V^*(s)$ is given by:

$$V^*(s) = \max_a Q^*(s, a)$$

$$V^*(s) = \max(10, -1, 0, 11) = 11$$

Submit

You have used 1 of 2 attempts

i Answers are displayed within the problem

Bellman Equation for Q function

1/1 point (graded)

Let s' be a state that can be reached from s by taking the action a_1 . Also assume that

$$T(s, a_1, s') = 1R(s, a_1, s') = 5\gamma = 0.5$$

Enter the value of $V^*(s')$ below:

10

✓ Answer: 10

Solution:

Note that since T denotes probabilities, the following must be true:

$$\sum_{s'} T(s, a, s') = 1$$

. Also,

$$Q^*(s, a) = \sum_{s'} T(s, a, s') (R(s, a, s') + \gamma V^*(s'))$$

Since, $T(s, a_1, s') = 1$ and $\sum_{s'} T(s, a, s') = 1$, we would have $T(s, a_1, s'') = 0 \quad \forall s'' \neq s'$.

The above equation would then reduce as follows

$$Q^*(s, a_1) = T(s, a_1, s') (R(s, a_1, s') + \gamma V^*(s'))$$

$$10 = 1 * (5 + 0.5 * V^*(s'))$$

$$V^*(s') = 5/0.5 = 10$$

Submit

You have used 2 of 3 attempts

i Answers are displayed within the problem

Discussion

[Hide Discussion](#)

Topic: Unit 5 Reinforcement Learning (2 weeks) ;Lecture 17. Reinforcement Learning 1 / 6.
Bellman Equations

[Add a Post](#)

Show all posts ▼

by recent activity ▼

? The values are missing spacing and are hard to read

1

© All Rights Reserved