

[Course](#)

[Progress](#)

[Dates](#)

[Discussion](#)

[Syllabus](#)

[Outline](#)

[laff routines](#)

[Community](#)

 [Course](#) / [Week 10: Vector Spaces, Orthogonality, and Lin...](#) / [10.3 Orthogonal Vectors...](#)



< Previous







Next >

# 10.3.1 Orthogonal Vectors

 Bookmark this page

Week 10 due Dec 16, 2023 07:42 IST   Completed

# 10.3.1 Orthogonal Vectors

## Video

[Start of transcript. Skip to the end.](#)



Dr. Robert van de Geijn: Next we're going to revisit orthogonal vectors. Now we talked a little bit about orthogonal vectors in week one when we talked about the dot product. And we linked the dot product of two vectors being equal to 0, two vectors being orthogonal or perpendicular.

## Video

[Download video file](#)

## Transcripts

- [Download SubRip \(.srt\) file](#)
- [Download Text \(.txt\) file](#)

## Reading Assignment

0 points possible (ungraded)  
Read Unit 10.3.1 of the notes. [\[LINK\]](#)

☒ Done

✓

Submit

✓ Correct

## Discussion

Topic: Week 10 / 10.3.1

Hide Discussion

Add a Post

Show all posts	by recent activity
<input checked="" type="checkbox"/> Question on lecture note 10.3.1	5
Greetings, I was curious why under the section 10.3.1 - orthogonal vectors in the page 364 of our lecture note (pdf), you restrict the $x, y$ to non-...	

## Homework 10.3.1.1

Calculator

Homework 10.3.1.1

4/4 points (graded)

For each of the following, indicate whether the vectors are orthogonal:

•  $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$  and  $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$

TRUE

▼

✔ Answer: TRUE

•  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$

TRUE

▼

✔ Answer: TRUE

• The unit basis vectors  $e_i$  and  $e_j$ .

Sometimes

▼

✔ Answer: Sometimes

•  $\begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}$  and  $\begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix}$

Always

▼

✔ Answer: Always

$\begin{pmatrix} 1 \\ -1 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$	True because $\begin{pmatrix} 1 \\ -1 \end{pmatrix}^T \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 0$
$\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$	True because $\begin{pmatrix} 1 \\ 0 \end{pmatrix}^T \begin{pmatrix} 0 \\ 1 \end{pmatrix} = 0$
The unit basis vectors $e_i$ and $e_j$ .	Sometimes because $e_i^T e_j = 0$ if $i \neq j$ but $e_i^T e_j = 1$ if $i = j$ .
$\begin{pmatrix} c \\ s \end{pmatrix}$ and $\begin{pmatrix} -s \\ c \end{pmatrix}$	Always because $\begin{pmatrix} c \\ s \end{pmatrix}^T \begin{pmatrix} -s \\ c \end{pmatrix} = 0$

Submit

ⓘ Answers are displayed within the problem

Homework 10.3.1.2

1/1 point (graded)

Let  $A \in \mathbb{R}^{m \times n}$ . Let  $a_i^T$  be a row of  $A$  and  $x \in \mathcal{N}(A)$ . Then  $a_i$  is orthogonal to  $x$ .

Always

▼

✔ Answer: Always

Answer: Always Since  $x \in \mathcal{N}(A)$ ,  $Ax = 0$ . But then, partitioning  $A$  by rows,

$$0 = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix} = Ax = \begin{pmatrix} a_0^T \\ a_1^T \\ \vdots \\ a_i^T \end{pmatrix} x = \begin{pmatrix} a_0^T x \\ a_1^T x \\ \vdots \\ a_i^T x \end{pmatrix}.$$

< Previous

Next >



## edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

## Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)
- [Cookie Policy](#)
- [Your Privacy Choices](#)

## Connect

- [Idea Hub](#)
- [Contact Us](#)
- [Help Center](#)
- [Security](#)
- [Media Kit](#)



© 2023 edX LLC. All rights reserved.  
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)