

[Course](#)

[Progress](#)

[Dates](#)

[Discussion](#)

[Syllabus](#)

[Outline](#)

[laff routines](#)

[Community](#)



[Course](#) / [Week 8: More on Matrix Inversion](#) / [8.2 Gauss-Jordan Elimination](#)



| | | | | | | | | |
|------------|---|---|---|---|---|---|---|--------|
| < Previous |  ✓ |  ✓ |  ✓ |  |  |  |  | Next > |
|------------|---|---|---|---|---|---|---|--------|

8.2.4 Computing the Inverse of A via Gauss-Jordan Elimination

 Bookmark this page

Week 8 due Nov 26, 2023 15:12 IST

8.2.4 Computing the Inverse of A via Gauss-Jordan Elimination

Summary

Gauss-Jordan elimination can be used to compute A^{-1} .

23 / 1

▶ 5:18 / 5:18

▶ 2.0x

🔊

🔗

📄

🗨

Video
📄 [Download video file](#)

Transcripts
📄 [Download SubRip \(.srt\) file](#)
📄 [Download Text \(.txt\) file](#)

Reading Assignment

0 points possible (ungraded)
Read Unit 8.2.4 of the notes. [\[LINK\]](#)

☒ Done

✔

Submit

📘 Answers are displayed within the problem

Discussion

Topic: Week 8 / 8.2.4

Hide Discussion

🧮 Calculator

Show all posts

by recent activity

There are no posts in this topic yet.

✖

Homework 8.2.4.1

45/45 points (graded)
Evaluate

• $\left(\begin{array}{ccc|ccc} 1 & 0 & 0 & -2 & 2 & -5 \\ 1 & 1 & 0 & 2 & -3 & 7 \\ -2 & 0 & 1 & -4 & 3 & -7 \end{array}\right) = \left(\begin{array}{ccc|ccc} -2 & 2 & -5 & \beta_{0,0} & \beta_{0,1} & \beta_{0,2} \\ 0 & -1 & 2 & \beta_{1,0} & \beta_{1,1} & \beta_{1,2} \\ 0 & -1 & 3 & \beta_{2,0} & \beta_{2,1} & \beta_{2,2} \end{array}\right)$

| | | | |
|--|---------------------------------------|-------------------------------------|-------------------------------------|
| $\left(\begin{array}{ccc ccc} \beta_{0,0} & \beta_{0,1} & \beta_{0,2} & 1 & 0 & 0 \\ \beta_{1,0} & \beta_{1,1} & \beta_{1,2} & 1 & 1 & 0 \\ \beta_{2,0} & \beta_{2,1} & \beta_{2,2} & -2 & 0 & 1 \end{array}\right) =$ | <div>1</div> <div>✓ Answer: 1</div> | <div>0</div> <div>✓ Answer: 0</div> | <div>0</div> <div>✓ Answer: 0</div> |
| | <div>1</div> <div>✓ Answer: 1</div> | <div>1</div> <div>✓ Answer: 1</div> | <div>0</div> <div>✓ Answer: 0</div> |
| | <div>-2</div> <div>✓ Answer: -2</div> | <div>0</div> <div>✓ Answer: 0</div> | <div>1</div> <div>✓ Answer: 1</div> |

• $\left(\begin{array}{ccc|ccc} 1 & 2 & 0 & -2 & 2 & -5 \\ 0 & 1 & 0 & 0 & -1 & 2 \\ 0 & -1 & 1 & 0 & -1 & 3 \end{array}\right) = \left(\begin{array}{ccc|ccc} -2 & 0 & -1 & \beta_{0,0} & \beta_{0,1} & \beta_{0,2} \\ 0 & -1 & 2 & \beta_{1,0} & \beta_{1,1} & \beta_{1,2} \\ 0 & 0 & 1 & \beta_{2,0} & \beta_{2,1} & \beta_{2,2} \end{array}\right)$

| | | | |
|---|---------------------------------------|---------------------------------------|-------------------------------------|
| $\left(\begin{array}{ccc ccc} \beta_{0,0} & \beta_{0,1} & \beta_{0,2} & 3 & 2 & 0 \\ \beta_{1,0} & \beta_{1,1} & \beta_{1,2} & 1 & 1 & 0 \\ \beta_{2,0} & \beta_{2,1} & \beta_{2,2} & -3 & -1 & 1 \end{array}\right) =$ | <div>3</div> <div>✓ Answer: 3</div> | <div>2</div> <div>✓ Answer: 2</div> | <div>0</div> <div>✓ Answer: 0</div> |
| | <div>1</div> <div>✓ Answer: 1</div> | <div>1</div> <div>✓ Answer: 1</div> | <div>0</div> <div>✓ Answer: 0</div> |
| | <div>-3</div> <div>✓ Answer: -3</div> | <div>-1</div> <div>✓ Answer: -1</div> | <div>1</div> <div>✓ Answer: 1</div> |

• $\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & -2 & 0 & -1 \\ 0 & 1 & -2 & 0 & -1 & 2 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array}\right) = \left(\begin{array}{ccc|ccc} -2 & 0 & 0 & \beta_{0,0} & \beta_{0,1} & \beta_{0,2} \\ 0 & -1 & 0 & \beta_{1,0} & \beta_{1,1} & \beta_{1,2} \\ 0 & 0 & 1 & \beta_{2,0} & \beta_{2,1} & \beta_{2,2} \end{array}\right)$

| | | | |
|--|-------------------------------------|-------------------------------------|---------------------------------------|
| $\left(\begin{array}{ccc ccc} \beta_{0,0} & \beta_{0,1} & \beta_{0,2} & 0 & 1 & 1 \\ \beta_{1,0} & \beta_{1,1} & \beta_{1,2} & 7 & 3 & -2 \\ \beta_{2,0} & \beta_{2,1} & \beta_{2,2} & & & \end{array}\right) =$ | <div>0</div> <div>✓ Answer: 0</div> | <div>1</div> <div>✓ Answer: 1</div> | <div>1</div> <div>✓ Answer: 1</div> |
| | <div>7</div> <div>✓ Answer: 7</div> | <div>3</div> <div>✓ Answer: 3</div> | <div>-2</div> <div>✓ Answer: -2</div> |

-3

✓ Answer: -3

-1

✓ Answer: -1

1

✓ Answer: 1

• $\begin{pmatrix} -\frac{1}{2} & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} -2 & 0 & 0 & | & 0 & 1 & 1 \\ 0 & -1 & 0 & | & 7 & 3 & -2 \\ 0 & 0 & 1 & | & -3 & -1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & | & \beta_{0,0} & \beta_{0,1} & \beta_{0,2} \\ 0 & 1 & 0 & | & \beta_{1,0} & \beta_{1,1} & \beta_{1,2} \\ 0 & 0 & 1 & | & \beta_{2,0} & \beta_{2,1} & \beta_{2,2} \end{pmatrix}$

$\begin{pmatrix} \beta_{0,0} & \beta_{0,1} & \beta_{0,2} \\ \beta_{1,0} & \beta_{1,1} & \beta_{1,2} \\ \beta_{2,0} & \beta_{2,1} & \beta_{2,2} \end{pmatrix} =$

0

✓ Answer: 0

-1/2

Answer: -1/2

✓

-1/2

Answer: -1/2

✓

-7

✓ Answer: -7

-3

Answer: -3

✓

2

Answer: 2

✓

-3

✓ Answer: -3

-1

Answer: -1

✓

1

Answer: 1

✓

• $\begin{pmatrix} -2 & 2 & -5 \\ 2 & -3 & 7 \\ -4 & 3 & -7 \end{pmatrix} \begin{pmatrix} 0 & -\frac{1}{2} & -\frac{1}{2} \\ -7 & -3 & 2 \\ -3 & -1 & 1 \end{pmatrix} = \begin{pmatrix} \beta_{0,0} & \beta_{0,1} & \beta_{0,2} \\ \beta_{1,0} & \beta_{1,1} & \beta_{1,2} \\ \beta_{2,0} & \beta_{2,1} & \beta_{2,2} \end{pmatrix}$

$\begin{pmatrix} \beta_{0,0} & \beta_{0,1} & \beta_{0,2} \\ \beta_{1,0} & \beta_{1,1} & \beta_{1,2} \\ \beta_{2,0} & \beta_{2,1} & \beta_{2,2} \end{pmatrix} =$

1

✓ Answer: 1

0

✓ Answer: 0

0

✓ Answer: 0

0

✓ Answer: 0

1

✓ Answer: 1

0

✓ Answer: 0

0

✓ Answer: 0

0

✓ Answer: 0

1

✓ Answer: 1

Submit

i Answers are displayed within the problem

Homework 8.2.4.2

1/1 point (graded)
In this exercise, you will use MATLAB to compute the inverse of a matrix using the techniques discussed in this unit.

Initialize

```
A = [
-2  2 -5
 2 -3  7
-4  3 -7 ]
```

Calculator

| | |
|---|---|
| Create an appended matrix by appending the identity | <div>A_appended = [A eye(size(A))]</div> |
| Create the first Gauss transform to introduce zeros in the first column (fill in the ?s). | <div>G0 = [1 0 0 ? 1 0 ? 0 1]</div> |
| Apply the Gauss transform to the appended system | <div>A0 = G0 * A_appended</div> |
| Create the second Gauss transform to introduce zeros in the second column | <div>G1 = [1 ? 0 0 1 0 0 ? 1]</div> |
| Apply the Gauss transform to the appended system | <div>A1 = G1 * A0</div> |
| Create the third Gauss transform to introduce zeros in the third column | <div>G2 = [1 0 ? 0 1 ? 0 0 1]</div> |
| Apply the Gauss transform to the appended system | <div>A2 = G2 * A1</div> |
| Create a diagonal matrix to set the diagonal elements to one | <div>D3 = [-1/2 0 0 0 -1 0 0 0 1]</div> |
| Apply the diagonal matrix to the appended system | <div>A3 = D3 * A2</div> |
| Extract the (updated) appended columns | <div>Ainv = A3(:, 4:6)</div> |
| Check that the inverse was computed | <div>A * Ainv</div> |

The result should be a 3×3 identity matrix.

☒ Done/Skip



Homework 8.2.4.2 Answer.m.

Submit

i Answers are displayed within the problem

Homework 8.2.4.3

18/18 points (graded)
Compute

$\begin{pmatrix} 3 & 2 & 9 \\ -3 & -3 & -14 \\ 3 & 1 & 3 \end{pmatrix}^{-1} =$

5/3

✓ Answer: 5/3

-11

✓ Answer: -11

2

✓ Answer: 2

1

✓ Answer: 1

-6

✓ Answer: -6

1

✓ Answer: 1

-1/3

✓ Answer: -1/3

5

✓ Answer: 5

-1

✓ Answer: -1

.

$\begin{pmatrix} 2 & -3 & 4 \\ 2 & -2 & 3 \\ 6 & -7 & 9 \end{pmatrix}^{-1} =$

-3/2

✓

Answer: -3/2

0

✓

Answer: 0

1

✓

Answer: 1

1/2

✓

Answer: 1/2

3

✓

Answer: 3

2

✓

Answer: 2

1/2

✓

Answer: 1/2

-1

✓

Answer: -1

-1

✓

Answer: -1

.

Submit

i Answers are displayed within the problem

Homework 8.2.4.4

1/1 point (graded)
Assume below that all matrices and vectors are partitioned “conformally” so that the operations make sense.

$$\left(\begin{array}{c|c|c} I & -u_{01} & 0 \\ \hline 0 & 1 & 0 \end{array} \right) \left(\begin{array}{c|c|c|c|c} D_{00} & a_{01} & A_{02} & & B_{00} & 0 & 0 \\ \hline 0 & \tilde{u}_{01} & \tilde{A}_{02} & & \tilde{B}_{02} & 1 & 0 \end{array} \right)$$

Calculator

$$\left(\begin{array}{c|c|c} \mathbf{0} & \mathbf{1} & \mathbf{0} \\ \hline 0 & -l_{21} & I \end{array}\right) \left(\begin{array}{c|c|c|c|c|c} \mathbf{0} & \alpha_{11} & \mathbf{a}_{12} & & \mathbf{u}_{10} & \mathbf{1} & \mathbf{0} \\ \hline 0 & \mathbf{a}_{21} & \mathbf{A}_{22} & & \mathbf{B}_{20} & 0 & I \end{array}\right)$$
$$= \left(\begin{array}{c|c|c|c|c|c|c} D_{00} & \mathbf{a}_{01} - \alpha_{11} \mathbf{u}_{01} & \mathbf{A}_{02} - \mathbf{u}_{01} \mathbf{a}_{12}^T & & \mathbf{B}_{00} - \mathbf{u}_{01} \mathbf{b}_{10}^T & -\mathbf{u}_{01} & 0 \\ \hline 0 & \alpha_{11} & \mathbf{a}_{12}^T & & \mathbf{b}_{10}^T & 1 & 0 \\ \hline 0 & \mathbf{a}_{21} - \alpha_{11} l_{21} & \mathbf{A}_{22} - l_{21} \mathbf{a}_{12}^T & & \mathbf{B}_{20} - l_{21} \mathbf{b}_{10}^T & -l_{21} & I \end{array}\right)$$

Always

✔ Answer: Always

Submit

ⓘ

Answers are displayed within the problem

Homework 8.2.4.5

1/1 point (graded)

Assume below that all matrices and vectors are partitioned “conformally” so that the operations make sense and that $\alpha_{11} \neq 0$.

Choose

- $\mathbf{u}_{01} := \mathbf{a}_{01} / \alpha_{11}$; and
 - $l_{21} := \mathbf{a}_{21} / \alpha_{11}$.

Consider the following expression:

$$\left(\begin{array}{c|c|c} I & -\mathbf{u}_{01} & 0 \\ \hline 0 & 1 & 0 \\ \hline 0 & -l_{21} & I \end{array}\right) \left(\begin{array}{c|c|c|c|c|c|c} D_{00} & \mathbf{a}_{01} & \mathbf{A}_{02} & & \mathbf{B}_{00} & 0 & 0 \\ \hline 0 & \alpha_{11} & \mathbf{a}_{12}^T & & \mathbf{b}_{10}^T & 1 & 0 \\ \hline 0 & \mathbf{a}_{21} & \mathbf{A}_{22} & & \mathbf{B}_{20} & 0 & I \end{array}\right)$$
$$= \left(\begin{array}{c|c|c|c|c|c|c} D_{00} & 0 & \mathbf{A}_{02} - \mathbf{u}_{01} \mathbf{a}_{12}^T & & \mathbf{B}_{00} - \mathbf{u}_{01} \mathbf{b}_{10}^T & -\mathbf{u}_{01} & 0 \\ \hline 0 & \alpha_{11} & \mathbf{a}_{12}^T & & \mathbf{b}_{10}^T & 1 & 0 \\ \hline 0 & 0 & \mathbf{A}_{22} - l_{21} \mathbf{a}_{12}^T & & \mathbf{B}_{20} - l_{21} \mathbf{b}_{10}^T & -l_{21} & I \end{array}\right)$$

Always

✔ Answer: Always

Submit

ⓘ

Answers are displayed within the problem

The above observations justify the following two algorithms for Gauss-Jordan elimination" for inverting a matrix.

Algorithm: $[A, B] := \text{GJ_INVERSE_PART1}(A, B)$

Partition

$A \rightarrow \left(\begin{array}{c|c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array}\right), B \rightarrow \left(\begin{array}{c|c} B_{TL} & B_{TR} \\ \hline B_{BL} & B_{BR} \end{array}\right)$

where A_{TL} is 0×0 , B_{TL} is 0×0

while $m(A_{TL}) < m(A)$ do

Repartition

$\left(\begin{array}{c|c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array}\right) \rightarrow \left(\begin{array}{c|c|c} A_{00} & \mathbf{a}_{01} & A_{02} \\ \hline \mathbf{a}_{10}^T & \alpha_{11} & \mathbf{a}_{12}^T \\ \hline A_{20} & \mathbf{a}_{21} & A_{22} \end{array}\right), \left(\begin{array}{c|c} B_{TL} & B_{TR} \\ \hline B_{BL} & B_{BR} \end{array}\right) \rightarrow \left(\begin{array}{c|c|c} B_{00} & \mathbf{b}_{01} & B_{02} \\ \hline \mathbf{b}_{10}^T & \beta_{11} & \mathbf{b}_{12}^T \\ \hline B_{20} & \mathbf{b}_{21} & B_{22} \end{array}\right)$

where α_{11} is 1×1 , β_{11} is 1×1

Calculator

| | | | |
|--------------------------------|-------------------------------------|-------------------------------------|---------------------|
| $a_{01} := a_{01}/\alpha_{11}$ | $A_{02} := A_{02} - a_{01}a_{12}^T$ | $B_{00} := B_{00} - a_{01}b_{10}^T$ | $b_{01} := -a_{01}$ |
| $a_{21} := a_{21}/\alpha_{11}$ | $A_{22} := A_{22} - a_{21}a_{12}^T$ | $B_{20} := B_{20} - a_{21}b_{10}^T$ | $b_{21} := -a_{21}$ |

(Note: a_{01} and a_{21} on the left need to be updated first.)

$a_{01} := 0$ (zero vector)

$a_{21} := 0$ (zero vector)

Continue with

$$\left(\begin{array}{c|c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array}\right) \leftarrow \left(\begin{array}{c|c|c} A_{00} & a_{01} & A_{02} \\ \hline a_{10}^T & \alpha_{11} & a_{12}^T \\ \hline A_{20} & a_{21} & A_{22} \end{array}\right), \left(\begin{array}{c|c} B_{TL} & B_{TR} \\ \hline B_{BL} & B_{BR} \end{array}\right) \leftarrow \left(\begin{array}{c|c|c} B_{00} & b_{01} & B_{02} \\ \hline b_{10}^T & \beta_{11} & b_{12}^T \\ \hline B_{20} & b_{21} & B_{22} \end{array}\right)$$

endwhile

Algorithm: $[A, B] := \text{GJ_INVERSE_PART2}(A, B)$

Partition $A \rightarrow \left(\begin{array}{c|c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array}\right), B \rightarrow \left(\begin{array}{c|c} B_{TL} & B_{TR} \\ \hline B_{BL} & B_{BR} \end{array}\right)$

where A_{TL} is 0×0 , B_{TL} is 0×0

while $m(A_{TL}) < m(A)$ **do**

Repartition

$$\left(\begin{array}{c|c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array}\right) \rightarrow \left(\begin{array}{c|c|c} A_{00} & a_{01} & A_{02} \\ \hline a_{10}^T & \alpha_{11} & a_{12}^T \\ \hline A_{20} & a_{21} & A_{22} \end{array}\right), \left(\begin{array}{c|c} B_{TL} & B_{TR} \\ \hline B_{BL} & B_{BR} \end{array}\right) \rightarrow \left(\begin{array}{c|c|c} B_{00} & b_{01} & B_{02} \\ \hline b_{10}^T & \beta_{11} & b_{12}^T \\ \hline B_{20} & b_{21} & B_{22} \end{array}\right)$$

where α_{11} is 1×1 , β_{11} is 1×1

$$b_{10}^T := b_{10}^T/\alpha_{11}$$

$$\beta_{11} := \beta_{11}/\alpha_{11}$$

$$b_{12}^T := b_{12}^T/\alpha_{11}$$

$$\alpha_{11} := 1$$

Continue with

$$\left(\begin{array}{c|c} A_{TL} & A_{TR} \\ \hline A_{BL} & A_{BR} \end{array}\right) \leftarrow \left(\begin{array}{c|c|c} A_{00} & a_{01} & A_{02} \\ \hline a_{10}^T & \alpha_{11} & a_{12}^T \\ \hline A_{20} & a_{21} & A_{22} \end{array}\right), \left(\begin{array}{c|c} B_{TL} & B_{TR} \\ \hline B_{BL} & B_{BR} \end{array}\right) \leftarrow \left(\begin{array}{c|c|c} B_{00} & b_{01} & B_{02} \\ \hline b_{10}^T & \beta_{11} & b_{12}^T \\ \hline B_{20} & b_{21} & B_{22} \end{array}\right)$$

endwhile

< Previous

Next >

© All Rights Reserved



edX

[About](#)

[Affiliates](#)

[edX for Business](#)

[Open edX](#)

[Careers](#)

[News](#)

Calculator

Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)
- [Cookie Policy](#)
- [Your Privacy Choices](#)

Connect

- [Idea Hub](#)
- [Contact Us](#)
- [Help Center](#)
- [Security](#)
- [Media Kit](#)



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)