



< Previous	 	 	 	 	Next >
-------------------------------	---	---	---	---	---------------------------

F.3.2 Final Questions 3-4

 Bookmark this page

Final Exam due Jan 5, 2024 00:12 IST Completed

F.3.2 Final Questions 3-4

Discussion

Topic: Final / F.3.2

Hide Discussion

Add a Post

Show all posts ▾

by recent activity ▾

There are no posts in this topic yet.

✕

Question 3

9/9 points (graded)

Compute the inverse of the following matrix: $A = \begin{pmatrix} 1 & 3 & 0 \\ -1 & 0 & 1 \\ 0 & -1 & -1 \end{pmatrix}$.

<div>-1/2</div> <div>✓</div> <div>Answer: -1/2</div>	<div>-3/2</div> <div>✓</div> <div>Answer: -3/2</div>	<div>-3/2</div> <div>✓</div> <div>Answer: -3/2</div>
<div>1/2</div> <div>✓</div> <div>Answer: 1/2</div>	<div>1/2</div> <div>✓</div> <div>Answer: 1/2</div>	<div>1/2</div> <div>✓</div> <div>Answer: 1/2</div>
<div>-1/2</div> <div>✓</div> <div>Answer: -1/2</div>	<div>-1/2</div> <div>✓</div> <div>Answer: -1/2</div>	<div>-3/2</div> <div>✓</div> <div>Answer: -3/2</div>

$$A^{-1} = \begin{pmatrix} -1/2 & -3/2 & -3/2 \\ 1/2 & 1/2 & 1/2 \\ -1/2 & -1/2 & -3/2 \end{pmatrix}$$

Submit

i Answers are displayed within the problem

Question 4

34/34 points (graded)

Compute the inverses of the following matrices

1. $A = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix}$.

 Calculator

0

✓ Answer: 0

0

✓ Answer: 0

1

✓ Answer: 1

0

✓ Answer: 0

1

✓ Answer: 0

0

✓ Answer: 0

0

✓ Answer: 0

0

✓ Answer: 1

1

✓ Answer: 1

0

✓ Answer: 0

0

✓ Answer: 0

0

✓ Answer: 0

0

✓ Answer: 0

0

✓ Answer: 0

1

✓ Answer: 1

0

✓ Answer: 0

$A^{-1} =$

It helps to remember that if P is a permutation matrix, then $P^{-1} = P^T$. $A^{-1} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$

2. $L_0^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 2 & 0 & 1 \end{pmatrix}$, $L_1^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix}$, $U^{-1} = \begin{pmatrix} 2 & -1 & 0 \\ 0 & -1 & -2 \\ 0 & 0 & 1 \end{pmatrix}$, and $A = L_0 L_1 U$. Then $A^{-1} =$

3

✓

Answer: 3

-1

✓

Answer: -1

1

✓

Answer: 1

-1

✓

Answer: -1

-3

✓

Answer: -3

1

✓

Answer: 1

0

✓

Answer: 0

-2

✓

Answer: -2

1

✓

Answer: 1

It helps to remember that

$$A^{-1} = (L_0 L_1 U)^{-1} = U^{-1} L_1^{-1} L_0^{-1}.$$

So,

$$A^{-1} = \underbrace{\begin{pmatrix} 2 & -1 & 0 \\ 0 & -1 & -2 \\ 0 & 0 & 1 \end{pmatrix}}_{U^{-1}} \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix}}_{L_1^{-1}} \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 2 & 0 & 1 \end{pmatrix}}_{L_0^{-1}}$$

Next, it helps to think through what happens when you multiply

$$\left(\begin{array}{c|c|c} c_0 & c_1 & c_2 \end{array} \right) \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix} = \left(\begin{array}{c|c|c} c_0 & c_1 + c_2 & c_2 \end{array} \right)$$

and

$$\begin{pmatrix} 1 & 0 & 0 \end{pmatrix}$$

$$\left(\begin{array}{c|c|c} c_0 & c_1 & c_2 \end{array} \right) \left(\begin{array}{ccc} -1 & 1 & 0 \\ 2 & 0 & 1 \end{array} \right) = \left(\begin{array}{c|c|c} c_0 - c_1 + 2c_2 & c_1 & c_2 \end{array} \right)$$

Hence

$$\begin{aligned} A^{-1} &= \underbrace{\left(\begin{array}{c|c|c} 2 & -1 & 0 \\ 0 & -1 & -2 \\ 0 & 0 & 1 \end{array} \right) \left(\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{array} \right)}_{\left(\begin{array}{c|c|c} 2 & -1+0 & 0 \\ 0 & -1-2 & -2 \\ 0 & 0+1 & 1 \end{array} \right) = \left(\begin{array}{ccc} 2 & -1 & 0 \\ 0 & -3 & -2 \\ 0 & 1 & 1 \end{array} \right)} \left(\begin{array}{ccc} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 2 & 0 & 1 \end{array} \right) \\ &= \underbrace{\left(\begin{array}{c|c|c} 2 & -1+0 & 0 \\ 0 & -1-2 & -2 \\ 0 & 0+1 & 1 \end{array} \right)}_{\left(\begin{array}{ccc} 2 & -1 & 0 \\ 0 & -3 & -2 \\ 0 & 1 & 1 \end{array} \right)} \left(\begin{array}{ccc} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 2 & 0 & 1 \end{array} \right) \\ &= \left(\begin{array}{c|c|c} 2 - (-1)(-1) + (2)(0) & -1 & 0 \\ 0 + (-1)(-3) + (2)(-2) & -3 & -2 \\ 0 + (-1)(1) + (2)(1) & 1 & 1 \end{array} \right) = \left(\begin{array}{ccc} 3 & -1 & 0 \\ -1 & -3 & -2 \\ 1 & 1 & 1 \end{array} \right) \end{aligned}$$

3. $D = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 5 & 2 \\ 0 & 2 & 1 \end{pmatrix}$. Then $D^{-1} =$

1

✓

Answer: 1

0

✓

Answer: 0

0

✓

Answer: 0

0

✓

Answer: 0

1

✓

Answer: 1

-2

✓

Answer: -2

0

✓

Answer: 0

-2

✓

Answer: -2

5

✓

Answer: 5

(Hint: How is the inverse of matrix $\begin{pmatrix} 1 & 0 \\ 0 & B \end{pmatrix}$ related to B^{-1} ?)

$$\begin{pmatrix} 1 & 0 \\ 0 & B \end{pmatrix}^{-1} = \begin{pmatrix} 1 & 0 \\ 0 & B^{-1} \end{pmatrix}$$

Now,

$$\begin{aligned} B^{-1} &= \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}^{-1} = \frac{1}{(5)(1) - (2)(2)} \begin{pmatrix} 1 & -2 \\ -2 & 5 \end{pmatrix} \\ &= \frac{1}{1} \begin{pmatrix} 1 & -2 \\ -2 & 5 \end{pmatrix} = \begin{pmatrix} 1 & -2 \\ -2 & 5 \end{pmatrix}. \end{aligned}$$

Hence

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 5 & 2 \\ 0 & 2 & 1 \end{pmatrix}^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -2 \\ 0 & -2 & 5 \end{pmatrix}$$

Submit

< Previous

Next >

© All Rights Reserved



edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)
- [Cookie Policy](#)
- [Your Privacy Choices](#)

Connect

- [Idea Hub](#)
- [Contact Us](#)
- [Help Center](#)
- [Security](#)
- [Media Kit](#)



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)