



[Course](#) > [Unit 1: Fourier Series](#) > [1. Introduction to Fourier Series](#) > 2. Motivation


**Audit Access Expires Jun 24, 2020**

You lose all access to this course, including your progress, on Jun 24, 2020.

Upgrade by Jun 7, 2020 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

## 2. Motivation

### Why Fourier Series?



▶ 0:00 / 8:52

▶ 1.50x 🔊 🗑️ 📄 🗨️

## Video

[Download video file](#)

## Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

The exponential response formula (ERF) is a quick method for finding the response to any input that is expressible in terms of an exponential, sine, or cosine function. (We will review this method on the next page.) What happens when the input is not one of these?



If the input function  $f(t)$  is periodic (of period  $2\pi$ ), we can express the function (where it is continuous) as an infinite sum of sines and cosines. This series representation is called a **Fourier series** :

$$f(t) = c_0 + \sum_{n=1}^{\infty} [a_n \cos(nt) + b_n \sin(nt)], \quad c_0, a_n, b_n \text{ real constants.}$$

(Note that here we are giving an expression that only works for  $2\pi$ -periodic functions. There is a formula that works more generally, which we will learn in the next lecture.)

The benefit is, given an inhomogeneous linear differential equation  $P(D)y = f(t)$ , we can apply ERF to every term in the Fourier series.

Input		Response
$b_n \sin(nt)$	$\rightsquigarrow$	$b_n y_n^{(s)}(t)$
$a_n \cos(nt)$	$\rightsquigarrow$	$a_n y_n^{(c)}(t)$
$c_0$	$\rightsquigarrow$	$c_1$

Taking the sum of all of the input terms we get an approximation to the input  $f(t)$ . Therefore by superposition (since our differential equation is linear), the sum of the terms on the right hand side is a series representation of the system response.

Input		Response
$f(t)$	$\rightsquigarrow$	$c_1 + \sum_{n=1}^{\infty} a_n y_n^{(c)}(t) + b_n y_n^{(s)}(t)$

### Program for solving inhomogeneous DEs with periodic input

1. Find a Fourier series approximation to the input signal (*What we will learn today.*)
2. The system response is the sum of the response to each  $\sin nt$  and  $\cos nt$  input. (*Explore in 2 lectures from now.*)


**Question:** Is it really useful to have a particular solution that is expressed as an infinite sum?



**Answer:** It is useful when you can determine which terms in the sum are dominant.

But first things first: we must define the Fourier series, determine how to compute it. In order to define the Fourier series, we start by reviewing the components necessary to define them – periodic functions.

## Overview of lecture goals



(Caption will be displayed when you start playing the video.)

0:00 / 0:00

1.50x



## Video

[Download video file](#)

## Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

## 2. Motivation

**Topic:** Unit 1: Fourier Series / 2. Motivation

Hide Discussion

Add a Post

Show all posts ▼

by recent activity ▼

? [STAFF: Download lecture videos](#)

I have problems downloading almost all of the lecture videos in the Unit 1. Could you help make it simpler to download these videos. Much appreciated... Thank you...

4

? [I cant see the equations](#)

11

Learn About Verified Certificates

© All Rights Reserved

