

## MITx: 6.041x Introduction to Probability - The Science of Uncertainty

■ Bookmarks

- Unit 0: Overview
- Entrance Survey
- Unit 1: Probability models and axioms
- Unit 2: Conditioning and independence
- Unit 3: Counting
- Unit 4: Discrete random variables
- Exam 1
- Unit 5: Continuous random variables

Unit 8: Limit theorems and classical statistics > Lec. 20: An introduction to classical statistics > Lec 20 An introduction to classical statistics vertical6

**■** Bookmark

## Exercise: ML estimation

(1/1 point)

Let K be a Poisson random variable with parameter  $\lambda$ : its PMF is

$$p_K(k;\lambda) = rac{\lambda^k e^{-\lambda}}{k!}, \qquad ext{for } k=0,1,2,\ldots.$$

What is the ML estimate of  $\lambda$  based on a single observation K=k? (Your answer should be an algebraic function of k using standard notation .)

k

**V** 

Answer: k

Answer:

- Unit 6: Further topics on random variables
- Unit 7: Bayesian inference
- ▶ Exam 2
- ▼ Unit 8: Limit theorems and classical statistics

Unit overview

Lec. 18: Inequalities, convergence, and the Weak Law of Large Numbers

Exercises 18 due Apr 27, 2016 at 23:59 UTC

Lec. 19: The Central Limit Theorem (CLT)

Exercises 19 due Apr 27, 2016 at 23:59 UTC

Lec. 20: An introduction to classical statistics

Exercises 20 due Apr 27, 2016 at 23:59 UTC

We maximize the logarithm of the PMF, which is  $k\ln\lambda-\lambda-\ln(k!)$ . Setting the derivative of this expression with respect to  $\lambda$  to 0, we obtain  $(k/\lambda)-1=0$ , so that  $\hat{\lambda}_{ML}=k$ .

You have used 1 of 2 submissions

Solved problems

Additional theoretical material

**Problem Set 8** 

Problem Set 8 due Apr 27, 2016 at 23:59 UTC

**Unit summary** 

© All Rights Reserved



© edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

















