

Ţ <u>Help</u>

sandipan_dey ~

Next >

<u>Syllabus</u> laff routines **Community Discussion** <u>Outline</u> <u>Course</u> <u>Progress</u> <u>Dates</u>

☆ Course / Week 1: Vectors in Linear Algebra / 1.3 Simple Vector Operations

(1)

1.3.4 Vector Subtraction


□ Bookmark this page

Previous

■ Calculator

Week 1 due Oct 5, 2023 03:12 IST Completed

1.3.4 Vector Subtraction



Start of transcript. Skip to the end.

Dr. Robert van de Geijn: Addition is important.

Scaling is important.

And the next question, of course, becomes how do you

subtract two vectors?

We're going to see that that's just a matter of combining

▶ 0:00 / 0:00

X CC 66 ▶ 2.0x

Video

▲ Download video file

Transcripts

- **▲** Download Text (.txt) file

Reading Assignment

0 points possible (ungraded) Read Unit 1.3.4 of the notes. [LINK]



Done



Submit

Discussion

Topic: Week 1 / 1.3.4

Hide Discussion

by recent activity >

Add a Post

Show all posts

Assignment not show as in completed in overall course even after all assignments submitted

Is anyone else having this problem? In this a several other sections, even after the assignments are done, submitted and the page itself shows a...

Homework 1.3.4.1

1/1 point (graded)

For $x \in \mathbb{R}^n, x-x=0$

⊞ Calculator

	✓ Answer: Always	
xplanation		
anscripted in final	section of this week	
<u> DF of Answer in Vic</u>		
Submit		
Submit		
	splayed within the problem	
	splayed within the problem	
• Answers are disonmework 1.3.4	4.2	
• Answers are disonmework 1.3.4	4.2	
Answers are discomework 1.3.4 point (graded) or $x,y\in\mathbb{R}^n$, x	4.2	
Answers are discomework 1.3.4 point (graded) or $x,y\in\mathbb{R}^n$, x — ometimes	4.2 $y = y - x$. Answer: Sometimes	a same but the direction is apposite since $m - u = (-1) \times (u - m)$
Answers are discomework 1.3.4 point (graded) or $x,y \in \mathbb{R}^n$, x ometimes	4.2 $y = y - x$. Answer: Sometimes is to notice that the length is the	e same, but the direction is opposite, since $x-y=(-1) imes (y-x)$
Answers are discomework 1.3.4 point (graded) or $x,y\in\mathbb{R}^n$, x ometimes	4.2 $y = y - x$. Answer: Sometimes is to notice that the length is the	e same, but the direction is opposite, since $x-y=(-1) imes (y-x)$
Answers are discomework 1.3.4 point (graded) or $x,y \in \mathbb{R}^n$, x ometimes	4.2 $y = y - x$. Answer: Sometimes is to notice that the length is the	e same, but the direction is opposite, since $x-y=(-1) imes (y-x)$

© All Rights Reserved



edX


<u>About</u>

Affiliates

edX for Business

<u>Open edX</u>

Careers



<u>News</u>

Legal

Terms of Service & Honor Code

Privacy Policy

Accessibility Policy

Trademark Policy

<u>Sitemap</u>

Cookie Policy

Your Privacy Choices

Connect

<u>Idea Hub</u>

Contact Us

Help Center

<u>Security</u>

Media Kit















© 2023 edX LLC. All rights reserved.

深圳市恒宇博科技有限公司 <u>粤ICP备17044299号-2</u>