**Data Analysis: Statistical Modeling and Computation in Applications** 

<u>Help</u>

sandipan\_dey ~

<u>Course</u>

**Progress** 

<u>Dates</u>

**Discussion** 

Resources






Next >



☐ Bookmark this page

Previous

## Fitting regressions to timeseries



Start of transcript. Skip to the end.

Prof Jegelka: Now that you've seen a few statistical models--

and these were actually all linear models--

let's think about how we could start fitting any

of these models to our data.

So we'll first take a regression

**▶** 0:00 / 0:00

▶ 2.0x

×

cc 66

Video

Download video file

## **Transcripts**

<u>Download SubRip (.srt) file</u> <u>Download Text (.txt) file</u>

### Discussion

Preview

**Topic:** Module 4: Time Series: Statistical Models / 8. Fitting regressions to timeseries

Add a Post

**Hide Discussion** 

# ✓ All Posts If we have stationarity, shouldn't the homogeneity in variance be guaranteed? question posted 2 months ago by Syed SB title This post is visible to everyone. vilgalys (Staff) 2 months ago - marked as answer 2 months ago by Syed SB Yes, that's correct! Homogeneity in variance is a component of stationarity. I believe what Prof. Jegelka is saying is that this is one of the prerequisites for running a time series regression. Add a comment

© All Rights Reserved



# edX

**About** 

**Affiliates** 

edX for Business

Open edX

Careers

**News** 

# Legal

Terms of Service & Honor Code

Privacy Policy

**Accessibility Policy** 

**Trademark Policy** 

<u>Sitemap</u>

# **Connect**

**Blog** 

Contact Us

Help Center

Media Kit

**Donate** 















© 2021 edX Inc. All rights reserved.

深圳市恒宇博科技有限公司 <u>粤ICP备17044299号-2</u>