



[Course](#) > [Unit 1: Fourier Series](#) > [2. Properties of Fourier Series \(of Period 2L\)](#)

> 11. Antiderivative of a Fourier series

Audit Access Expires Jun 24, 2020

You lose all access to this course, including your progress, on Jun 24, 2020.

Upgrade by Jun 7, 2020 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)


11. Antiderivative of a Fourier series

Suppose that f is a piecewise differentiable periodic function, and that F is an antiderivative of f . (If f has jump discontinuities, one can still define $F(t) := \int_0^t f(\tau) d\tau + C$, but at the jump discontinuities F will be only continuous, not differentiable.)

The function F might not be periodic. For example, if f is a function of period 2 such that

$$f(t) := \begin{cases} 2, & \text{if } 0 < t < 1, \\ -1 & \text{if } -1 < t < 0, \end{cases}$$

then $F(t)$ creeps upward over time.



An even easier example: if $f(t) = 1$, then $F(t) = t + C$ for some C , so $F(t)$ is not periodic.

But if the constant term $a_0/2$ in the Fourier series of f is 0, then F is periodic, and its Fourier series can be obtained by taking the simplest antiderivative of each cosine and sine term, and adding an overall $+C$, where C is the average value of F .

Problem 11.1 Let $T(t)$ be the triangle wave of period 2 and amplitude 1: so that $T(t) = |t|$ for $-1 \leq t \leq 1$. Find the Fourier series of $T(t)$.

Solution: We could use the Fourier coefficient formula. But instead, notice that $T(t)$ has slope -1 on $(-1, 0)$ and slope 1 on $(0, 1)$, so $T(t)$ is an antiderivative of the period 2 square wave

$$\text{Sq}(\pi t) = \sum_{n \geq 1, \text{odd}} \frac{4}{n\pi} \sin n\pi t.$$

Taking an antiderivative termwise (and using that the average value of $T(t)$ is $1/2$) gives

$$\begin{aligned} T(t) &= \frac{1}{2} + \sum_{n \geq 1, \text{odd}} \frac{4}{n\pi} \left(\frac{-\cos n\pi t}{n\pi} \right) \\ &= \frac{1}{2} - \sum_{n \geq 1, \text{odd}} \frac{4}{n^2 \pi^2} \cos n\pi t. \end{aligned}$$

Warning: If a periodic function $f(t)$ is not continuous, it will not be an antiderivative of any piecewise differentiable function, so you cannot find the Fourier series of $f(t)$ by integration.

Remark 11.2 The Fourier series for a function with discontinuities can (formally) be differentiated term by term, but **the result will not converge**. For example, the term-wise derivative of the Fourier series for $Sq(t)$ is

$$\frac{4}{\pi} \sum_{n \text{ odd}} \cos(nt).$$

This does not converge anywhere (since the n th term does not even vanish as $n \rightarrow \infty$). However, note that it is possible to make sense of this series, and of the anti-derivative of the square wave function, in terms of Dirac's delta functions and the theory of distributions — seen in more advanced courses than this one.

Integrate to find the Fourier series (*)

2/2 points (graded)

Find the Fourier series of the function $f(t) = t^2$ defined on $[-1, 1]$.

First find the constant term.

$$\frac{a_0}{2} = \boxed{\frac{1}{3}} \quad \checkmark \text{ Answer: } 1/3$$

Next, find the remaining terms of the Fourier series in terms of n :

$$\frac{a_0}{2} + \sum_{n=1}^{\infty} \boxed{4*(-1)^n/(n^2*\pi^2)*\cos(n*\pi*t)} \quad \checkmark \text{ Answer: } 4*(-1)^n*\cos(n*\pi*t)/(n^2*\pi^2)$$

Solution:

The function $f(t) = t^2$, $-1 < t < 1$ is even of period 2. This function is continuous, so it could be the antiderivative of another function. In particular, note that $f(t)$ is the antiderivative of the period 2 sawtooth wave $g(t) = 2t$, $-1 < t < 1$.



The Fourier series for the 2π -periodic sawtooth wave $W(u)$ is

$$W(u) = 2 \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \sin(nu).$$

Therefore, scaling by $u = \pi t$ so that when $u = \pi$, we have $t = 1$ we get that

$$g(t) = \frac{2}{\pi} W(u) = \frac{2}{\pi} W(\pi t) = \frac{4}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \sin(n\pi t).$$

Finally, to find the Fourier series for $f(t)$, we integrate the Fourier series for $g(t)$ term-by-term.

$$\begin{aligned} \int g(t) dt &= \frac{4}{\pi} \sum_{n=1}^{\infty} \int \frac{(-1)^{n+1}}{n} \sin(n\pi t) dt \\ &= \frac{4}{\pi} \sum_{n=1}^{\infty} \frac{-(-1)^{n+1}}{n^2 \pi} \cos(n\pi t) \\ &= \frac{4}{\pi^2} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} \cos(n\pi t). \end{aligned}$$

To find the constant term, we find the average value of $f(t)$ on the interval $-1 < t < 1$.

$$\frac{a_0}{2} = \frac{\int_{-1}^1 t^2 dt}{1 - (-1)} = \frac{1}{2} \left(\frac{1 - (-1)}{3} \right) = \frac{1}{3}.$$

Submit

You have used 3 of 10 attempts



i Answers are displayed within the problem

11. Antiderivative of a Fourier series

Hide Discussion

Topic: Unit 1: Fourier Series / 11. Antiderivative of a Fourier series

Add a Post

◀ All Posts

Improvement s to the grader?

discussion posted 24 days ago by [jsimko](#)

Just noticed some improvements with the grader. Looks like partial credit is now given for a mistake with sign when the rest of the answer is correct.

Nice!

This post is visible to everyone.

Add a Response

1 response

jfrench (Staff)

23 days ago

:) You speak, we listen. Thanks for the feedback. I realized it could be confusing and was too strict in interpretation.

so, we have to keep the half point and continue. right?

posted 19 days ago by [vcassati](#)

You can keep submitting answers until you run out of attempts or get all of the points (whichever comes first). It looks like 10 attempts are allowed for the problem on this page.



posted 19 days ago by [Steve Nicodemus](#)

Add a comment

Showing all responses

Add a response:

Preview

Submit

[Learn About Verified Certificates](#)

© All Rights Reserved

