

<u>Help</u>

sandipan\_dey >

Next >

**Discussion Progress** <u>Dates</u> <u>Calendar</u> <u>Notes</u> <u>Course</u>

☆ Course / Unit 3: Optimization / Lecture 11: Lagrange Multipliers



You are taking "Exam (Timed, No Correctness Feedback)" as a timed exam. Show more



22:47:55






□ Bookmark this page



#### **Summarize**

### **Summary of argument**



Start of transcript. Skip to the end.

PROFESSOR: So somebody made a comment to me

recently that in learning this stuff, something that's tricky

. . . . . .

So you learn some new stuff, and then there's an argument,

and there's a bunch of little steps in that argument.

0:00 / 0:00 " 2.0x X CC


#### Video

Download video file

### **Transcripts**

Download SubRip (.srt) file Download Text (.txt) file

We have been exploring a function  $f\left(x,y
ight)$  constrained to a region R. The gradient of the function  $f\left(x,y
ight)$  and the region R, a circle of radius 1 centered at the point (2,0) are pictured below.



We have defined the function  $g\left(x,y
ight)=\left(x-2
ight)^{2}+y^{2}$  , whose level curve  $g\left(x,y
ight)=1$  defines the boundary of the region  $oldsymbol{R}$ .

What we've discovered so far:

- ullet abla g is perpendicular to the boundary of R everywhere (because we've defined g so that its level curve is this boundary)
- ullet abla f is perpendicular to the boundary of R at the maximum point
- At the maximum point of  $m{f}$  on this region  $m{R}$ ,  $m{
  abla} m{f}$  is pointing in the same direction (o

■ Calculator

This holds more generally.

**Theorem** The maximum (or minimum) of a function  $f\left(x,y
ight)$  constrained to a level curve  $g\left( x,y\right) =c$  occurs where the gradient of the function is parallel to the gradient of the constraint equation:

$$abla f = \lambda 
abla g$$

for a real number  $\lambda$ .


**Pro tip:** You must enable 3rd party cookies so that you can interact with the mathlet below.

Change functions and constraint equations using the dropdown menus, and observe the gradient vector of the function whose level curves are shown and the gradient vector of the constraint curve at the same point.

### ► Lagrange Multipliers **4**

Equation 1 
$$\hspace{0.1cm} extstyle \hspace{0.1cm} \hspace{0.1cm} z=f(x,y)=rac{x^2+y^2}{4}$$

Constraint 1 
$$\hspace{0.1cm} extstyle
otag \hspace{0.1cm} extstyle
otag \hspace{0.1cm$$





#### PLEASE RATE THIS MATHLET

(Use a one star to five star rating scale.)

- $\circ \Leftrightarrow$
- 0 \$ \$

**■** Calculator





© All Rights Reserved



# edX

**About** 

**Affiliates** 

edX for Business

Open edX

**Careers** 

<u>News</u>

# Legal

Terms of Service & Honor Code

**Privacy Policy** 

**Accessibility Policy** 

**Trademark Policy** 

<u>Sitemap</u>

## **Connect**



<u>Blog</u>

Contact Us

Help Center

Media Kit

**Donate** 















© 2021 edX Inc. All rights reserved.

深圳市恒宇博科技有限公司 <u>粤ICP备17044299号-2</u>