

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).




[Course](#) > [Unit 3 Neural networks \(2.5 weeks\)](#) > [Homework 4](#) > 3. Backpropagation

3. Backpropagation

Extension Note: Homework 4 due date has been extended by 1 day to **July 27 23:59UTC**.

One of the key steps for training multi-layer neural networks is stochastic gradient descent. We will use the back-propagation algorithm to compute the gradient of the loss function with respect to the model parameters.

Consider the L -layer neural network below:



In the following problems, we will use the following notation: b_j^l is the bias of the j^{th} neuron in the l^{th} layer, a_j^l is the activation of j^{th} neuron in the l^{th} layer, and w_{jk}^l is the weight for the connection from the k^{th} neuron in the $(l-1)^{\text{th}}$ layer to the j^{th} neuron in the l^{th} layer.

If the activation function is f and the loss function we are minimizing is C , then the equations describing the network are:

$$a_j^l = f \left(\sum_k w_{jk}^l a_k^{l-1} + b_j^l \right)$$

$$\text{Loss} = C(a^L)$$

For $l = 1, \dots, L$.

Computing the Error

2/2 points (graded)

Let the weighted inputs to the d neurons in layer l be defined as $z^l \equiv w^l a^{l-1} + b^l$, where $z^l \in \mathbb{R}^d$. As a result, we can also write the activation of layer l as $a^l \equiv f(z^l)$, and the "error" of neuron j in layer l as $\delta_j^l \equiv \frac{\partial C}{\partial z_j^l}$. Let $\delta^l \in \mathbb{R}^d$ denote the full vector of errors associated with layer l .

Back-propagation will give us a way of computing δ^l for every layer.

Assume there are d outputs from the last layer (i.e. $a^L \in \mathbb{R}^d$). What is δ_j^L for the last layer?

☒ $\frac{\partial C}{\partial a_j^L} f'(z_j^L)$ ✓

☐ $\sum_{k=1}^d \frac{\partial C}{\partial a_k^L} f'(z_j^L)$

☐ $\frac{\partial C}{\partial a_j^L}$

☐ $f'(z_j^L)$

What is δ_j^l for all $l \neq L$?

☒ $\sum_k w_{kj}^{l+1} \delta_k^{l+1} f'(z_j^l) \checkmark$

☐ $\delta_k^{l+1} f'(z_j^l)$

☐ $\sum_k w_{jk}^{l-1} \delta_j^{l-1} f'(z_j^l)$

☐ $\sum_k w_{kj}^{l+1} \delta_k^{l+1} f(z_j^l)$

Solution:

We make use of the chain rule.

1. By definition, $\delta_j^L = \frac{\partial C}{\partial a_j^L} \frac{\partial a_j^L}{\partial z_j^L} = \frac{\partial C}{\partial a_j^L} f'(z_j^L)$.

2. We have:

$$\begin{aligned} \delta_j^l &= \frac{\partial C}{\partial z_j^l} \\ &= \sum_k \frac{\partial C}{\partial z_k^{l+1}} \frac{\partial z_k^{l+1}}{\partial z_j^l} \end{aligned}$$

$$= \sum_k \frac{\partial z_k^{l+1}}{\partial z_j^l} \delta_k^{l+1}$$

Then we have $z_k^{l+1} = \sum_j w_{kj}^{l+1} a_j^l + b_k^{l+1} = \sum_j w_{kj}^{l+1} f(z_j^l) + b_k^{l+1}$. Taking the derivative of this with respect to z_j^l gives $w_{kj}^{l+1} f'(z_j^l)$.

Combining the two gives the final answer: $\delta_j^l = \sum_k w_{kj}^{l+1} \delta_k^{l+1} f'(z_j^l)$.

Submit

You have used 2 of 2 attempts

i Answers are displayed within the problem

Parameter Derivatives

2/2 points (graded)

During SGD we are interested in relating the errors computed by back-propagation to the quantities of real interest: the partial derivatives of the loss with respect to our parameters. Here that is $\frac{\partial C}{\partial w_{jk}^l}$ and $\frac{\partial C}{\partial b_j^l}$.

What is $\frac{\partial C}{\partial w_{jk}^l}$? Write in terms of the variables a_k^{l-1} , w_j^l , b_j^l , and δ_j^l if necessary.

Example of writing superscripts and subscripts:

δ_j^l for δ_j^l

$w_{jk}^{(l)}$ for $w_{jk}^{(l)}$

$$\frac{\partial C}{\partial w_{jk}^{(l)}} = \delta_j^{(l)} a_k^{(l-1)}$$

✓ Answer: $a_k^{(l-1)} \delta_j^{(l)}$

$$\delta_j^{(l)} \cdot a_k^{(l-1)}$$

What is $\frac{\partial C}{\partial b_j^{(l)}}$? Write in terms of the variables $a_k^{(l-1)}$, w_j^l , b_j^l , and δ_j^l if necessary.

$$\frac{\partial C}{\partial b_j^{(l)}} = \delta_j^{(l)}$$

✓ Answer: $\delta_j^{(l)}$

$$\delta_j^{(l)}$$

STANDARD NOTATION

Solution:

$$1. \frac{\partial C}{\partial w_{jk}^{(l)}} = \frac{\partial C}{\partial z_j^{(l)}} \frac{\partial z_j^{(l)}}{\partial w_{jk}^{(l)}} = a_k^{(l-1)} \delta_j^{(l)}$$

$$2. \frac{\partial C}{\partial b_j^{(l)}} = \frac{\partial C}{\partial z_j^{(l)}} \frac{\partial z_j^{(l)}}{\partial b_j^{(l)}} = 1 * \delta_j^{(l)}$$

Submit

You have used 1 of 5 attempts

i Answers are displayed within the problem

Activation Functions: Sigmoid

4/4 points (graded)

Recall that there are several different possible choices of activation functions f . Let's get more familiar with them and their gradients.

What is the derivative of the sigmoid function, $\sigma(z) = \frac{1}{1+e^{-z}}$? Please write your answer in terms of e and z :

$$e^{-z}/(1+e^{-z})^2$$

✓ Answer: $e^{-z} / (1 + e^{-z})^2$

$$\frac{e^{-z}}{(1+e^{-z})^2}$$

Which of the following is true of $\sigma'(z)$ as $\|z\|$ gets large?

☐ Its magnitude becomes large.

☒ Its magnitude becomes small. ✓

☐ It suffers from high variance.

What is the derivative of the ReLU function, $\text{ReLU}(z) = \max(0, z)$ for $z > 0$?

✓ Answer: 1

For $z < 0$?

✓ Answer: 0

STANDARD NOTATION**Solution:**

$\sigma'(z) = \sigma(z)(1 - \sigma(z))$. As z gets large in magnitude, the sigmoid function saturates, and the gradient approaches zero.

ReLU is a simple activation function. Above zero, it has a constant gradient of 1. Below zero, it is always zero.

You have used 1 of 5 attempts

i Answers are displayed within the problem

Simple Network

4/4 points (graded)

Consider a simple 2-layer neural network with a single neuron in each layer. The loss function is the quadratic loss:

$C = \frac{1}{2}(y - t)^2$, where y is the prediction and t is the target.

Starting with input x we have:

- $z_1 = w_1 x$
- $a_1 = \text{ReLU}(z_1)$
- $z_2 = w_2 a_1 + b$
- $y = \sigma(z_2)$
- $C = \frac{1}{2}(y - t)^2$

Consider a target value $t = 1$ and input value $x = 3$. The weights and bias are $w_1 = 0.01$, $w_2 = -5$, and $b = -1$.

Please provide numerical answers accurate to at least three decimal places.

What is the loss?

0.28842841648243966

✓ Answer: 0.28842841648243966

What are the derivatives with respect to the parameters?

$\frac{\partial C}{\partial w_1} =$ 2.0809165621704553

✓ Answer: 2.0809165621704553

$$\frac{\partial C}{\partial w_2} =$$

✓ Answer: -0.00416183312434091

$$\frac{\partial C}{\partial b} =$$

✓ Answer: -0.13872777081136367

STANDARD NOTATION

Solution:

Using the chain rule, we have:

- $\frac{\partial C}{\partial w_1} = \frac{\partial C}{\partial y} \frac{\partial y}{\partial z_2} \frac{\partial z_2}{\partial a_1} \frac{\partial a_1}{\partial z_1} \frac{\partial z_1}{\partial w_1} = (y - t) y (1 - y) w_2 \mathbf{1}\{z_1 > 0\} x$
- $\frac{\partial C}{\partial w_2} = \frac{\partial C}{\partial y} \frac{\partial y}{\partial z_2} \frac{\partial z_2}{\partial w_2} = (y - t) y (1 - y) a_1$
- $\frac{\partial C}{\partial b} = (y - t) y (1 - y)$

Submit

You have used 1 of 5 attempts

i Answers are displayed within the problem

SGD

1/1 point (graded)

Referring to the previous problem, what is the update rule for w_1 in the SGD algorithm with step size η ? Write in terms of w_1 , η , and $\frac{\partial C}{\partial w_1}$; enter the latter as (partialC)/(partialw_1), noting the lack of space in the variable names:

Next $w_1 =$ ✓ Answer: $w_1 - \eta * (\text{partialC})/(\text{partial}w_1)$

STANDARD NOTATION

Solution:

The definition of the simple SGD update rule is $\text{new_parameter} = \text{old_parameter} - \text{learning_rate} * \text{derivative of loss w.r.t old parameter}$.

Submit

You have used 1 of 5 attempts

i Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 3 Neural networks (2.5 weeks):Homework 4 / 3. Backpropagation

Add a Post

Show all posts ▼

by recent activity ▼

? [Simple Network, Derivative](#)

1

✓ [Staff]Coarse grader?	4
💬 Simple Network I'm taking the derivatives w.r.t each parameter, and the expressions are very long. Are we perhaps supposed to plug-in the values we're given, or...	10
💬 Simple Network any clues? I'm at the third try. I got the Loss and $\partial C/\partial w_2$ but didn't figure out $\partial C/\partial w_1$ and $\partial C/\partial b$	8
? [STAFF] Parameter Derivatives	6
💬 [staff] Simple network, why is my second answer wrong? All others are correct? Is this a notation problem again? Why is my second answer wrong? I calculated several times. I want to know if this is a typo or a calculation mistake.	4
? computing the error part 2 reset	3
💬 Can the course follow the same notation convention please? May I suggest a little improvement to the course? As stated in the title, please state a set of notation convention at the beginning of the course, a...	3
? [Simple Network]: To staff, can you look into my entered answer since it takes the derivatives with respect to w_1 , w_2 and b respectively but the system states that there is w_1 and w_2 , b are invalid input	2
💬 Learning Source - Hint for those who still struggle to understand our course material, watch this video. Took some time to understand the mechanics behind by this vid...	8
💬 Backpropagation -- a little history, and a cautionary tale Backpropagation is one of those ideas like Bayes Rule that's "whoa, why didn't I think of that" ...*after* you see it. It's just the plain old ordinary c... 👤 Community TA	4
💬 What the $\sigma(x)$ function in $y=\sigma(z_2)$? What the $\sigma(x)$ function in $y=\sigma(z_2)$?	3
[Staff] Online Derivatives	

☒ Is there any objection to us looking up the requested derivatives online? i.e. ReLU and Sigmoid. Additional note: Well, I just looked them up online...

3

Simple Network - puzzled with rejection of my submission

© All Rights Reserved