

< Previous

✓

✓

✓

✓

Next >

4.2.5 Problem Set: Robertson's Autocatalytic Reaction

 Bookmark this page

Now we will consider the use of an implicit method, specifically Backward Euler, to solve an initial value problem in which \underline{f} depends nonlinearly on \underline{u} . The following IVP is a model of an autocatalytic chemical reaction:

$$\frac{dY_0}{dt} = -k_0 Y_0 + k_2 Y_1 Y_2 \tag{4.37}$$

$$\frac{dY_1}{dt} = k_0 Y_0 - k_1 Y_1^2 - k_2 Y_1 Y_2 \tag{4.38}$$

$$\frac{dY_2}{dt} = k_1 Y_1^2 \tag{4.39}$$

where Y_i is the mass fraction of species i and k_i are constants. This IVP was first presented by Robertson in 1966 as an example of a stiff system and hence is known now as Robertson's problem. We will use the following values,

$$k_0 = 4.0\text{E-}2\text{ s}^{-1}, \quad k_1 = 3.0\text{E}7\text{ s}^{-1}, \quad k_2 = 1.0\text{E}4\text{ s}^{-1} \tag{4.40}$$

Following our standard IVP notation, the state vector we choose as,

$$\underline{u} = [Y_0, Y_1, Y_2]^T \tag{4.41}$$

For the initial condition at $t = 0$, we will use,

$$\underline{u}(0) = [1.0, 0.0, 0.0]^T \tag{4.42}$$

Thus, only Y_0 is present initially.

Complete the following steps:

1. Implement the `calc_all` method in `solve_robertson.py` to calculate \underline{f} and $\partial \underline{f} / \partial \underline{u}$ given \underline{u} . See the docstring for details.

Further, complete the implementation of `test_calc_all` which will test that your implementation of `calc_all` is correct. Specifically, within `test_calc_all`, call `calc_all` to determine \underline{f} and $\partial \underline{f} / \partial \underline{u}$ for $\underline{u} = [0.5, 0.05, 0.45]^T$ (and use the same values of k_0 , k_1 , and k_2 given above). The correct values of `f` and `f_u` are already included in `test_calc_all` and are named `f_correct` and `f_u_correct`.

Discussions

All posts sorted by recent activity

The main block of `solve_robertson.py` calls `test_calc_all`. When, `test_calc_all` is working, uncomment the next line in the main block beginning `time_FE = ...`, in preparation for the next part.

2. Implement the `plot_Y` method in `solve_robertson.py` to produce plots of the

< Previous

Next >

`solve_robertson.py`, the desired plot of the Forward Euler method (using $\Delta t = 1.0\text{E-}4$



edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)
- [Cookie Policy](#)
- [Your Privacy Choices](#)

Connect

- [Idea Hub](#)
- [Contact Us](#)
- [Help Center](#)
- [Security](#)
- [Media Kit](#)



`myNewton` in `mynonlin solver.py`. A tester is provided which can be checked by running

