

[Course](#)

[Progress](#)

[Dates](#)

[Discussion](#)

[Syllabus](#)

[Outline](#)

[laff routines](#)

[Community](#)

[🏠](#) [Course](#) / [Week 12: Eigenvalues and Eigenvectors](#) / [12.2 Getting Started](#)



< Previous	 	 	 	 		Next >
------------	---	---	---	---	---	--------

12.2.4 Eigenvalues and Vectors of 3×3 Matrices

 Bookmark this page

Week 12 due Dec 29, 2023 10:42 IST Completed

12.2.4 Eigenvalues and Vectors of 3 x 3 Matrices

Video 12.2.4 Part 1

Start of transcript. Skip to the end.

▶ 0:00 / 0:00

▶ 2.0x

🔊

⌂

CC

“

Dr. Robert van de Geijn: So let's move right along

and look at how to find the eigenvalues and vectors of a 3 by 3 matrix.

Let's review a little bit.

Here we have a 3 by 3 diagonal matrix.

How do you find its eigenvalues and eigenvectors?

Go and do the homework, and see me in

Video

📄 [Download video file](#)

Transcripts

📄 [Download SubRip \(.srt\) file](#)

📄 [Download Text \(.txt\) file](#)

Reading Assignment

0 points possible (ungraded)
Read Unit 12.2.4 of the notes. [\[LINK\]](#)

☒ Done

✓

Submit

📘 Answers are displayed within the problem

Discussion

Topic: Week 12 / 12.2.4

Hide Discussion

Add a Post

Show all posts ▼

by recent activity ▼

? Question

Hi Professors! What does it mean when a matrix is defective?

5

Homework 12.2.4.1

🧮 Calculator

Homework 12.2.4.1

4/4 points (graded)

Let $A = \begin{pmatrix} 3 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{pmatrix}$. Then which of the following are true:

- $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ is an eigenvector associated with eigenvalue **3**.

TRUE

✓ Answer: TRUE

Just multiply it out.

- $\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ is an eigenvector associated with eigenvalue **−1**.

TRUE

✓ Answer: TRUE

Just multiply it out.

- $\begin{pmatrix} 0 \\ \chi_1 \\ 0 \end{pmatrix}$, where $\chi_1 \neq 0$ is a scalar, is an eigenvector associated with eigenvalue **−1**.

TRUE

✓ Answer: TRUE

Just multiply it out.

- $\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$ is an eigenvector associated with eigenvalue **2**.

TRUE

✓ Answer: TRUE

Just multiply it out.

Submit

Answers are displayed within the problem

Homework 12.2.4.2

10.0/10.0 points (graded)

Let $A = \begin{pmatrix} \alpha_{0,0} & 0 & 0 \\ 0 & \alpha_{1,1} & 0 \\ 0 & 0 & \alpha_{2,2} \end{pmatrix}$. Then which of the following are true:

- $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ is an eigenvector associated with eigenvalue $\alpha_{0,0}$.

TRUE

✓ Answer: TRUE

Just multiply it out.

Calculator

- $\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ is an eigenvector associated with eigenvalue $\alpha_{1,1}$.

TRUE

✔ Answer: TRUE

Just multiply it out.

- $\begin{pmatrix} 0 \\ \chi_1 \\ 0 \end{pmatrix}$ where $\chi_1 \neq 0$ is an eigenvector associated with eigenvalue $\alpha_{1,1}$.

TRUE

✔ Answer: TRUE

Just multiply it out.

- $\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$ is an eigenvector associated with eigenvalue $\alpha_{2,2}$.

TRUE

✔ Answer: TRUE


Just multiply it out.

Submit

📘 Answers are displayed within the problem

Video 12.2.4 Part 2

[Start of transcript. Skip to the end.](#)



Dr. Robert van de Geijn: So hopefully, you remembered that the diagonal elements of a diagonal matrix are its eigenvalues, and then the eigenvectors can be found as the unit basis vectors.

What about this one?

Here we have a 3 by 3 upper triangular matrix.

Video

📄 [Download video file](#)

Transcripts

📄 [Download SubRip \(.srt\) file](#)

📄 [Download Text \(.txt\) file](#)

🧮 Calculator

Homework 12.2.4.3

10.0/10.0 points (graded)

Let $A = \begin{pmatrix} 3 & 1 & -1 \\ 0 & -1 & 2 \\ 0 & 0 & 2 \end{pmatrix}$. Then which of the following are true:

- $3, -1$, and 2 are eigenvalues of A .
- $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ is an eigenvector associated with eigenvalue 3 .

TRUE

✔ Answer: TRUE

Just multiply it out.

- $\begin{pmatrix} -1/4 \\ 1 \\ 0 \end{pmatrix}$ is an eigenvector associated with eigenvalue -1 .

TRUE

✔ Answer: TRUE

Just multiply it out.

- $\begin{pmatrix} -1/4\chi_1 \\ \chi_1 \\ 0 \end{pmatrix}$ where $\chi_1 \neq 0$ is an eigenvector associated with eigenvalue -1 .

TRUE

✔ Answer: TRUE

Just multiply it out.

- $\begin{pmatrix} 1/3 \\ 2/3 \\ 1 \end{pmatrix}$ is an eigenvector associated with eigenvalue 2 .

TRUE

✔ Answer: TRUE

Just multiply it out.

Submit

i Answers are displayed within the problem

Video 12.2.4 Part 3


[Start of transcript. Skip to the end.](#)



Dr. Robert van de Geijn: So this

▲

Calculator



▶ 0:00 / 0:00

▶ 2.0x

🔊

🔍

CC

“ ”

should have remembered that again, the diagonal elements of a triangular matrix are its eigenvalues.

And then once you subtract off the eigenvalue from the diagonal, you should be able to reduce the system to row echelon form and find the eigenvectors associated with the eigenvalues.

Video

Download video file

Transcripts

Download SubRip (.srt) file

Download Text (.txt) file

Homework 12.2.4.4

1/1 point (graded)

Let $A = \begin{pmatrix} \alpha_{0,0} & \alpha_{0,1} & \alpha_{0,2} \\ 0 & \alpha_{1,1} & \alpha_{1,2} \\ 0 & 0 & \alpha_{2,2} \end{pmatrix}$. Then the eigenvalues of this matrix are $\alpha_{0,0}$, $\alpha_{1,1}$, and $\alpha_{2,2}$.

TRUE Answer: TRUE

Submit

Answers are displayed within the problem

Homework 12.2.4.5

1/1 point (graded)

Consider $A = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$. Which of the following is true about this matrix:

(Mark all correct answers)

- ☒ $(1, \begin{pmatrix} 1 \\ 3 \\ 1 \end{pmatrix})$ is an eigenpair.
- ☒ $(0, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix})$ is an eigenpair.
- ☒ $(0, \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix})$ is an eigenpair.
- ☒ This matrix is defective.



Answer:

$$\det\left(\begin{pmatrix} 1-\lambda & 0 & 0 \\ 2 & -\lambda & 1 \\ 1 & 0 & -\lambda \end{pmatrix}\right) = \underbrace{[(1-\lambda)(-\lambda)^2 + 0 + 0]}_{=0} - \underbrace{[0 + 0 + 0]}_{=0}.$$

Calculator

$$\lambda^3 - \lambda^2 = \underbrace{\underbrace{-(1-\lambda)\lambda^2}_{\lambda^3 - \lambda^2}}_{-(1-\lambda)\lambda^2} = 0$$

So, $\lambda_0 = \lambda_1 = 0$ is a double root, while $\lambda_2 = 1$ is the third root.

<p>$\lambda_2 = 1$:</p> $A - \lambda_2 I = \begin{pmatrix} 0 & 0 & 0 \\ 2 & -1 & 1 \\ 1 & 0 & -1 \end{pmatrix}$ <p>We wish to find a nonzero vector in the null space:</p> $\begin{pmatrix} 0 & 0 & 0 \\ 2 & -1 & 1 \\ 1 & 0 & -1 \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$ <p>By examination, I noticed that</p> $\begin{pmatrix} 0 & 0 & 0 \\ 2 & -1 & 1 \\ 1 & 0 & -1 \end{pmatrix} \begin{pmatrix} 1 \\ 3 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$ <p>Eigenpair:</p> $\left(1, \begin{pmatrix} 1 \\ 3 \\ 1 \end{pmatrix}\right).$	<p>$\lambda_0 = \lambda_1 = 0$:</p> $A - 0I = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$ <p>Reducing this to row-echelon form gives us the matrix</p> $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}.$ <p>Notice that there is only one free variable, and hence this matrix is defective! The sole linearly independent eigenvector associated with $\lambda = 0$ is</p> $\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}.$ <p>Eigenpair:</p> $\left(0, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}\right)$
--	--



edX

- About
- Affiliates
- edX for Business
- Open edX
- Careers
- News

Legal

- Terms of Service & Honor Code
- Privacy Policy
- Accessibility Policy
- Trademark Policy
- Sitemap
- Cookie Policy
- Your Privacy Choices

Connect

[Idea Hub](#)

[Contact Us](#)

[Help Center](#)

[Security](#)

[Media Kit](#)



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)