

Introduction

- Start with a probability distribution $f(\mathbf{y}|\boldsymbol{\theta})$ for the data $\mathbf{y} = (y_1, \dots, y_n)$ given a vector of unknown parameters $\boldsymbol{\theta} = (\theta_1, \dots, \theta_K)$, and add a **prior** distribution $p(\boldsymbol{\theta}|\boldsymbol{\eta})$, where $\boldsymbol{\eta}$ is a vector of **hyperparameters**

Introduction

- Start with a probability distribution $f(\mathbf{y}|\boldsymbol{\theta})$ for the data $\mathbf{y} = (y_1, \dots, y_n)$ given a vector of unknown parameters $\boldsymbol{\theta} = (\theta_1, \dots, \theta_K)$, and add a **prior** distribution $p(\boldsymbol{\theta}|\boldsymbol{\eta})$, where $\boldsymbol{\eta}$ is a vector of **hyperparameters**
- Inference for $\boldsymbol{\theta}$ is based on its **posterior** distribution,

$$\begin{aligned} p(\boldsymbol{\theta}|\mathbf{y}, \boldsymbol{\eta}) &= \frac{p(\mathbf{y}, \boldsymbol{\theta}|\boldsymbol{\eta})}{p(\mathbf{y}|\boldsymbol{\eta})} = \frac{p(\mathbf{y}, \boldsymbol{\theta}|\boldsymbol{\eta})}{\int p(\mathbf{y}, \mathbf{u}|\boldsymbol{\eta}) d\mathbf{u}} \\ &= \frac{f(\mathbf{y}|\boldsymbol{\theta})p(\boldsymbol{\theta}|\boldsymbol{\eta})}{\int f(\mathbf{y}|\mathbf{u})p(\mathbf{u}|\boldsymbol{\eta}) d\mathbf{u}} = \frac{f(\mathbf{y}|\boldsymbol{\theta})p(\boldsymbol{\theta}|\boldsymbol{\eta})}{m(\mathbf{y}|\boldsymbol{\eta})}. \end{aligned}$$

Introduction

- Start with a probability distribution $f(\mathbf{y}|\boldsymbol{\theta})$ for the data $\mathbf{y} = (y_1, \dots, y_n)$ given a vector of unknown parameters $\boldsymbol{\theta} = (\theta_1, \dots, \theta_K)$, and add a **prior** distribution $p(\boldsymbol{\theta}|\boldsymbol{\eta})$, where $\boldsymbol{\eta}$ is a vector of **hyperparameters**
- Inference for $\boldsymbol{\theta}$ is based on its **posterior** distribution,

$$\begin{aligned} p(\boldsymbol{\theta}|\mathbf{y}, \boldsymbol{\eta}) &= \frac{p(\mathbf{y}, \boldsymbol{\theta}|\boldsymbol{\eta})}{p(\mathbf{y}|\boldsymbol{\eta})} = \frac{p(\mathbf{y}, \boldsymbol{\theta}|\boldsymbol{\eta})}{\int p(\mathbf{y}, \mathbf{u}|\boldsymbol{\eta}) d\mathbf{u}} \\ &= \frac{f(\mathbf{y}|\boldsymbol{\theta})p(\boldsymbol{\theta}|\boldsymbol{\eta})}{\int f(\mathbf{y}|\mathbf{u})p(\mathbf{u}|\boldsymbol{\eta}) d\mathbf{u}} = \frac{f(\mathbf{y}|\boldsymbol{\theta})p(\boldsymbol{\theta}|\boldsymbol{\eta})}{m(\mathbf{y}|\boldsymbol{\eta})}. \end{aligned}$$

- We refer to this formula as **Bayes' Theorem**. Note its similarity to the definition of conditional probability,

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{P(B|A)P(A)}{P(B)}$$

Example 2.1

- Consider the normal (Gaussian) likelihood,
 $f(y|\theta) = N(y|\theta, \sigma^2)$, $y \in \mathfrak{R}$, $\theta \in \mathfrak{R}$, and $\sigma > 0$ **known**. Take
 $p(\theta|\boldsymbol{\eta}) = N(\theta|\mu, \tau^2)$, where $\mu \in \mathfrak{R}$ and $\tau > 0$ are known
hyperparameters, so that $\boldsymbol{\eta} = (\mu, \tau)$. Then

$$p(\theta|y) = N\left(\theta \mid \frac{\sigma^2\mu + \tau^2y}{\sigma^2 + \tau^2}, \frac{\sigma^2\tau^2}{\sigma^2 + \tau^2}\right) .$$

Example 2.1

- Consider the normal (Gaussian) likelihood, $f(y|\theta) = N(y|\theta, \sigma^2)$, $y \in \mathbb{R}$, $\theta \in \mathbb{R}$, and $\sigma > 0$ **known**. Take $p(\theta|\boldsymbol{\eta}) = N(\theta|\mu, \tau^2)$, where $\mu \in \mathbb{R}$ and $\tau > 0$ are known hyperparameters, so that $\boldsymbol{\eta} = (\mu, \tau)$. Then

$$p(\theta|y) = N\left(\theta \mid \frac{\sigma^2\mu + \tau^2y}{\sigma^2 + \tau^2}, \frac{\sigma^2\tau^2}{\sigma^2 + \tau^2}\right).$$

- Write $B = \frac{\sigma^2}{\sigma^2 + \tau^2}$, and note that $0 < B < 1$. Then:

Example 2.1

- Consider the normal (Gaussian) likelihood, $f(y|\theta) = N(y|\theta, \sigma^2)$, $y \in \mathbb{R}$, $\theta \in \mathbb{R}$, and $\sigma > 0$ **known**. Take $p(\theta|\boldsymbol{\eta}) = N(\theta|\mu, \tau^2)$, where $\mu \in \mathbb{R}$ and $\tau > 0$ are known hyperparameters, so that $\boldsymbol{\eta} = (\mu, \tau)$. Then

$$p(\theta|y) = N\left(\theta \mid \frac{\sigma^2\mu + \tau^2y}{\sigma^2 + \tau^2}, \frac{\sigma^2\tau^2}{\sigma^2 + \tau^2}\right).$$

- Write $B = \frac{\sigma^2}{\sigma^2 + \tau^2}$, and note that $0 < B < 1$. Then:
 - $E(\theta|y) = B\mu + (1 - B)y$, a **weighted average** of the prior mean and the observed data value, with weights determined sensibly by the variances.

Example 2.1

- Consider the normal (Gaussian) likelihood, $f(y|\theta) = N(y|\theta, \sigma^2)$, $y \in \Re$, $\theta \in \Re$, and $\sigma > 0$ **known**. Take $p(\theta|\boldsymbol{\eta}) = N(\theta|\mu, \tau^2)$, where $\mu \in \Re$ and $\tau > 0$ are known hyperparameters, so that $\boldsymbol{\eta} = (\mu, \tau)$. Then

$$p(\theta|y) = N\left(\theta \mid \frac{\sigma^2\mu + \tau^2y}{\sigma^2 + \tau^2}, \frac{\sigma^2\tau^2}{\sigma^2 + \tau^2}\right).$$

- Write $B = \frac{\sigma^2}{\sigma^2 + \tau^2}$, and note that $0 < B < 1$. Then:
 - $E(\theta|y) = B\mu + (1 - B)y$, a **weighted average** of the prior mean and the observed data value, with weights determined sensibly by the variances.
 - $Var(\theta|y) = B\tau^2 \equiv (1 - B)\sigma^2$, **smaller** than τ^2 and σ^2 .

Example 2.1

- Consider the normal (Gaussian) likelihood, $f(y|\theta) = N(y|\theta, \sigma^2)$, $y \in \Re$, $\theta \in \Re$, and $\sigma > 0$ **known**. Take $p(\theta|\boldsymbol{\eta}) = N(\theta|\mu, \tau^2)$, where $\mu \in \Re$ and $\tau > 0$ are known hyperparameters, so that $\boldsymbol{\eta} = (\mu, \tau)$. Then

$$p(\theta|y) = N\left(\theta \mid \frac{\sigma^2\mu + \tau^2y}{\sigma^2 + \tau^2}, \frac{\sigma^2\tau^2}{\sigma^2 + \tau^2}\right).$$

- Write $B = \frac{\sigma^2}{\sigma^2 + \tau^2}$, and note that $0 < B < 1$. Then:
 - $E(\theta|y) = B\mu + (1 - B)y$, a **weighted average** of the prior mean and the observed data value, with weights determined sensibly by the variances.
 - $Var(\theta|y) = B\tau^2 \equiv (1 - B)\sigma^2$, **smaller** than τ^2 and σ^2 .
 - **Precision** (which is like "information") **is additive**:
$$Var^{-1}(\theta|y) = Var^{-1}(\theta) + Var^{-1}(y|\theta).$$

Sufficiency still helps

- **Lemma:** If $S(\mathbf{y})$ is **sufficient** for θ , then $p(\theta|\mathbf{y}) = p(\theta|s)$, so we may work with s instead of the entire dataset \mathbf{y} .

Sufficiency still helps


- **Lemma:** If $S(\mathbf{y})$ is **sufficient** for θ , then $p(\theta|\mathbf{y}) = p(\theta|s)$, so we may work with s instead of the entire dataset \mathbf{y} .
- **Example 2.2:** Consider again the normal/normal model where we now have an independent sample of size n from $f(\mathbf{y}|\theta)$. Since $S(\mathbf{y}) = \bar{y}$ is sufficient for θ , we have that $p(\theta|\mathbf{y}) = p(\theta|\bar{y})$.

Sufficiency still helps

- **Lemma:** If $S(\mathbf{y})$ is **sufficient** for θ , then $p(\theta|\mathbf{y}) = p(\theta|s)$, so we may work with s instead of the entire dataset \mathbf{y} .
- **Example 2.2:** Consider again the normal/normal model where we now have an independent sample of size n from $f(\mathbf{y}|\theta)$. Since $S(\mathbf{y}) = \bar{y}$ is sufficient for θ , we have that $p(\theta|\mathbf{y}) = p(\theta|\bar{y})$.
- But since we know that $f(\bar{y}|\theta) = N(\theta, \sigma^2/n)$, previous slide implies that


$$\begin{aligned} p(\theta|\bar{y}) &= N\left(\theta \mid \frac{(\sigma^2/n)\mu + \tau^2\bar{y}}{(\sigma^2/n) + \tau^2}, \frac{(\sigma^2/n)\tau^2}{(\sigma^2/n) + \tau^2}\right) \\ &= N\left(\theta \mid \frac{\sigma^2\mu + n\tau^2\bar{y}}{\sigma^2 + n\tau^2}, \frac{\sigma^2\tau^2}{\sigma^2 + n\tau^2}\right). \end{aligned}$$

Example: $\mu = 2, \bar{y} = 6, \tau = \sigma = 1$




- When $n = 1$ the prior and likelihood receive equal weight, so the posterior mean is $4 = \frac{2+6}{2}$.

Example: $\mu = 2, \bar{y} = 6, \tau = \sigma = 1$



- When $n = 1$ the prior and likelihood receive equal weight, so the posterior mean is $4 = \frac{2+6}{2}$.
- When $n = 10$ the data dominate the prior, resulting in a posterior mean much closer to \bar{y} .

Example: $\mu = 2, \bar{y} = 6, \tau = \sigma = 1$



- When $n = 1$ the prior and likelihood receive equal weight, so the posterior mean is $4 = \frac{2+6}{2}$.
- When $n = 10$ the data dominate the prior, resulting in a posterior mean much closer to \bar{y} .
- The posterior variance also shrinks as n gets larger; the posterior collapses to a point mass on \bar{y} as $n \rightarrow \infty$.

Three-stage Bayesian model

- If we are unsure as to the proper value of the hyperparameter η , the natural Bayesian solution would be to quantify this uncertainty in a **third-stage** distribution, sometimes called a **hyperprior**.

Three-stage Bayesian model

- If we are unsure as to the proper value of the hyperparameter η , the natural Bayesian solution would be to quantify this uncertainty in a **third-stage** distribution, sometimes called a **hyperprior**.
- Denoting this distribution by $h(\eta)$, the desired posterior for θ is now obtained by marginalizing over θ **and** η :

$$\begin{aligned} p(\boldsymbol{\theta}|\mathbf{y}) &= \frac{p(\mathbf{y}, \boldsymbol{\theta})}{p(\mathbf{y})} = \frac{\int p(\mathbf{y}, \boldsymbol{\theta}, \eta) d\eta}{\int \int p(\mathbf{y}, \mathbf{u}, \eta) d\eta d\mathbf{u}} \\ &= \frac{\int f(\mathbf{y}|\boldsymbol{\theta})p(\boldsymbol{\theta}|\eta)h(\eta) d\eta}{\int \int f(\mathbf{y}|\mathbf{u})p(\mathbf{u}|\eta)h(\eta) d\eta d\mathbf{u}} . \end{aligned}$$

Hierarchical modeling

- The hyperprior for η might itself depend on a collection of unknown parameters λ , resulting in a generalization of our three-stage model to one having a third-stage prior $h(\eta|\lambda)$ and a **fourth**-stage hyperprior $g(\lambda)$...

Hierarchical modeling

- The hyperprior for η might itself depend on a collection of unknown parameters λ , resulting in a generalization of our three-stage model to one having a third-stage prior $h(\eta|\lambda)$ and a **fourth**-stage hyperprior $g(\lambda)$...
- This enterprise of specifying a model over several levels is called **hierarchical modeling**, which is often helpful when the data are **nested**:

Hierarchical modeling

- The hyperprior for η might itself depend on a collection of unknown parameters λ , resulting in a generalization of our three-stage model to one having a third-stage prior $h(\eta|\lambda)$ and a **fourth**-stage hyperprior $g(\lambda)$...
- This enterprise of specifying a model over several levels is called **hierarchical modeling**, which is often helpful when the data are **nested**:
- **Example:** Test scores Y_{ijk} for student k in classroom j of school i :

$$Y_{ijk}|\theta_{ij} \sim N(\theta_{ij}, \sigma^2)$$

$$\theta_{ij}|\mu_i \sim N(\mu_i, \tau^2)$$

$$\mu_i|\lambda \sim N(\lambda, \kappa^2)$$

Adding $p(\lambda)$ and possibly $p(\sigma^2, \tau^2, \kappa^2)$ completes the specification!

Prediction

- Returning to two-level models, we often write

$$p(\boldsymbol{\theta}|\mathbf{y}) \propto f(\mathbf{y}|\boldsymbol{\theta})p(\boldsymbol{\theta}) ,$$

since the likelihood may be multiplied by any constant (or any function of \mathbf{y} alone) without altering $p(\boldsymbol{\theta}|\mathbf{y})$.

Prediction

- Returning to two-level models, we often write

$$p(\boldsymbol{\theta}|\mathbf{y}) \propto f(\mathbf{y}|\boldsymbol{\theta})p(\boldsymbol{\theta}) ,$$

since the likelihood may be multiplied by any constant (or any function of \mathbf{y} alone) without altering $p(\boldsymbol{\theta}|\mathbf{y})$.

- If y_{n+1} is a future observation, independent of \mathbf{y} given $\boldsymbol{\theta}$, then the **predictive** distribution for y_{n+1} is

$$p(y_{n+1}|\mathbf{y}) = \int f(y_{n+1}|\boldsymbol{\theta})p(\boldsymbol{\theta}|\mathbf{y})d\boldsymbol{\theta} ,$$

thanks to the conditional independence of y_{n+1} and \mathbf{y} .

Prediction

- Returning to two-level models, we often write

$$p(\boldsymbol{\theta}|\mathbf{y}) \propto f(\mathbf{y}|\boldsymbol{\theta})p(\boldsymbol{\theta}) ,$$

since the likelihood may be multiplied by any constant (or any function of \mathbf{y} alone) without altering $p(\boldsymbol{\theta}|\mathbf{y})$.

- If y_{n+1} is a future observation, independent of \mathbf{y} given $\boldsymbol{\theta}$, then the **predictive** distribution for y_{n+1} is

$$p(y_{n+1}|\mathbf{y}) = \int f(y_{n+1}|\boldsymbol{\theta})p(\boldsymbol{\theta}|\mathbf{y})d\boldsymbol{\theta} ,$$

thanks to the conditional independence of y_{n+1} and \mathbf{y} .

- The naive frequentist would use $f(y_{n+1}|\hat{\boldsymbol{\theta}})$ here, which is correct only for large n (i.e., when $p(\boldsymbol{\theta}|\mathbf{y})$ is a point mass at $\hat{\boldsymbol{\theta}}$).

Prior Distributions

- Suppose we require a prior distribution for
 θ = true proportion of U.S. men who are HIV-positive.

Prior Distributions

- Suppose we require a prior distribution for
 θ = true proportion of U.S. men who are HIV-positive.
- We cannot appeal to the usual long-term frequency notion of probability – it is not possible to even imagine “running the HIV epidemic over again” and reobserving θ . Here θ is random only because it is unknown **to us**.

Prior Distributions

- Suppose we require a prior distribution for
 θ = true proportion of U.S. men who are HIV-positive.
- We cannot appeal to the usual long-term frequency notion of probability – it is not possible to even imagine “running the HIV epidemic over again” and reobserving θ . Here θ is random only because it is unknown **to us**.
- Bayesian analysis is predicated on such a belief in **subjective probability** and its quantification in a prior distribution $p(\theta)$. But:

Prior Distributions

- Suppose we require a prior distribution for
 θ = true proportion of U.S. men who are HIV-positive.
- We cannot appeal to the usual long-term frequency notion of probability – it is not possible to even imagine “running the HIV epidemic over again” and reobserving θ . Here θ is random only because it is unknown **to us**.
- Bayesian analysis is predicated on such a belief in **subjective probability** and its quantification in a prior distribution $p(\theta)$. But:
 - How to create such a prior?

Prior Distributions

- Suppose we require a prior distribution for
 θ = true proportion of U.S. men who are HIV-positive.
- We cannot appeal to the usual long-term frequency notion of probability – it is not possible to even imagine “running the HIV epidemic over again” and reobserving θ . Here θ is random only because it is unknown **to us**.
- Bayesian analysis is predicated on such a belief in **subjective probability** and its quantification in a prior distribution $p(\theta)$. But:
 - How to create such a prior?
 - Are “objective” choices available?

Elicited Priors

- **Histogram approach:** Assign probability masses to the “possible” values in such a way that their sum is 1, and their relative contributions reflect the experimenter’s prior beliefs as closely as possible.

Elicited Priors

- **Histogram approach:** Assign probability masses to the “possible” values in such a way that their sum is 1, and their relative contributions reflect the experimenter’s prior beliefs as closely as possible.
 - **BUT:** Awkward for continuous or unbounded θ .

Elicited Priors

- **Histogram approach:** Assign probability masses to the “possible” values in such a way that their sum is 1, and their relative contributions reflect the experimenter’s prior beliefs as closely as possible.
 - **BUT:** Awkward for continuous or unbounded θ .
- **Matching a functional form:** Assume that the prior belongs to a parametric distributional family $p(\theta|\eta)$, choosing η so that the result matches the elicitee’s true prior beliefs as nearly as possible.

Elicited Priors

- **Histogram approach:** Assign probability masses to the “possible” values in such a way that their sum is 1, and their relative contributions reflect the experimenter’s prior beliefs as closely as possible.
 - **BUT:** Awkward for continuous or unbounded θ .
- **Matching a functional form:** Assume that the prior belongs to a parametric distributional family $p(\theta|\eta)$, choosing η so that the result matches the elicitee’s true prior beliefs as nearly as possible.
 - This approach limits the effort required of the elicitee, and also overcomes the finite support problem inherent in the histogram approach...

Elicited Priors

- **Histogram approach:** Assign probability masses to the “possible” values in such a way that their sum is 1, and their relative contributions reflect the experimenter’s prior beliefs as closely as possible.
 - **BUT:** Awkward for continuous or unbounded θ .
- **Matching a functional form:** Assume that the prior belongs to a parametric distributional family $p(\theta|\eta)$, choosing η so that the result matches the elicitee’s true prior beliefs as nearly as possible.
 - This approach limits the effort required of the elicitee, and also overcomes the finite support problem inherent in the histogram approach...
 - **BUT:** it may not be possible for the elicitee to “shoehorn” his or her prior beliefs into any of the standard parametric forms.

Conjugate Priors

- Defined as one that leads to a posterior distribution belonging to the same distributional family as the prior.

Conjugate Priors

- Defined as one that leads to a posterior distribution belonging to the same distributional family as the prior.
- **Example 2.5:** Suppose that X is distributed $\text{Poisson}(\theta)$, so that

$$f(x|\theta) = \frac{e^{-\theta} \theta^x}{x!}, \quad x \in \{0, 1, 2, \dots\}, \quad \theta > 0.$$

Conjugate Priors

- Defined as one that leads to a posterior distribution belonging to the same distributional family as the prior.
- Example 2.5:** Suppose that X is distributed $\text{Poisson}(\theta)$, so that

$$f(x|\theta) = \frac{e^{-\theta} \theta^x}{x!}, \quad x \in \{0, 1, 2, \dots\}, \quad \theta > 0.$$

- A reasonably flexible prior for θ having support on the positive real line is the *Gamma*(α, β) distribution,

$$p(\theta) = \frac{\theta^{\alpha-1} e^{-\theta/\beta}}{\Gamma(\alpha) \beta^\alpha}, \quad \theta > 0, \alpha > 0, \beta > 0,$$

Conjugate Priors

- The posterior is then

$$\begin{aligned} p(\theta|x) &\propto f(x|\theta)p(\theta) \\ &\propto \left(e^{-\theta}\theta^x\right) \left(\theta^{\alpha-1}e^{-\theta/\beta}\right) \\ &= \theta^{x+\alpha-1}e^{-\theta(1+1/\beta)} . \end{aligned}$$

Conjugate Priors

- The posterior is then

$$\begin{aligned} p(\theta|x) &\propto f(x|\theta)p(\theta) \\ &\propto \left(e^{-\theta}\theta^x\right) \left(\theta^{\alpha-1}e^{-\theta/\beta}\right) \\ &= \theta^{x+\alpha-1}e^{-\theta(1+1/\beta)} . \end{aligned}$$

- But this form is proportional to a *Gamma*(α' , β'), where

$$\alpha' = x + \alpha \quad \text{and} \quad \beta' = (1 + 1/\beta)^{-1}.$$

Since this is the **only** function proportional to our form that integrates to 1 and density functions uniquely determine distributions, $p(\theta|x)$ must indeed be *Gamma*(α' , β'), and the gamma is the **conjugate family** for the Poisson likelihood.

Notes on conjugate priors

- Can often guess the conjugate prior by looking at the likelihood as a function of θ , instead of x .

Notes on conjugate priors

- Can often guess the conjugate prior by looking at the likelihood as a function of θ , instead of x .
- In higher dimensions, priors that are **conditionally** conjugate are often available (and helpful).

Notes on conjugate priors

- Can often guess the conjugate prior by looking at the likelihood as a function of θ , instead of x .
- In higher dimensions, priors that are **conditionally** conjugate are often available (and helpful).
- a finite **mixture** of conjugate priors may be sufficiently flexible (allowing multimodality, heavier tails, etc.) while still enabling simplified posterior calculations.

Noninformative Prior

– is one that does not favor one θ value over another

• Examples:

Noninformative Prior

– is one that does not favor one θ value over another

● Examples:

● $\Theta = \{\theta_1, \dots, \theta_n\} \Rightarrow p(\theta_i) = 1/n, i = 1, \dots, n$

Noninformative Prior

– is one that does not favor one θ value over another

● Examples:

● $\Theta = \{\theta_1, \dots, \theta_n\} \Rightarrow p(\theta_i) = 1/n, \quad i = 1, \dots, n$

● $\Theta = [a, b], \quad -\infty < a < b < \infty$
 $\Rightarrow p(\theta) = 1/(b - a), \quad a < \theta < b$

Noninformative Prior

– is one that does not favor one θ value over another

• Examples:

• $\Theta = \{\theta_1, \dots, \theta_n\} \Rightarrow p(\theta_i) = 1/n, \quad i = 1, \dots, n$

• $\Theta = [a, b], \quad -\infty < a < b < \infty$
 $\Rightarrow p(\theta) = 1/(b - a), \quad a < \theta < b$

• $\Theta = (-\infty, \infty) \Rightarrow p(\theta) = c, \quad \text{any } c > 0$

Noninformative Prior

– is one that does not favor one θ value over another

Examples:

• $\Theta = \{\theta_1, \dots, \theta_n\} \Rightarrow p(\theta_i) = 1/n, i = 1, \dots, n$

• $\Theta = [a, b], -\infty < a < b < \infty$
 $\Rightarrow p(\theta) = 1/(b - a), a < \theta < b$

• $\Theta = (-\infty, \infty) \Rightarrow p(\theta) = c, \text{ any } c > 0$

This is an **improper** prior (does not integrate to 1), but its use can still be legitimate if

$\int f(\mathbf{x}|\theta)d\theta = K < \infty$, since then

$$p(\theta|\mathbf{x}) = \frac{f(\mathbf{x}|\theta) \cdot c}{\int f(\mathbf{x}|\theta) \cdot c d\theta} = \frac{f(\mathbf{x}|\theta)}{K},$$

so the posterior is just the **renormalized likelihood!**

Jeffreys Prior

- another noninformative prior, given in the univariate case by

$$p(\theta) = [I(\theta)]^{1/2} ,$$

where $I(\theta)$ is the expected Fisher information in the model, namely

$$I(\theta) = -E_{\mathbf{x}|\theta} \left[\frac{\partial^2}{\partial \theta^2} \log f(\mathbf{x}|\theta) \right] .$$

Jeffreys Prior

- another noninformative prior, given in the univariate case by

$$p(\theta) = [I(\theta)]^{1/2} ,$$

where $I(\theta)$ is the expected Fisher information in the model, namely

$$I(\theta) = -E_{\mathbf{x}|\theta} \left[\frac{\partial^2}{\partial \theta^2} \log f(\mathbf{x}|\theta) \right] .$$

- Unlike the uniform, the Jeffreys prior is **invariant to 1-1 transformations**. That is, computing the Jeffreys prior for some 1-1 transformation $\gamma = g(\theta)$ directly produces the same answer as computing the Jeffreys prior for θ and subsequently performing the usual Jacobian transformation to the γ scale (see p.54, problem 7).

Other Noninformative Priors

- When $f(x|\theta) = f(x - \theta)$ (location parameter family),

$$p(\theta) = 1, \theta \in \mathbb{R}$$

is invariant under location transformations ($Y = X + c$).

Other Noninformative Priors

- When $f(x|\theta) = f(x - \theta)$ (location parameter family),

$$p(\theta) = 1, \theta \in \mathbb{R}$$

is invariant under location transformations ($Y = X + c$).

- When $f(x|\sigma) = \frac{1}{\sigma} f(\frac{x}{\sigma})$, $\sigma > 0$ (scale parameter family),

$$p(\sigma) = \frac{1}{\sigma}, \sigma > 0$$

is invariant under scale transformations ($Y = cX$, $c > 0$).

Other Noninformative Priors

- When $f(x|\theta) = f(x - \theta)$ (location parameter family),

$$p(\theta) = 1, \theta \in \mathbb{R}$$

is invariant under location transformations ($Y = X + c$).

- When $f(x|\sigma) = \frac{1}{\sigma} f(\frac{x}{\sigma})$, $\sigma > 0$ (scale parameter family),

$$p(\sigma) = \frac{1}{\sigma}, \sigma > 0$$

is invariant under scale transformations ($Y = cX$, $c > 0$).

- When $f(x|\theta, \sigma) = \frac{1}{\sigma} f(\frac{x-\theta}{\sigma})$ (location-scale family), prior “independence” suggests

$$p(\theta, \sigma) = \frac{1}{\sigma}, \theta \in \mathbb{R}, \sigma > 0.$$

Bayesian Inference: Point Estimation

- **Easy!** Simply choose an appropriate distributional summary: posterior **mean**, **median**, or **mode**.

Bayesian Inference: Point Estimation

- **Easy!** Simply choose an appropriate distributional summary: posterior **mean**, **median**, or **mode**.
- **Mode** is often easiest to compute (no integration), but is often least representative of “middle”, especially for one-tailed distributions.

Bayesian Inference: Point Estimation

- **Easy!** Simply choose an appropriate distributional summary: posterior **mean**, **median**, or **mode**.
- **Mode** is often easiest to compute (no integration), but is often least representative of “middle”, especially for one-tailed distributions.
- **Mean** has the opposite property, tending to “chase” heavy tails (just like the sample mean \bar{X})

Bayesian Inference: Point Estimation

- **Easy!** Simply choose an appropriate distributional summary: posterior **mean**, **median**, or **mode**.
- **Mode** is often easiest to compute (no integration), but is often least representative of “middle”, especially for one-tailed distributions.
- **Mean** has the opposite property, tending to “chase” heavy tails (just like the sample mean \bar{X})
- **Median** is probably the best compromise overall, though can be awkward to compute, since it is the solution θ^{median} to

$$\int_{-\infty}^{\theta^{median}} p(\theta|x) d\theta = \frac{1}{2} .$$

Example: The General Linear Model

- Let \mathbf{Y} be an $n \times 1$ data vector, X an $n \times p$ matrix of covariates, and adopt the likelihood and prior structure,

$$\mathbf{Y}|\boldsymbol{\beta} \sim N_n(X\boldsymbol{\beta}, \Sigma) \text{ and } \boldsymbol{\beta} \sim N_p(A\boldsymbol{\alpha}, V)$$

Example: The General Linear Model

- Let \mathbf{Y} be an $n \times 1$ data vector, X an $n \times p$ matrix of covariates, and adopt the likelihood and prior structure,

$$\mathbf{Y}|\boldsymbol{\beta} \sim N_n(X\boldsymbol{\beta}, \Sigma) \text{ and } \boldsymbol{\beta} \sim N_p(A\boldsymbol{\alpha}, V)$$

- Then the posterior distribution of $\boldsymbol{\beta}|\mathbf{Y}$ is

$$\boldsymbol{\beta}|\mathbf{Y} \sim N(D\mathbf{d}, D) \text{ , where}$$

$$D^{-1} = X^T \Sigma^{-1} X + V^{-1} \text{ and } \mathbf{d} = X^T \Sigma^{-1} \mathbf{Y} + V^{-1} A\boldsymbol{\alpha}.$$

Example: The General Linear Model

- Let \mathbf{Y} be an $n \times 1$ data vector, X an $n \times p$ matrix of covariates, and adopt the likelihood and prior structure,

$$\mathbf{Y}|\boldsymbol{\beta} \sim N_n(X\boldsymbol{\beta}, \Sigma) \text{ and } \boldsymbol{\beta} \sim N_p(A\boldsymbol{\alpha}, V)$$

- Then the posterior distribution of $\boldsymbol{\beta}|\mathbf{Y}$ is

$$\boldsymbol{\beta}|\mathbf{Y} \sim N(D\mathbf{d}, D) \text{ , where}$$

$$D^{-1} = X^T \Sigma^{-1} X + V^{-1} \text{ and } \mathbf{d} = X^T \Sigma^{-1} \mathbf{Y} + V^{-1} A\boldsymbol{\alpha}.$$

- $V^{-1} = 0$ delivers a “flat” prior; if $\Sigma = \sigma^2 I_p$, we get

$$\boldsymbol{\beta}|\mathbf{Y} \sim N\left(\hat{\boldsymbol{\beta}}, \sigma^2(X'X)^{-1}\right) \text{ , where}$$

$$\hat{\boldsymbol{\beta}} = (X'X)^{-1}X'\mathbf{y} \iff \text{usual likelihood approach!}$$

Bayesian Inference: Interval Estimation

- The Bayesian analogue of a frequentist CI is referred to as a **credible set**: a $100 \times (1 - \alpha)\%$ credible set for θ is a subset C of Θ such that

$$1 - \alpha \leq P(C|\mathbf{y}) = \int_C p(\boldsymbol{\theta}|\mathbf{y}) d\boldsymbol{\theta} .$$

Bayesian Inference: Interval Estimation

- The Bayesian analogue of a frequentist CI is referred to as a **credible set**: a $100 \times (1 - \alpha)\%$ credible set for θ is a subset C of Θ such that

$$1 - \alpha \leq P(C|\mathbf{y}) = \int_C p(\boldsymbol{\theta}|\mathbf{y}) d\boldsymbol{\theta} .$$

- In continuous settings, we can obtain coverage **exactly** $1 - \alpha$ at **minimum size** via the **highest posterior density (HPD)** credible set,

$$C = \{\boldsymbol{\theta} \in \Theta : p(\boldsymbol{\theta}|\mathbf{y}) \geq k(\alpha)\} ,$$

where $k(\alpha)$ is the **largest** constant such that

$$P(C|\mathbf{y}) \geq 1 - \alpha .$$

Interval Estimation (cont'd)

- Simpler alternative: the **equal-tail** set, which takes the $\alpha/2$ - and $(1 - \alpha/2)$ -quantiles of $p(\theta|\mathbf{y})$.

Interval Estimation (cont'd)

- Simpler alternative: the **equal-tail** set, which takes the $\alpha/2$ - and $(1 - \alpha/2)$ -quantiles of $p(\theta|\mathbf{y})$.
- Specifically, consider q_L and q_U , the $\alpha/2$ - and $(1 - \alpha/2)$ -quantiles of $p(\theta|\mathbf{y})$:

$$\int_{-\infty}^{q_L} p(\theta|\mathbf{y})d\theta = \alpha/2 \quad \text{and} \quad \int_{q_U}^{\infty} p(\theta|\mathbf{y})d\theta = \alpha/2 .$$

Then clearly $P(q_L < \theta < q_U|\mathbf{y}) = 1 - \alpha$; our confidence that θ lies in (q_L, q_U) is $100 \times (1 - \alpha)\%$. Thus this interval is a $100 \times (1 - \alpha)\%$ credible set (**“Bayesian CI”**) for θ .

Interval Estimation (cont'd)

- Simpler alternative: the **equal-tail** set, which takes the $\alpha/2$ - and $(1 - \alpha/2)$ -quantiles of $p(\theta|\mathbf{y})$.
- Specifically, consider q_L and q_U , the $\alpha/2$ - and $(1 - \alpha/2)$ -quantiles of $p(\theta|\mathbf{y})$:


$$\int_{-\infty}^{q_L} p(\theta|\mathbf{y})d\theta = \alpha/2 \quad \text{and} \quad \int_{q_U}^{\infty} p(\theta|\mathbf{y})d\theta = \alpha/2 .$$

Then clearly $P(q_L < \theta < q_U|\mathbf{y}) = 1 - \alpha$; our confidence that θ lies in (q_L, q_U) is $100 \times (1 - \alpha)\%$. Thus this interval is a $100 \times (1 - \alpha)\%$ credible set (**“Bayesian CI”**) for θ .

- This interval is relatively easy to compute, and enjoys a direct interpretation (**“The probability that θ lies in (q_L, q_U) is $(1 - \alpha)$ ”**) that the frequentist interval does not.


Interval Estimation: Example

Using a $\text{Gamma}(2, 1)$ posterior distribution and $k(\alpha) = 0.1$:




Equal tail interval is a bit **wider**, but **easier to compute** (just two gamma quantiles), and also **transformation invariant**.

Ex: $Y \sim \text{Bin}(10, \theta)$, $\theta \sim U(0, 1)$, $y_{obs} = 7$



Ex: $Y \sim \text{Bin}(10, \theta)$, $\theta \sim U(0, 1)$, $y_{obs} = 7$




Plot $\text{Beta}(y_{obs} + 1, n - y_{obs} + 1) = \text{Beta}(8, 4)$ posterior in R/S:

```
> theta <- seq(from=0, to=1, length=101)
```

```
> yobs <- 7; n <- 10
```

```
> plot(theta, dbeta(theta, yobs+1, n-yobs+1), type="l")
```

Ex: $Y \sim \text{Bin}(10, \theta)$, $\theta \sim U(0, 1)$, $y_{obs} = 7$



Plot $\text{Beta}(y_{obs} + 1, n - y_{obs} + 1) = \text{Beta}(8, 4)$ posterior in R/S:

```
> theta <- seq(from=0, to=1, length=101)
> yobs <- 7; n <- 10
> plot(theta, dbeta(theta, yobs+1, n-yobs+1), type="l")
```

Add 95% equal-tail Bayesian CI (dotted vertical lines):

```
> abline(v=qbeta(.5, yobs+1, n-yobs+1))
> abline(v=qbeta(c(.025, .975), yobs+1, n-yobs+1), lty=2)
```

Bayesian hypothesis testing

- Classical approach bases accept/reject decision on

$$\text{p-value} = P\{T(\mathbf{Y}) \text{ more "extreme" than } T(\mathbf{y}_{obs}) | \boldsymbol{\theta}, H_0\} ,$$

where “extremeness” is in the direction of H_A

Bayesian hypothesis testing

- Classical approach bases accept/reject decision on

$$\text{p-value} = P\{T(\mathbf{Y}) \text{ more "extreme" than } T(\mathbf{y}_{obs}) | \boldsymbol{\theta}, H_0\} ,$$

where “extremeness” is in the direction of H_A

- Several *troubles* with this approach:

Bayesian hypothesis testing

- Classical approach bases accept/reject decision on

$$\text{p-value} = P\{T(\mathbf{Y}) \text{ more "extreme" than } T(\mathbf{y}_{obs}) | \boldsymbol{\theta}, H_0\} ,$$

where “extremeness” is in the direction of H_A

- Several *troubles* with this approach:
 - hypotheses must be *nested*

Bayesian hypothesis testing

- Classical approach bases accept/reject decision on

$$\text{p-value} = P\{T(\mathbf{Y}) \text{ more "extreme" than } T(\mathbf{y}_{obs}) | \boldsymbol{\theta}, H_0\} ,$$

where “extremeness” is in the direction of H_A

- Several *troubles* with this approach:
 - hypotheses must be *nested*
 - p-value can only offer evidence *against* the null

Bayesian hypothesis testing

- Classical approach bases accept/reject decision on

$$\text{p-value} = P\{T(\mathbf{Y}) \text{ more "extreme" than } T(\mathbf{y}_{obs}) | \boldsymbol{\theta}, H_0\} ,$$

where “extremeness” is in the direction of H_A

- Several *troubles* with this approach:
 - hypotheses must be *nested*
 - p-value can only offer evidence *against* the null
 - p-value is *not* the “probability that H_0 is true” (but is often erroneously interpreted this way)

Bayesian hypothesis testing

- Classical approach bases accept/reject decision on
 $\text{p-value} = P\{T(\mathbf{Y}) \text{ more "extreme" than } T(\mathbf{y}_{obs}) | \boldsymbol{\theta}, H_0\}$,
where “extremeness” is in the direction of H_A
- Several *troubles* with this approach:
 - hypotheses must be *nested*
 - p-value can only offer evidence *against* the null
 - p-value is *not* the “probability that H_0 is true” (but is often erroneously interpreted this way)
 - As a result of the dependence on “more extreme” $T(\mathbf{Y})$ values, two experiments with different *designs* but identical likelihoods could result in different p-values, *violating the Likelihood Principle!*

Bayesian hypothesis testing (cont'd)

- Bayesian approach: Select the model with the largest posterior probability, $P(M_i|\mathbf{y}) = p(\mathbf{y}|M_i)p(M_i)/p(\mathbf{y})$,

where
$$p(\mathbf{y}|M_i) = \int f(\mathbf{y}|\boldsymbol{\theta}_i, M_i)\pi_i(\boldsymbol{\theta}_i)d\boldsymbol{\theta}_i .$$

Bayesian hypothesis testing (cont'd)

- Bayesian approach: Select the model with the largest posterior probability, $P(M_i|\mathbf{y}) = p(\mathbf{y}|M_i)p(M_i)/p(\mathbf{y})$,

$$\text{where } p(\mathbf{y}|M_i) = \int f(\mathbf{y}|\boldsymbol{\theta}_i, M_i)\pi_i(\boldsymbol{\theta}_i)d\boldsymbol{\theta}_i .$$

- For two models, the quantity commonly used to summarize these results is the **Bayes factor**,

$$BF = \frac{P(M_1|\mathbf{y})/P(M_2|\mathbf{y})}{P(M_1)/P(M_2)} = \frac{p(\mathbf{y} | M_1)}{p(\mathbf{y} | M_2)} ,$$

i.e., the likelihood ratio if both hypotheses are simple

Bayesian hypothesis testing (cont'd)

- Bayesian approach: Select the model with the largest posterior probability, $P(M_i|\mathbf{y}) = p(\mathbf{y}|M_i)p(M_i)/p(\mathbf{y})$,

$$\text{where } p(\mathbf{y}|M_i) = \int f(\mathbf{y}|\boldsymbol{\theta}_i, M_i)\pi_i(\boldsymbol{\theta}_i)d\boldsymbol{\theta}_i .$$

- For two models, the quantity commonly used to summarize these results is the **Bayes factor**,

$$BF = \frac{P(M_1|\mathbf{y})/P(M_2|\mathbf{y})}{P(M_1)/P(M_2)} = \frac{p(\mathbf{y} | M_1)}{p(\mathbf{y} | M_2)} ,$$

i.e., the likelihood ratio if both hypotheses are simple

- **Problem:** If $\pi_i(\boldsymbol{\theta}_i)$ is **improper**, then $p(\mathbf{y}|M_i)$ necessarily is as well \implies **BF is not well-defined!**...

Bayesian hypothesis testing (cont'd)

When the BF is not well-defined, several alternatives:

- Modify the definition of BF : partial Bayes factor, fractional Bayes factor (text, pp.41-42)

Bayesian hypothesis testing (cont'd)

When the BF is not well-defined, several alternatives:

- Modify the definition of BF : partial Bayes factor, fractional Bayes factor (text, pp.41-42)
- Switch to the conditional predictive distribution,

$$f(y_i | \mathbf{y}_{(i)}) = \frac{f(\mathbf{y})}{f(\mathbf{y}_{(i)})} = \int f(y_i | \boldsymbol{\theta}, \mathbf{y}_{(i)}) p(\boldsymbol{\theta} | \mathbf{y}_{(i)}) d\boldsymbol{\theta} ,$$

which will be proper if $p(\boldsymbol{\theta} | \mathbf{y}_{(i)})$ is. Assess model fit via plots or a suitable summary (say, $\prod_{i=1}^n f(y_i | \mathbf{y}_{(i)})$).

Bayesian hypothesis testing (cont'd)

When the BF is not well-defined, several alternatives:

- **Modify the definition of BF :** partial Bayes factor, fractional Bayes factor (text, pp.41-42)
- Switch to the **conditional predictive distribution**,

$$f(y_i | \mathbf{y}_{(i)}) = \frac{f(\mathbf{y})}{f(\mathbf{y}_{(i)})} = \int f(y_i | \boldsymbol{\theta}, \mathbf{y}_{(i)}) p(\boldsymbol{\theta} | \mathbf{y}_{(i)}) d\boldsymbol{\theta} ,$$

which will be proper if $p(\boldsymbol{\theta} | \mathbf{y}_{(i)})$ is. Assess model fit via plots or a suitable summary (say, $\prod_{i=1}^n f(y_i | \mathbf{y}_{(i)})$).

- **Penalized likelihood criteria:** the Akaike information criterion (**AIC**), Bayesian information criterion (**BIC**), or Deviance information criterion (**DIC**).

Bayesian hypothesis testing (cont'd)

When the BF is not well-defined, several alternatives:

- **Modify the definition of BF :** partial Bayes factor, fractional Bayes factor (text, pp.41-42)
- Switch to the **conditional predictive distribution**,

$$f(y_i|\mathbf{y}_{(i)}) = \frac{f(\mathbf{y})}{f(\mathbf{y}_{(i)})} = \int f(y_i|\boldsymbol{\theta}, \mathbf{y}_{(i)})p(\boldsymbol{\theta}|\mathbf{y}_{(i)})d\boldsymbol{\theta} ,$$

which will be proper if $p(\boldsymbol{\theta}|\mathbf{y}_{(i)})$ is. Assess model fit via plots or a suitable summary (say, $\prod_{i=1}^n f(y_i|\mathbf{y}_{(i)})$).

- **Penalized likelihood criteria:** the Akaike information criterion (**AIC**), Bayesian information criterion (**BIC**), or Deviance information criterion (**DIC**).
- **IOU on all this – Chapter 6!**

Example: Consumer preference data

- Suppose 16 taste testers compare two types of ground beef patty (one stored in a deep freeze, the other in a less expensive freezer). The food chain is interested in whether storage in the higher-quality freezer translates into a "substantial improvement in taste."

Example: Consumer preference data

- Suppose 16 taste testers compare two types of ground beef patty (one stored in a deep freeze, the other in a less expensive freezer). The food chain is interested in whether storage in the higher-quality freezer translates into a "substantial improvement in taste."
- **Experiment:** In a test kitchen, the patties are defrosted and prepared by a single chef/statistician, who randomizes the order in which the patties are served in double-blind fashion.

Example: Consumer preference data

- Suppose 16 taste testers compare two types of ground beef patty (one stored in a deep freeze, the other in a less expensive freezer). The food chain is interested in whether storage in the higher-quality freezer translates into a "substantial improvement in taste."
- **Experiment:** In a test kitchen, the patties are defrosted and prepared by a single chef/statistician, who randomizes the order in which the patties are served in double-blind fashion.
- **Result:** 13 of the 16 testers state a preference for the more expensive patty.

Example: Consumer preference data

● **Likelihood:** Let

θ = prob. consumers prefer more expensive patty

$$Y_i = \begin{cases} 1 & \text{if tester } i \text{ prefers more expensive patty} \\ 0 & \text{otherwise} \end{cases}$$

Example: Consumer preference data

● **Likelihood:** Let

θ = prob. consumers prefer more expensive patty

$$Y_i = \begin{cases} 1 & \text{if tester } i \text{ prefers more expensive patty} \\ 0 & \text{otherwise} \end{cases}$$

● Assuming **independent testers** and **constant θ** , then if $X = \sum_{i=1}^{16} Y_i$, we have $X|\theta \sim \text{Binomial}(16, \theta)$,

$$f(x|\theta) = \binom{16}{x} \theta^x (1 - \theta)^{16-x} .$$

Example: Consumer preference data

- **Likelihood:** Let

θ = prob. consumers prefer more expensive patty

$$Y_i = \begin{cases} 1 & \text{if tester } i \text{ prefers more expensive patty} \\ 0 & \text{otherwise} \end{cases}$$


- Assuming **independent testers** and **constant θ** , then if $X = \sum_{i=1}^{16} Y_i$, we have $X|\theta \sim \text{Binomial}(16, \theta)$,

$$f(x|\theta) = \binom{16}{x} \theta^x (1 - \theta)^{16-x} .$$

- The **beta** distribution offers a conjugate family, since


$$p(\theta) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \theta^{\alpha-1} (1 - \theta)^{\beta-1} .$$

Three 'minimally informative' priors




The posterior is then $Beta(x + \alpha, 16 - x + \beta)...$

Three corresponding posteriors



● Note ordering of posteriors; consistent with priors.

Three corresponding posteriors



- Note ordering of posteriors; consistent with priors.
- All three produce 95% equal-tail credible intervals that exclude 0.5 \Rightarrow there **is** an improvement in taste.

Posterior summaries

Prior distribution	Posterior quantile			$P(\theta > .6 x)$
	.025	.500	.975	
$Beta(.5, .5)$	0.579	0.806	0.944	0.964
$Beta(1, 1)$	0.566	0.788	0.932	0.954
$Beta(2, 2)$	0.544	0.758	0.909	0.930

Posterior summaries

Prior distribution	Posterior quantile			$P(\theta > .6 x)$
	.025	.500	.975	
$Beta(.5, .5)$	0.579	0.806	0.944	0.964
$Beta(1, 1)$	0.566	0.788	0.932	0.954
$Beta(2, 2)$	0.544	0.758	0.909	0.930

- Suppose we define “*substantial* improvement in taste” as $\theta \geq 0.6$. Then under the uniform prior, the Bayes factor in favor of $M_1 : \theta \geq 0.6$ over $M_2 : \theta < 0.6$ is

$$BF = \frac{0.954/0.046}{0.4/0.6} = 31.1 ,$$

or fairly strong evidence (adjusted odds about 30:1) in favor of a substantial improvement in taste.