<u>Help</u>

sandipan_dey >

<u>Course</u>

Progress


<u>Dates</u>

Discussion

MO Index

☆ Course / 9 Convergence, Accuracy, and an IVP Solver / 9.2 Accuracy and Convergence





MO2.4 MO2.6 MO2.7

Let's consider the error for the Forward Euler simulation of coffee cooling. Figure 9.1 shows that the error decreases by about a factor of 10 for a factor of 10 decrease in Δt from 50 to 5 minutes. This would seem to indicate a first order (p=1) method since the error seems proportional to Δt .


To see more certainly that Forward Euler is p=1, we plot the maximum error (e_{\max}) versus a range of Δt in Figure 9.2. This plot is done using a log-log scale because the slope then is directly the observed order of accuracy. To see that this is true, recall that the definition of the order of accuracy says that,

$$e_{\max} = O(\Delta t^p) \tag{9.4}$$

$$\log e_{\max} = O(\log(\Delta t^p)) \tag{9.5}$$

$$\Rightarrow \log e_{\max} = O(p \log \Delta t) \tag{9.6}$$

Thus, p is the slope of the $\log e_{\max}$ versus $\log \Delta t$ plot. As can be seen in Figure 9.2, the slope is 1.



edX for Business

Discussions

All posts sorted by recent activity

© All Rights Reserved

Open ed Xious Next > <u>Careers</u> **News**

Legal

Terms of Service & Honor Code

Privacy Policy

Accessibility Policy

<u>Trademark Policy</u>

<u>Sitemap</u>

Cookie Policy

Your Privacy Choices

Connect

Blog

Contact Us

Help Center

<u>Security</u>

Media Kit















© 2023 edX LLC. All rights reserved.

深圳市恒宇博科技有限公司 <u>粤ICP备17044299号-2</u>

transcript. Skip to the end.