



MITx: 15.053x Optimization Methods in Business Analytics



Bookmarks

▸ General Information

▸ Week 1

▸ Week 2

▼ Week 3

Lecture

Lecture questions due Sep 27, 2016 at 19:30 IST

**Recitation****Problem Set 3**

Homework 3 due Sep 27, 2016 at 19:30 IST



Week 3 > Recitation > Practice Problem 2



Bookmark

PART A

Consider two non-negative integral variables x_1 and x_2 . Suppose that $x_1 \leq 60$ and $x_2 \leq 50$. Model the following constraint using an additional binary variable w :

$$x_1 < 30 \text{ or } x_2 \geq 20.$$

Note that the first inequality is strict. Select the correct way of modeling this “OR” constraint using linear and integer constraints. Select the (one) correct answer.

☐ $x_1 \leq 29 - M(1 - w)$
 $x_2 \geq 20 + Mw$

☒ $x_1 \leq 29 + M(1 - w)$ ✓
 $x_2 \geq 20 - Mw$

☐ $x_1 \leq 30 - M(1 - w)$
 $x_2 \geq 20 + Mw$

● $x_1 \leq 30 + M(1 - w)$
 $x_2 \geq 20 - Mw$

EXPLANATION

Solution

$$x_1 \leq 29 + M(1 - w)$$
$$x_2 \geq 20 - Mw$$

It is often difficult to work with strict inequalities when modeling MIPs (Mixed Integer/Linear Programs). In this case, x_1 is known to be integer valued. We transformed " $x_1 < 30$ " into " $x_1 \leq 29$."

PART B

How small can M be in the correct answer for PART A?

✓ Answer: 31

31

EXPLANATION

Solution**31**

In order to make the constraint $x_1 \leq 29 + M(1 - w)$ redundant when $w = 0$, one needs M to be at least **31**. In order to make the constraint $x_2 \geq 20 - Mw$ be redundant when $w = 1$, we need M to be at least **20**.

© All Rights Reserved



© 2016 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

POWERED BY
OPENedX®

