



## <u>Lecture 14: Wald's Test, Likelihood</u> <u>Ratio Test, and Implicit Hypothesis</u>

Course > Unit 4 Hypothesis testing > Test

> 7. Wald's Test in 1 Dimension

### **Audit Access Expires Dec 24, 2019**

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. **Upgrade now** 

## 7. Wald's Test in 1 Dimension

# **Comparing Quantiles**

1/1 point (graded)

Let 
$$Z \sim \mathcal{N}\left(0,1
ight)$$
 . Then  $Z^2 \sim \chi_1^2$  .

The **quantile**  $q_{\alpha}\left(\chi_{1}^{2}\right)$  of the  $\chi_{1}^{2}-$  distibution is the number such that

$$\mathbf{P}\left(Z^{2}>q_{lpha}\left(\chi_{1}^{2}
ight)
ight)=lpha.$$

Find the quantiles of the  $\chi_1^2$  distribution in terms of the quantiles of the normal distribution. That is, write  $q_{\alpha}\left(\chi_1^2\right)$  in terms of  $q_{\alpha'}\left(\mathcal{N}\left(0,1\right)\right)$  where  $\alpha'$  is a function of  $\alpha$ .

(Enter **q(alpha)** for the quantile  $q_{lpha}\left(\mathcal{N}\left(0,1
ight)
ight)$  of the normal distribution.)

$$q_{lpha}\left(\chi_{1}^{2}
ight)= egin{pmatrix} (\mathsf{q}(\mathsf{alpha/2}))^{2} & & \checkmark \text{ Answer: } (\mathsf{q}(\mathsf{alpha/2}))^{2} \end{pmatrix}$$

#### STANDARD NOTATION

### **Solution:**

Since  $Z^2>q$  for any  $q>0\,$  if and only if  $|Z|>\sqrt{q},\,$  we have

$$P\left(Z^2>q_lpha\left(\chi_1^2
ight)
ight)\,=\,P\left(|Z|>\sqrt{q_lpha\left(\chi_1^2
ight)}
ight)\,=\,lpha.$$

Since  $Z\sim\mathcal{N}\left(0,1
ight),\,P(|Z|>\sqrt{q_{lpha}\left(\chi_{1}^{2}
ight)}
ight)\,=\,lpha\,$  if and only if

$$\sqrt{q_{lpha}\left(\chi_{1}^{2}
ight)}=q_{lpha/2}\left(\mathcal{N}\left(0,1
ight)
ight)$$

Hence  $q_{lpha}\left(\chi_{1}^{2}
ight)=q_{lpha/2}(\mathcal{N}\left(0,1
ight))^{2}.$ 

For example, for  $\alpha=5\%$ , using a table (e.g. https://people.richland.edu/james/lecture/m170/tbl-chi.html) or software (e.g. R), we have

$$egin{aligned} q_{0.05}\left(\chi_1^2
ight) &pprox 3.84. \ &\left(q_{0.025}\left(\mathcal{N}\left(0,1
ight)
ight)
ight)^2 &pprox \left(1.96
ight)^2 pprox 3.84. \end{aligned}$$

Submit


You have used 3 of 3 attempts

• Answers are displayed within the problem

Video note: In the video below at 5:27, Prof Rigollet misprinted on the board: the bottom inequality should read:

$$\sqrt{n}rac{\left|\hat{ heta}- heta
ight|}{\sigma}\,>\,q_{lpha/2}\left(\mathcal{N}\left(0,1
ight)
ight).$$

# Wald's Test Continued



### Wald's Test in 1 dimension

In 1 dimension, Wald's Test coincides with the two-sided test based on on the asymptotic normality of the MLE.

Given the hypotheses

$$H_0: heta^* = heta_0$$

$$H_1: heta^* 
eq heta_0,$$

a two-sided test of level  $\alpha$ , based on the asymptotic normality of the MLE, is

$$\psi_{lpha}=\mathbf{1}\left(\sqrt{nI\left( heta_{0}
ight)}\left|\hat{ heta}^{\mathrm{MLE}}- heta_{0}
ight|>q_{lpha/2}\left(\mathcal{N}\left(0,1
ight)
ight)
ight)$$

where the Fisher information  $I( heta_0)^{-1}$  is the asymptotic variance of  $\hat{ heta}^{ ext{MLE}}$  under the null hypothesis.

On the other hand, a Wald's test of level  $\, \alpha \,$  is

$$egin{array}{lcl} \psi_{lpha}^{
m Wald} & = & \mathbf{1} \left( n I\left( heta_0
ight) \left( \hat{ heta}^{
m MLE} - heta_0 
ight)^2 \, > \, q_lpha\left(\chi_1^2
ight) 
ight) \ \\ & = & \mathbf{1} \left( \sqrt{n I\left( heta_0
ight)} \, \left| \hat{ heta}^{
m MLE} - heta_0 
ight| \, > \, \sqrt{q_lpha\left(\chi_1^2
ight)} 
ight). \end{array}$$

Using the result from the problem above, we see that the two-sided test of level  $\alpha$  is the same as Wald's test at level  $\alpha$ .

Discussion

**Hide Discussion** 


**Topic:** Unit 4 Hypothesis testing:Lecture 14: Wald's Test, Likelihood Ratio Test, and Implicit Hypothesis Test / 7. Wald's Test in 1 Dimension

Add a Post

**♦** All Posts

Q(alpha)

question posted 8 days ago by **nbourbon** 



Learn About Verified Certificates

© All Rights Reserved