



[Course](#) > [Unit 4 Hypothesis testing](#) > [Homework 7](#) > 4. One-sided Test vs Wald's Test

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

4. One-sided Test vs Wald's Test

In the problems on this page, $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Exp}(\lambda)$, where $\lambda > 0$ is an unknown parameter. In this series of problems, we will compare two tests for the following null and alternative hypotheses:

$$H_0 : \lambda \leq 1$$

$$H_1 : \lambda > 1.$$

MLE and Fisher Information for an Exponential Statistical Model

2/2 points (graded)

What is the MLE $\hat{\lambda}$ for an exponential statistical model?

(Enter **barX_n** for \bar{X}_n .)

Generating Speech Output

1/barX_n



What is the Fisher information $I(\lambda)$ for an exponential statistical model?

$I(\lambda) =$

1/lambda^2



STANDARD NOTATION

Submit

You have used 1 of 4 attempts

✓ Correct (2/2 points)

Test Statistic Based on the MLE for an Exponential Statistical Model

1/1 point (graded)

Assume that the technical conditions hold so that the MLE $\hat{\lambda}_n^{MLE}$ of an exponential statistical model is asymptotically normal. Then it follows that

$$\frac{\sqrt{n}(\hat{\lambda}_n^{MLE} - \lambda)}{g(\hat{\lambda}_n^{MLE})} \xrightarrow[n \rightarrow \infty]{(d)} N(0, 1)$$

where $g(\hat{\lambda}_n^{MLE})$ is an expression that depends on $\hat{\lambda}_n^{MLE}$.

What is $g(\hat{\lambda}_n^{MLE})$?

(Enter **hatlambda** for $\hat{\lambda}_n^{MLE}$.)

$g(\hat{\lambda}_n^{MLE}) =$

hatlambda



Generating Speech Output

RD NOTATION

Submit

You have used 2 of 4 attempts

✓ Correct (1/1 point)

Evaluating the Test Based on the MLE

1/1 point (graded)

Let us define the test statistic

$$T_n = \frac{\sqrt{n}(\hat{\lambda}_n^{MLE} - 1)}{g(\hat{\lambda}_n^{MLE})}$$

where $g(\hat{\lambda}_n^{MLE})$ is the expression from the previous problem.

We define the test $\psi = \mathbf{1}(T_n > \tau)$, where τ is chosen so that ψ is a test at asymptotic level $\alpha = 0.05$. Suppose we observe $\bar{X}_n = 0.83$.

Does the test ψ **reject** or **fail to reject** H_0 on this data set? **Use** $n = 100$.

☐ Fail to reject

☒ Reject



Submit

You have used 1 of 1 attempt

✓ Correct (1/1 point)

Generating Speech Output

Wald's Test

2/2 points (graded)

Recall the test-statistic T_n from the previous problem, and let T_n^{Wald} denote the test-statistic associated to Wald's test for the hypotheses H_0 and H_1 .

Express T_n^{Wald} in terms of T_n .

(Enter **T_n** for T_n .)

$T_n^{Wald} =$ ✓

T_n^2

Which of the following is true about T_n^{Wald} if we assume that $\lambda = 1$?

☐ T_n^{Wald} is distributed as $\mathcal{N}(0, 1)$.

☐ T_n^{Wald} is asymptotically distributed as χ_2^2 .

☐ T_n^{Wald} is distributed as χ_1^2 .

☒ T_n^{Wald} is asymptotically distributed as χ_1^2 .



STANDARD NOTATION

Submit

You have used 1 of 4 attempts

✓ Correct (2/2 points)

Generating Speech Output

Evaluating Wald's Test on a Sample Data Set

1/1 point (graded)

Consider the test $\psi^{Wald} = \mathbf{1}(T_n^{Wald} > \tau)$ where τ is set so that the test ψ^{Wald} has asymptotic level 0.05. Suppose you observe $\bar{X}_n = 0.83$.

Does the test ψ^{Wald} **reject** or **fail to reject** on the given data set? **Use** $n = 100$.

☒ Fail to reject

☐ Reject



Submit

You have used 1 of 1 attempt

✓ Correct (1/1 point)

Discussion

Hide Discussion

Topic: Unit 4 Hypothesis testing:Homework 7 / 4. One-sided Test vs Wald's Test

Add a Post

Show all posts ▼

by recent activity ▼

There are no posts in this topic yet.

✕

Learn About Verified Certificates

© All Rights Reserved

Generating Speech Output