

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Course](#) > [Unit 2 Foundation of Inference](#) > [Lecture 5: Delta Method and Confidence Intervals](#)

10. Confidence Interval for an Exponential Statistical Model

### Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

## 10. Confidence Interval for an Exponential Statistical Model

### Confidence Interval for an Exponential Statistical Model

So you have that lambda is between lambda 1 plus lambda hat plus lambda q alpha over 2 divided by root n,

lambda hat minus lambda q alpha over 2 divided by root n.

I put on my plug-in hat, and there you go.

I have a new confidence interval which is my plug-in confidence interval.

I factor out my lambda hat, and I

get something which is centered about lambda hat--

**much easier.**

### Three solutions

1. The conservative bound:  no a priori way to bound  $\lambda$
2. We can solve for  $\lambda$ :

$$|\hat{\lambda} - \lambda| \leq \frac{q_{\alpha/2}\lambda}{\sqrt{n}} \iff \lambda \left(1 - \frac{q_{\alpha/2}}{\sqrt{n}}\right) \leq \hat{\lambda} \leq \lambda \left(1 + \frac{q_{\alpha/2}}{\sqrt{n}}\right)$$

$$\iff \frac{\hat{\lambda}}{1 + \frac{q_{\alpha/2}}{\sqrt{n}}} \leq \lambda \leq \frac{\hat{\lambda}}{1 - \frac{q_{\alpha/2}}{\sqrt{n}}}$$

It yields

$$\mathcal{I}_{\text{solve}} = \left[ \hat{\lambda} \left(1 + \frac{q_{\alpha/2}}{\sqrt{n}}\right)^{-1}, \hat{\lambda} \left(1 - \frac{q_{\alpha/2}}{\sqrt{n}}\right)^{-1} \right]$$

3. Plug-in yields

$$\mathcal{I}_{\text{plug-in}} = \left[ \hat{\lambda} \left(1 - \frac{q_{\alpha/2}}{\sqrt{n}}\right), \hat{\lambda} \left(1 + \frac{q_{\alpha/2}}{\sqrt{n}}\right) \right]$$



5:30 / 5:30

▶ 1.50x



#### Video

[Download video file](#)

#### Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

### Confidence interval Concept Check

1/1 point (graded)

As in the previous section, let  $X_1, \dots, X_n \stackrel{iid}{\sim} \exp(\lambda)$ . Let

$$\hat{\lambda}_n := \frac{n}{\sum_{i=1}^n X_i}$$

denote an estimator for  $\lambda$ . We know by now that  $\hat{\lambda}_n$  is a **consistent** and **asymptotically normal** estimator for  $\lambda$ .

Recall  $q_{\alpha/2}$  denote the  $1 - \alpha/2$  quantile of a standard Gaussian. By the Delta method:

$$\lambda \in \left[ \hat{\lambda}_n - \frac{q_{\alpha/2}\lambda}{\sqrt{n}}, \hat{\lambda}_n + \frac{q_{\alpha/2}\lambda}{\sqrt{n}} \right] =: \mathcal{I}$$

with probability  $1 - \alpha$ . However,  $\mathcal{I}$  is still **not** a confidence interval for  $\lambda$ .

Why is this the case?

- ☐  $\mathcal{I}$  is actually a confidence interval for  $1/\lambda$ , not  $\lambda$ .
- ☒ The endpoints of  $\mathcal{I}$  depend on the true parameter.
- ☐ A confidence interval is supposed to be random, but  $\mathcal{I}$ , as constructed, is not.
- ☐ As written, the left endpoint of  $\mathcal{I}$  may be larger than the right endpoint of  $\mathcal{I}$ , in which case  $\mathcal{I}$  would not even be a valid interval.



#### Solution:

The **second choice** is correct. The expression for the left and right endpoint of  $\mathcal{I}$  both depend on the true parameter  $\lambda$ . By definition, a confidence interval must be computed only using the data and other known quantities, but not the true parameter, which is unknown. Therefore  $\mathcal{I}$  is not a valid confidence interval.

Now we examine the incorrect choices.

- The first choice ' $\mathcal{I}$  is actually a confidence interval for  $1/\lambda$ , not  $\lambda$ ' is incorrect because, as already discussed,  $\mathcal{I}$  cannot be a confidence in the first place because its endpoints depend on the true parameter.

- The third choice 'A confidence interval is supposed to be random, but  $\mathcal{I}$ , as constructed, is not' is also incorrect. The randomness for  $\mathcal{I}$  comes from  $\hat{\lambda}_n$ , which is random because it depends on the sample. Recall that  $\hat{\lambda}_n$  is the reciprocal of the sample mean.
- The fourth choice 'As written, the left endpoint of  $\mathcal{I}$  may be larger than the right endpoint of  $\mathcal{I}$ , in which case  $\mathcal{I}$  would not be a valid interval' is also incorrect. Since  $q_{\alpha/2}$ ,  $\lambda$ , and  $\sqrt{n}$  are all positive numbers, it follows that

$$\hat{\lambda}_n - \frac{q_{\alpha/2}\lambda}{\sqrt{n}} < \hat{\lambda}_n + \frac{q_{\alpha/2}\lambda}{\sqrt{n}}.$$

Hence,  $\mathcal{I}$  is always a valid interval, just not a valid **confidence** interval.

Submit

You have used 1 of 3 attempts

**i** Answers are displayed within the problem

## Conservative confidence interval for a exponential model

1/1 point (graded)

This problem illustrates the failure of the 'conservative method' for constructing confidence intervals for an exponential statistical model.

As above, let  $X_1, \dots, X_n \stackrel{iid}{\sim} \exp(\lambda)$ . Let

$$\hat{\lambda}_n := \frac{n}{\sum_{i=1}^n X_i}$$

denote an estimator for  $\lambda$ .

Previously, we used the Delta method to show that for  $n$  sufficiently large

$$\lambda \in \left[ \hat{\lambda}_n - \frac{q_{\alpha/2}\lambda}{\sqrt{n}}, \hat{\lambda}_n + \frac{q_{\alpha/2}\lambda}{\sqrt{n}} \right] =: \mathcal{I}$$

where  $q_{\alpha/2}$  is the  $1 - \alpha/2$  quantile of a standard Gaussian.

Given an interval of the form  $\mathcal{I}$ , we can use the "conservative method" to find a confidence interval  $\mathcal{I}_{cons}$  for  $\lambda$  defined by

$$\mathcal{I}_{cons} := \left[ \hat{\lambda}_n - \max_{\lambda \in (0, \infty)} \frac{q_{\alpha/2}\lambda}{\sqrt{n}}, \hat{\lambda}_n + \max_{\lambda \in (0, \infty)} \frac{q_{\alpha/2}\lambda}{\sqrt{n}} \right].$$

Which of the following is  $\mathcal{I}_{cons}$ ?

☒  $(-\infty, \infty)$

☐ the empty interval

☐ the point  $\hat{\lambda}_n$

☐ Cannot be determined, since the exact form of  $\mathcal{I}_{cons}$  will depend on the particular sample.



**Solution:**

Observe that

$$\max_{\lambda \in (0, \infty)} \frac{q_{\alpha/2}\lambda}{\sqrt{n}} = \infty$$

. In this case, we take the max over the interval  $(0, \infty)$ , because a priori,  $\lambda$  can be any number in this interval. Therefore,

$$\mathcal{I}_{cons} = (-\infty, \infty).$$

**Remark:** Although  $(-\infty, \infty)$  is technically still a confidence interval (it even has level 100%!), it is not useful for statistical purposes because such a confidence interval gives no information about the location of the true parameter.

Submit

You have used 1 of 3 attempts

**i** Answers are displayed within the problem

## Explicit Confidence Intervals for an Exponential Statistical Model

6/6 points (graded)

Suppose you observe a sample data set consisting of  $n = 64$  inter-arrival times  $X_1, \dots, X_{64}$  for the subway, measured in minutes. As before, we assume the statistical model that  $X_1, \dots, X_{64} \stackrel{iid}{\sim} \exp(\lambda)$  for some unknown parameter  $\lambda > 0$ . In this data set, you observe that the sample mean is  $\frac{1}{64} \sum_{i=1}^{64} X_i = 7.8$ .

*Additional Instructions: For best results, please adhere to the following guidelines and reminders:*

1. For the upcoming calculations, please truncate  $q_{\alpha/2}$  at 2 decimal places, instead of a more exact value. For example, if  $q_{\alpha/2} = 3.84941$ , use 3.84 instead of 3.85 or 3.849.
2. Input answers truncated at 4 decimal places. For example, if your calculations yield 11.327458, use 11.3274 instead of 11.3275 or 11.32745.
3. You will be computing CIs at **asymptotic level 90%**.

Using the 'solve method' (**refer to the slide 'Three solutions'**), construct a confidence interval  $\mathcal{I}_{solve}$  with asymptotic level 90% for the unknown parameter  $\lambda$ .

[  ✓ Answer: 0.1063408 ,  ✓ Answer: 0.1613875 ]

Using the 'plug-in method'  $\mathcal{I}_{plug-in}$  (refer to the slide 'Three solutions'), construct a confidence interval with asymptotic level 90% for the unknown parameter  $\lambda$ .

[  ✓ Answer: 0.1018453 ,  ✓ Answer: 0.154565 ]

Which interval is narrower?

☐  $\mathcal{I}_{solve}$

☒  $\mathcal{I}_{plug-in}$



Which of these confidence intervals is centered about the sample estimate,  $\hat{\lambda}_n$ ?

☐  $\mathcal{I}_{solve}$

☒  $\mathcal{I}_{plug-in}$

☐ Both

☐ Neither



**Solution:**

The formula for  $\mathcal{I}_{solve}$  at asymptotic level  $1 - \alpha$  is given by

$$\mathcal{I}_{solve} = \left[ \widehat{\lambda}_n \left( 1 + \frac{q_{\alpha/2}}{\sqrt{n}} \right)^{-1}, \widehat{\lambda}_n \left( 1 - \frac{q_{\alpha/2}}{\sqrt{n}} \right)^{-1} \right].$$

We need to construct a confidence interval of (asymptotic) level 90%, so this implies that  $\alpha = 0.1$  and thus  $q_{\alpha/2} = q_{0.05} \approx 1.64$  (consulting a table for the standard Gaussian). Hence, for this data set,

$$\begin{aligned} \mathcal{I}_{solve} &= \left[ \frac{1}{7.8} \left( 1 + \frac{1.64}{\sqrt{64}} \right)^{-1}, \frac{1}{7.8} \left( 1 - \frac{1.64}{\sqrt{64}} \right)^{-1} \right] \\ &\approx [0.1064, 0.1613]. \end{aligned}$$

Next we compute  $\mathcal{I}_{plug-in}$ . The formula is given by

$$\mathcal{I}_{plug-in} = \left[ \widehat{\lambda}_n \left( 1 - \frac{q_{\alpha/2}}{\sqrt{n}} \right), \widehat{\lambda}_n \left( 1 + \frac{q_{\alpha/2}}{\sqrt{n}} \right) \right].$$

Thus for this data set,

$$\begin{aligned} \mathcal{I}_{plug-in} &= \left[ \frac{1}{7.8} \left( 1 - \frac{q_{0.05}}{\sqrt{64}} \right), \frac{1}{7.8} \left( 1 + \frac{q_{0.05}}{\sqrt{64}} \right) \right] \\ &\approx [0.1019, 0.1545]. \end{aligned}$$

Since

$$\begin{aligned} |0.1064 - 0.1613| &= 0.0549 \\ |0.1019 - 0.1545| &= 0.0526, \end{aligned}$$

this implies that  $\mathcal{I}_{plug-in}$  is the **narrower** confidence interval.

Finally,



$$((0.1064 + 0.1613) / 2) * 7.8 = 1.04403$$

$$(0.1019 + 0.1545) / 2) * 7.8 = 0.99996,$$

so we see that  $\mathcal{I}_{plug-in}$  is **centered** about  $\hat{\lambda}_n$ , while  $\mathcal{I}_{solve}$  is not. Alternatively, one can see directly from the formulas that  $\mathcal{I}_{plug-in}$  is always centered about  $\hat{\lambda}_n$  whereas  $\mathcal{I}_{solve}$  is **not** in general.

[Submit](#)

You have used 3 of 5 attempts

**i** Answers are displayed within the problem

## Discussion

[Hide Discussion](#)

**Topic:** Unit 2 Foundation of Inference:Lecture 5: Delta Method and Confidence Intervals / 10. Confidence Interval for an Exponential Statistical Model

[Add a Post](#)

Show all posts ▼

by recent activity ▼

? [Argument for the CI to not be of the form...](#)

1

[Learn About Verified Certificates](#)

© All Rights Reserved