

edX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Unit 0. Course Overview, Syllabus, Guidelines, and Homework on](#)

[Homework 0: Probability and Linear](#)

[Course](#) > [Prerequisites](#)

> [algebra Review](#)

> 6. Probability tables

### Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

## 6. Probability tables

### Gaussian probabilities

4/4 points (graded)

Let  $X \sim N(1, 2.25)$ . As a reminder, the 2.25 here represents the value of  $\sigma^2$ . Using the normal probability table below, compute the following probabilities:

Normal probability table

[Show](#)

$$\mathbf{P}(X > 1) = \boxed{0.5} \quad \checkmark \text{ Answer: } 0.5$$

$$\mathbf{P}(|X - 2| \leq 1) = \boxed{0.4087887802741321} \quad \checkmark \text{ Answer: } 0.4082$$

$$\mathbf{P}(X^2 > 4) = \boxed{0.2752426694951021} \quad \checkmark \text{ Answer: } 0.2752427$$

$$\mathbf{P}(X^2 - 2X - 1 > 0) = \boxed{0.3457785861511602} \quad \checkmark \text{ Answer: } 0.3457786$$

STANDARD NOTATION

### Solution:

First, note that for  $Z \sim \mathcal{N}(0, 1)$ ,  $x > 0$ , we have

$$\mathbf{P}(Z \leq -x) = \mathbf{P}(Z \geq x) = 1 - \mathbf{P}(Z \leq x),$$

and

$$\mathbf{P}(Z \geq x) = 1 - \mathbf{P}(Z \leq x).$$

Moreover, if  $X \sim \mathcal{N}(1, 2.25)$ , we can write it as  $X = 1.5Z + 1$ , where  $Z \sim \mathcal{N}(0, 1)$ . This allows us to reduce all probabilities to the ones that are listed in the table.

In particular,

$$\begin{aligned}\mathbf{P}(X > 1) &= \mathbf{P}(1.5Z + 1 > 1) = \mathbf{P}(1.5Z > 0) \\ &= \mathbf{P}(Z \geq 0) = 1 - \mathbf{P}(Z \leq 0) = 1 - 0.5000 = 0.5000,\end{aligned}$$

$$\begin{aligned}\mathbf{P}(|X - 2| \leq 1) &= \mathbf{P}(-1 \leq (X - 2) \leq 1) = \mathbf{P}(-1 \leq (1.5Z + 1 - 2) \leq 1) \\ &= \mathbf{P}(0 \leq 1.5Z \leq 2) \\ &\simeq \mathbf{P}(0 \leq Z \leq 1.33) \\ &= \mathbf{P}(Z \leq 1.33) - \mathbf{P}(Z \leq 0) \simeq 0.9082 - 0.5000 = 0.4082\end{aligned}$$

$$\begin{aligned}\mathbf{P}(X^2 > 4) &= \mathbf{P}(|X| > 2) = \mathbf{P}(|1.5Z + 1| > 2) \\ &= \mathbf{P}(1.5Z + 1 \leq -2) + \mathbf{P}(1.5Z + 1 \geq 2) \\ &= \mathbf{P}(Z \leq -2) + \mathbf{P}\left(Z \geq \frac{2}{3}\right) \\ &= 1 - \mathbf{P}(Z \leq 2) + 1 - \mathbf{P}\left(Z \leq \frac{2}{3}\right) \\ &\simeq 2 - 0.9772 - 0.7486 = 0.2742\end{aligned}$$

$$\begin{aligned}\mathbf{P}(X^2 - 2X - 1 > 0) &= \mathbf{P}((X - 1)^2 - 2 > 0) = \mathbf{P}(|X - 1| > \sqrt{2}) \\ &= \mathbf{P}(|1.5Z| > \sqrt{2})\end{aligned}$$

$$\begin{aligned} &= \mathbf{P}\left(Z > \frac{\sqrt{2}}{1.5}\right) + \mathbf{P}\left(Z < -\frac{\sqrt{2}}{1.5}\right) \\ &= 2 - 2\mathbf{P}\left(Z < \frac{\sqrt{2}}{1.5}\right) \\ &\simeq 2 - 2(0.8264) \\ &= 0.3472. \end{aligned}$$

Submit

You have used 2 of 3 attempts

**i** Answers are displayed within the problem

## Approximation of Binomial variables

1/1 point (graded)

Using the normal probability table, evaluate approximately  $\mathbf{P}(X > 400)$ , where  $X$  is a binomial random variable with parameters 1000 and .3.

Normal probability table

[Show](#) $\mathbf{P}(X > 400) \simeq$ 

0

✓ Answer: 0.0002

STANDARD NOTATION

**Solution:**

A binomial distribution with parameters  $(n, p)$  has expectation  $np$  and variance  $np(1 - p)$ . Hence, by the Central Limit Theorem, we have

$$\frac{1}{\sqrt{np(1 - p)}}(X - np) \xrightarrow{(D)} Z \sim \mathcal{N}(0, 1).$$

The probability in question can therefore be approximated by

$$\begin{aligned} \mathbf{P}(X > 400) &= \mathbf{P}\left(\frac{1}{\sqrt{1000 \times 0.3 \times 0.7}}(X - 300) > \frac{100}{\sqrt{1000 \times 0.3 \times 0.7}}\right) \\ &\simeq 1 - \mathbf{P}\left(Z \leq \frac{100}{\sqrt{1000 \times 0.3 \times 0.7}}\right) \\ &\simeq 1 - \mathbf{P}(Z \leq 6.90) \\ &\leq 1 - 0.9998 = 0.0002. \end{aligned}$$

Note: This is only an estimate, because the probability table ends here. In fact, the probability is approximately  $7 \times 10^{-12}$ .

You have used 1 of 3 attempts

**i** Answers are displayed within the problem

## Discussion

[Hide Discussion](#)

**Topic:** Unit 0. Course Overview, Syllabus, Guidelines, and Homework on  
**Prerequisites:**Homework 0: Probability and Linear algebra Review / 6. Probability tables

[Add a Post](#)[◀ All Posts](#)

### grader answer doesn't match given solution

discussion posted 7 days ago by anonymous

it's slightly off - just wanted to make sure that a green tick is actually correct - there's been some discrepancies in other courses.

Also - in this course, is it the convention to round up when doing the table lookup? Thanks.

This post is visible to everyone.

[Add a Response](#)

2 responses

**Cool7**

7 days ago

Grader answer looks like not using table lookup, but solution does use table lookup.

I think when we are told to use table, we'd better to use it. Sometimes grader accept both but more often grader only accept answer using table lookup method.

Add a comment

**IvanZinDC**

5 days ago



yeah, basically the table says that anything beyond a certain Z score is the same low probability.

Add a comment

Showing all responses

Add a response:

Preview

Submit

Learn About Verified Certificates

© All Rights Reserved