



Bookmarks

- ▶ Unit 0: Overview
- ▶ Entrance Survey
- ▶ Unit 1: Probability models and axioms
- ▶ Unit 2: Conditioning and independence
- ▶ Unit 3: Counting
- ▶ Unit 4: Discrete random variables
- ▶ Exam 1
- ▶ Unit 5: Continuous random variables
- ▼ **Unit 6: Further topics on random variables**
  - Unit overview
  - Lec. 11: Derived distributions**  
Exercises 11 due Mar 30, 2016 at 23:59 UTC
  - Lec. 12: Sums of independent r.v.'s; Covariance and correlation**  
Exercises 12 due Mar 30, 2016 at 23:59 UTC
  - Lec. 13: Conditional expectation and variance revisited; Sum of a random number of independent r.v.'s**  
Exercises 13 due Mar 30, 2016 at 23:59 UTC
  - Solved problems
  - Additional theoretical material
  - Problem Set 6**  
Problem Set 6 due Mar 30, 2016 at 23:59 UTC
  - Unit summary

Unit 6: Further topics on random variables &gt; Lec. 11: Derived distributions &gt; Lec 11 Derived distributions vertical4



Bookmark

## Exercise: Nonmonotonic functions

(4/4 points)

Suppose that  $\mathbf{X}$  is a continuous random variable and that  $\mathbf{Y} = \mathbf{X}^4$ . Then, for  $y \geq 0$ , we have

$$f_Y(y) = ay^b f_X(-cy^d) + ay^b f_X(cy^d),$$

for some  $a, b, d$ , and some  $c > 0$ . Find  $a, b, c$ , and  $d$ .

 $a =$  

✓ Answer: 0.25

 $b =$  

✓ Answer: -0.75

 $c =$  

✓ Answer: 1

 $d =$  

✓ Answer: 0.25

Answer:

We have, for  $y \geq 0$ ,

$$F_Y(y) = \mathbf{P}(Y \leq y) = \mathbf{P}(X^4 \leq y) = \mathbf{P}(-y^{1/4} \leq X \leq y^{1/4}) = F_X(y^{1/4}) - F_X(-y^{1/4})$$

By differentiating, and using also the chain rule, we obtain

$$f_Y(y) = f_X(y^{1/4}) \cdot \frac{1}{4} \cdot y^{-3/4} + f_X(-y^{1/4}) \cdot \frac{1}{4} \cdot y^{-3/4}.$$

Therefore,  $a = 1/4$ ,  $b = -3/4$ ,  $c = 1$ , and  $d = 1/4$ .

You have used 1 of 2 submissions



© edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

