



15. Worked example: odd periodic

[Course](#) > [Unit 1: Fourier Series](#) > [1. Introduction to Fourier Series](#) > function


Audit Access Expires Jun 24, 2020

You lose all access to this course, including your progress, on Jun 24, 2020.

Upgrade by Jun 7, 2020 to get unlimited access to the course as long as it exists on the site. [**Upgrade now**](#)

15. Worked example: odd periodic function

An odd, periodic example



▶ 7:20 / 7:20

▶ 1.50x 🔊 🗑️ CC 🗨️

Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Find the Fourier series of the 2π -periodic **sawtooth** wave



$$f(t) = t, -\pi < t < \pi.$$




Figure 5: The sawtooth wave.

Solution: The function is odd, therefore $a_n = 0$.

Using our simplified formula for b_n , we find


$$\begin{aligned} b_n &= \frac{2}{\pi} \int_0^\pi t \sin(nt) \, dt \\ &= \frac{2}{\pi} \left[\frac{-t \cos(nt)}{n} \Big|_0^\pi - \int_0^\pi \frac{-\cos(nt)}{n} \, dt \right] \\ &= \frac{2}{\pi} \left[-\frac{\pi(-1)^n}{n} + \frac{\sin(nt)}{n^2} \Big|_0^\pi \right] \\ &= \frac{2(-1)^{n+1}}{n}. \end{aligned}$$

Therefore the Fourier series for the sawtooth wave is



$$\sum_{n=1}^{\infty} \frac{2(-1)^{n+1}}{n} \sin(nt).$$

Graphical intuition



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Practice problem

3/3 points (graded)

Find the Fourier series of the 2π -periodic **triangle wave**, which is defined by

$$T(t) := \begin{cases} t, & \text{if } 0 < t < \pi, \\ -t & \text{if } -\pi < t < 0. \end{cases}$$




Figure 6: The triangle wave.

Note that this function is even, therefore $b_n = 0$.

$$\frac{a_0}{2} =$$

pi/2

✓ Answer: pi/2

$\frac{\pi}{2}$

$$a_n, n \text{ odd} = \boxed{-4/(\pi \cdot n^2)} \quad \checkmark \text{ Answer: } -4/(\pi \cdot n^2)$$

$$-\frac{4}{\pi \cdot n^2}$$

$$a_n, n \text{ even} = \boxed{0} \quad \checkmark \text{ Answer: } 0$$

$$0$$

[FORMULA INPUT HELP](#)

Solution:

The triangle wave is an even function, therefore the $b_n = 0$.

We use the fact that $a_0/2$ is the average value of the function on the interval $-\pi < t < \pi$, which is $\pi/2$.

To compute the rest of the terms, we do so directly from the simplified formulas:

$$a_n = \frac{2}{\pi} \int_0^{\pi} t \cos(nt) dt = \frac{2}{\pi} \left(\frac{(-1)^n - 1}{n^2} \right) = \begin{cases} \frac{-4}{\pi n^2} & n \text{ odd,} \\ 0 & n \text{ even.} \end{cases}$$

Therefore the Fourier series is given by

$$\frac{\pi}{2} - \frac{4}{\pi} \sum_{n \text{ odd}} \frac{\cos(nt)}{n^2}.$$

Submit

You have used 2 of 5 attempts

i Answers are displayed within the problem



15. Worked example: odd periodic function

Hide Discussion

Topic: Unit 1: Fourier Series / 15. Worked example: odd periodic function

Add a Post

Show all posts ▼

by recent activity ▼

- ✓ Why the term a_0 can not be calculated using the one of a_n ? 2
I was using the formula of a_n to compute a_0 by repacking n with 0. However, the result was not correct. Why is the computation of a_0 different from the formula using the ...
- ? 'Best' approximation to function over interval? 8
Prof Matcuk contrasts Taylor series approximating near a point with Fourier series approximating over an interval. In the case of the triangular wave only a few terms are nee...
- 💬 a_n odd 5
I think my answer is correct. Could you check it?
- 💬 Derivative of sawtooth 2
In this video, we calculated the Fourier series for the sawtooth. Jumping ahead I wanted to take the derivative of the series, evaluate it at $t=0$, and see that we get 1. Not havin...

Learn About Verified Certificates

© All Rights Reserved

