

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Unit 0. Course Overview, Syllabus, Guidelines, and Homework on Course](#) > [Prerequisites](#)

[Homework 0: Probability and Linear algebra Review](#)

9. Eigenvalues, Eigenvectors and Determinants(Optional)

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

9. Eigenvalues, Eigenvectors and Determinants(Optional)

Eigenvalues and Eigenvectors of a matrix (Optional)

0 points possible (ungraded)

Let $\mathbf{A} = \begin{pmatrix} 3 & 0 \\ \frac{1}{2} & 2 \end{pmatrix}$, $\mathbf{v} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ and $\mathbf{w} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$.

$\mathbf{A}\mathbf{v} = \lambda_1\mathbf{v}$, where $\lambda_1 =$

✓ Answer: 3 .

 $\mathbf{A}\mathbf{w} = \lambda_2\mathbf{w}$, where $\lambda_2 =$

✓ Answer: 2 .

Therefore, \mathbf{v} is an eigenvector of \mathbf{A} with eigenvalue λ_1 , and \mathbf{w} is an eigenvector of \mathbf{A} with eigenvalue λ_2 .

Solution:

$$\mathbf{A}\mathbf{v} = \begin{pmatrix} 3 & 0 \\ \frac{1}{2} & 2 \end{pmatrix} \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 6 \\ 3 \end{pmatrix} \implies \lambda_1 = 3$$

$$\mathbf{A}\mathbf{w} = \begin{pmatrix} 3 & 0 \\ \frac{1}{2} & 2 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 2 \end{pmatrix} \implies \lambda_2 = 2$$

You have used 1 of 3 attempts

i Answers are displayed within the problem

Geometric Interpretation of Eigenvalues and Eigenvectors (Optional)

0 points possible (ungraded)

Let $\mathbf{A} = \begin{pmatrix} 3 & 0 \\ \frac{1}{2} & 2 \end{pmatrix}$, $\mathbf{v} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ and $\mathbf{w} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$. Recall from the previous exercise that \mathbf{v} and \mathbf{w} are eigenvectors of \mathbf{A} .

Suppose $\mathbf{x} = \mathbf{v} + 2\mathbf{w} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$. Then $\mathbf{Ax} = s\mathbf{v} + t\mathbf{w}$, where:

$s =$  Answer: 3

and

$t =$  Answer: 4 .

In particular, s describes the amount that \mathbf{A} stretches \mathbf{x} in the direction of \mathbf{v} , and $\frac{t}{2}$ (note the "2" in front of \mathbf{w} in \mathbf{x}) describes the amount that \mathbf{A} stretches \mathbf{x} in the direction of \mathbf{w} .

Solution:

We have

$$\begin{aligned}\mathbf{Ax} &= \mathbf{A}(\mathbf{v} + 2\mathbf{w}) \\ &= \mathbf{Av} + 2\mathbf{Aw} \\ &= (3\mathbf{v}) + 2(2\mathbf{w}) \\ &= 3\mathbf{v} + 4\mathbf{w}.\end{aligned}$$

From this, we get $s = 3, t = 4$.

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Determinant and Eigenvalues (optional)

0 points possible (ungraded)

Recall that the **determinant** of a matrix indicates whether it is singular. For 2×2 matrices, it has the formula

$$\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ad - bc$$

but for larger matrices, the formula is more complicated.

What is the determinant of the matrix $\mathbf{A} = \begin{pmatrix} 3 & 0 \\ \frac{1}{2} & 2 \end{pmatrix}$?

6

✓ Answer: 6

On the other hand, what is the product of the eigenvalues λ_1, λ_2 of \mathbf{A} ? (We already computed this in the previous exercises.)

✓ Answer: 6

Solution:

Plugging into the formula directly gives $3 \cdot 2 - 0 \cdot \frac{1}{2} = 6$. On the other hand, the eigenvalues are $\lambda_1 = 3$, $\lambda_2 = 2$, so the product is 6. This is not a coincidence; for general $n \times n$ matrices, the **product of the eigenvalues is always equal to the determinant**.

You have used 1 of 3 attempts

i Answers are displayed within the problem

Trace and Eigenvalues

0 points possible (ungraded)

Recall that the **trace** of a matrix is the sum of the diagonal entries.

What is the trace of the matrix $\mathbf{A} = \begin{pmatrix} 3 & 0 \\ \frac{1}{2} & 2 \end{pmatrix}$?

✓ Answer: 5

On the other hand, what is the sum of the eigenvalues λ_1, λ_2 of \mathbf{A} ? (We already computed this in the previous exercises.)

✓ Answer: 5

Solution:

The diagonal sum is $3 + 2 = 5$. On the other hand, the eigenvalues are $\lambda_1 = 3, \lambda_2 = 2$, so the sum is 5. Just like the determinant, this is also not a coincidence. For general $n \times n$ matrices, the **sum of the eigenvalues is always equal to the trace of the matrix**.

You have used 1 of 3 attempts

i Answers are displayed within the problem

Nullspace (Optional)

0 points possible (ungraded)

If a (nonzero) vector is in the nullspace of a square matrix \mathbf{A} , is it an eigenvector of \mathbf{A} ?

✓ Answer: yes

Which of the following are equivalent to the statement that 0 is an eigenvalue for a given square matrix \mathbf{A} ? (Choose all that apply.)

☒ There exists a nonzero solution to $\mathbf{A}\mathbf{v} = \mathbf{0}$. ✓

☒ $\det(\mathbf{A}) = 0$ ✓☐ $\det(\mathbf{A}) \neq 0$ ☐ $\text{NS}(\mathbf{A}) = \mathbf{0}$ ☒ $\text{NS}(\mathbf{A}) \neq \mathbf{0}$ ✓**Solution:**

- If a vector \mathbf{v} is in the nullspace of \mathbf{A} , then $\mathbf{A}\mathbf{v} = \mathbf{0} = (0)\mathbf{v}$. So it is an eigenvector of \mathbf{A} associated to the eigenvalue 0.
- If 0 is an eigenvalue for a matrix \mathbf{A} , then by definition, there exists a nonzero solution to $\mathbf{A}\mathbf{v} = \mathbf{0}$; that is, $\text{NS}(\mathbf{A}) \neq \mathbf{0}$, and this only happens if and only if $\det(\mathbf{A}) = 0$.

You have used 1 of 3 attempts

i Answers are displayed within the problem

Discussion

Topic: Unit 0. Course Overview, Syllabus, Guidelines, and Homework on

Prerequisites:Homework 0: Probability and Linear algebra Review / 9. Eigenvalues, Eigenvectors and Determinants(Optional)

Add a Post

Show all posts ▼

by recent activity ▼

- ✓ [STAFF] Just curious 7
Is there any chance MITx will offer 18.06x? and/or 18.02x? Just wondering. I believe these subjects will really complement the MicroMasters Program.
- 💬 Boosting grade 3
If I complete the ungraded, will this contribute to my score, if the graded answers are low?
- ? Request extension to deadline 2
For those of us who are finishing the last optional but important week of the probability course these deadlines are tough.Quiz 0 is substantial. Pers...
- 💬 Sources 7
Could somebody recommend any good source to learn all this stuff? Preferably with detailed explanations of what is going on on each step of calcul...
- 💬 NS(A) notation in Nullspace (Optional.) 2

Learn About Verified Certificates

© All Rights Reserved