**Data Analysis: Statistical Modeling and Computation in Applications** 

<u>Help</u>

sandipan\_dey ~

<u>Course</u>

**Progress** 

<u>Dates</u>

**Discussion** 

Resources







## 5. Statistics of random walk

☐ Bookmark this page

Exercises due Nov 10, 2021 17:29 IST Completed

To compute the basic statistics (mean, variance, covariance) of the random walk, it is useful to write  $X_t$  as a sum of the perturbations  $\{W_h\}_{h=1}^t$  that accumulate over time:

$$egin{array}{ll} X_t &= X_{t-1} + W_t \ &= \left[ X_{t-2} + W_{t-1} 
ight] + W_t \ &dots \ &dots \ &= X_0 + \sum_{h=1}^t W_h. \end{array}$$

Similarly, for the random walk with drift  $Y_t$  we have:

$$egin{aligned} Y_t &= \delta + Y_{t-1} + W_t \ &= \delta + \left[\delta + Y_{t-2} + W_{t-1}
ight] + W_t \ &dots \ &= \delta \cdot t + Y_0 + \sum_{h=1}^t W_h. \end{aligned}$$

Using these representations we can find the marginal mean function, the covariance function and the autocorrelation function:

$$egin{aligned} \mu_X\left(t
ight) &= \mathbf{E}\left[X_t
ight] \ &= \mathbf{E}\left[X_0 + \sum_{h=1}^t W_h
ight] \ &= \mathbf{E}\left[X_0
ight] \end{aligned}$$

since  $W_t$  is white noise and has mean zero.

$$egin{aligned} \sigma_X^2\left(t
ight) &= \mathsf{Var}\left(X_t
ight) \ &= \mathsf{Var}\Big(X_0 + \sum_{h=1}^t W_h\Big) \ &= \mathsf{Var}\left(X_0
ight) + \sum_{h=1}^t \left[2\mathsf{Cov}\left(X_0, W_h
ight) + \mathsf{Var}\left(W_h
ight)
ight] + 2\sum_{1 \leq h < j \leq t} \mathsf{Cov}\left(W_h, W_j
ight) \ &= \mathsf{Var}\left(X_0
ight) + t \cdot \sigma_W^2 \end{aligned}$$

since  $W_h$  is uncorrelated with  $X_0$  and with  $W_j$  for j 
eq h.

$$egin{aligned} \gamma_X\left(s,t
ight) &= \mathsf{Cov}\left(X_s,X_t
ight) \ &= \mathsf{Cov}\Big(X_0 + \sum_{h=1}^s W_h,\; X_0 + \sum_{h=1}^t W_h\Big) \ &= \mathsf{Var}\left(X_0
ight) + \sum_{h=1}^{\min(s,t)} \mathsf{Var}\left(W_h
ight) \ &= \mathsf{Var}\left(X_0
ight) + \min\left(s,t
ight) \cdot \sigma_W^2 \end{aligned}$$

since  $W_h$  is uncorrelated with  $X_0$  and with  $W_j$  for j 
eq h.

Note that typically  $X_0$  is assumed to be deterministic, thus  $\mathsf{Var}(X_0) = 0$ .

#### Statistics of random walk

3/3 points (graded)

Let X be a random walk with random perturbations that have  $\sigma_W^2=1$ .

Compute  $\gamma_X$  (5, 10).

5 **✓ Answer:** 5

Compute  $\gamma_X$  (10, 15).

10 **Answer:** 10

Does the autocovariance function  $\gamma_{X}\left(s,t
ight)$  only depend on time gap |s-t|?

True





#### **Solution:**

From the above calculations,  $\gamma_X(5,10) = \min(5,10) \cdot 1 = 5$  and  $\gamma_X(10,15) = 10$ . Therefore, the autocovariance of a random walk is not stationary since the gap between time stamps in these two computations is the same but the covariances are different.

Submit

You have used 1 of 3 attempts

**1** Answers are displayed within the problem

#### Statistics of random walk with drift

As an exercise, repeat the calculation for the marginal mean function  $\mu_Y(t) = \mathbf{E}[Y_t]$ , the marginal variance function  $\sigma_Y^2(t) = \mathsf{Var}(Y_t)$  and the autocovariance function  $\gamma_Y(s,t) = \mathsf{Cov}(Y_s,Y_t)$  of the random walk with drift  $Y_t$ .

#### Computation of marginal mean, marginal variance, and autocovariance function

From the expression above we have

$$egin{aligned} \mu_{Y}\left(t
ight) &= \mathbf{E}\left[\delta t + Y_{0} + \sum_{h=1}^{t}W_{h}
ight] \ &= \delta t + \mathbf{E}\left[Y_{0}
ight] \end{aligned}$$

by linearity of expectation and zero mean property of white noise.

$$\mathsf{Var}\left(Y_{t}
ight) = \mathsf{Var}\Big(Y_{0} + \delta t + \sum_{t=1}^{t} W_{t}\Big)$$

$$= \mathsf{Var}\Big(Y_0 + \sum_{h=1}^t W_t\Big)$$

$$=\mathsf{Var}\left(Y_{0}
ight)+t\sigma_{W}^{2}$$

by property of the variance (that a shift of a distribution does not change the spread of the distribution) and assumptions about white noise.

Similarly,  $\mathsf{Cor}\left(Y_t,Y_s\right) = \mathsf{Var}\left(Y_0\right) + \min\left(t,s\right)\sigma_W^2$  .

<u>Hide</u>

## Random walk and stationarity

2/2 points (graded)

False

Is the random walk with drift process stationary?

True			



Is the random walk process stationary?

\_\_\_\_\_ True





## **Solution:**

False. The random walk is not stationary because the variance is growing with time and the autocovariance depends on the smallest of the two time stamps rather than on the difference.

Submit

You have used 1 of 1 attempt

Answers are displayed within the problem

## Differencing random walk

2/2 points (graded)

Consider the first difference  $abla Y_t$  of a random walk  $Y_t$  with drift.

Calculate the marginal mean function  $\mu_{\nabla Y}(t)$ , the marginal variance function  $\sigma_{\nabla Y}^2(t)$  and the autocovariance function  $\gamma_{\nabla Y}(s,t)$  of  $\nabla Y_t$ .

Select all correct statements.

 $igcup 
abla Y_t$  is random walk

 $igvee VY_t$  is white noise plus constant ( $\delta + W_t$ )

 $ightharpoonup \mu_{
abla Y}\left( t
ight)$  is constant

 $ec{arphi} \; \sigma^2_{
abla Y} \left( t 
ight)$  is constant

 $igwedge \gamma_{
abla Y}\left( s,t
ight) =0$ 

 $\bigcap$   $\gamma_{
abla Y}\left(s,t
ight)$  is constant, but not necessarily 0



Is  $abla Y_t$  a stationary time series?



True





#### **Solution:**

The first difference of the random walk is  $\nabla Y_t = Y_t - Y_{t-1} = \delta + W_t$  white noise. Therefore  $\mu_{\nabla Y}(t) = \delta$  is constant,  $\sigma^2_{\nabla Y}(t) = \sigma^2_W$  is constant, and  $\gamma_{\nabla Y}(s,t) = 0$  for  $s \neq t$ . Yes, a shifted white noise  $\{\delta + W_t\}$  is a stationary time series.

Submit

You have used 1 of 3 attempts

**1** Answers are displayed within the problem

## (Optional) Estimations of random walk model

0 points possible (ungraded)

If  $X_t$  is a random walk, would our estimators  $\hat{\sigma}_X^2\left(1\right)$  and  $\hat{\gamma}_X\left(h\right)$  be consistent for  $\sigma_X^2$  and  $\gamma_X\left(1,h\right)$ ?

True




False




### **Solution:**

No, consistency of the estimated acf function requires stationarity which random walk does not have.

We can illustrate this with a plot of an estimated acf:





© All Rights Reserved



# edX

**About** 

**Affiliates** 

edX for Business

Open edX

Careers

<u>News</u>

# Legal

Terms of Service & Honor Code

Privacy Policy

**Accessibility Policy** 

**Trademark Policy** 

<u>Sitemap</u>

## **Connect**

**Blog** 

**Contact Us** 

Help Center

Media Kit

**Donate** 















© 2021 edX Inc. All rights reserved.

深圳市恒宇博科技有限公司 <u>粤ICP备17044299号-2</u>