EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the Privacy Policy.





Homework 4: TV distance, KL-

Course > Unit 3 Methods of Estimation > Divergence, and Introduction to MLE > 4. Maximum likelihood estimators

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019. Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. **Upgrade now**

4. Maximum likelihood estimators **Instructions:**

Let X_1, \ldots, X_n be n i.i.d. random variables with pdf f_{θ} , where θ is an unknown parameter.

For each of the following questions, compute the likelihood function on paper and then find the maximum likelihood estimator for θ .

To encourage you to do the computations carefully rather than eliminate choices, you will only be given 1 or 2 attempts per question.

(a)

1/1 point (graded)

Compute the likelihood function and the maximum likelihood estimator for $\, heta\,$ for

$$f_{ heta}\left(x
ight)= au heta^{ au}x^{-\left(au+1
ight)}\mathbf{1}\left(x\geq heta
ight),\quad heta>0,$$

where $\tau > 0$ is a known constant.

max	X_i
	$-\iota$











Solution:

The likelihood function is

$$L = au^n heta^{n au} \prod_i X_i^{-(au+1)} \mathbf{1}\{\min_i X_i \geq heta\}$$

For $heta \leq \min_i X_i$, the log-likelihood function is

$$l=n\ln au+n au\ln heta-(au+1)\sum_{i=1}\ln X_i$$

Take the derivative with respect to θ :

$$\frac{\partial l}{\partial \theta} = \frac{n\tau}{\theta} > 0.$$

Thus, L is an increasing function on $(0,\min_i X_i]$, and is 0 for $\theta > \min_i X_i$. Therefore,

$$\hat{ heta} = \min_i X_i$$

Submit

You have used 1 of 2 attempts

• Answers are displayed within the problem

(b)

3/3 points (graded)

Compute the likelihood function and the maximum likelihood estimator for $\, heta\,$ for

$$f_{ heta}\left(x
ight) = \sqrt{ heta}x^{\sqrt{ heta}-1}\mathbf{1}\left(0 < x < 1
ight), \quad heta > 0.$$

You will find that the maximum likelihood estimator for θ is of the form

$$\hat{ heta}^{ ext{MLE}} \ = \ c_1 n^{c_2} \left(\sum_{i=1}^n \ln X_i
ight)^{c_3}.$$

Enter the numbers c_1, c_2, c_3 below.

Answer: 1

$$c_2 =$$
 2 $ightharpoonup$ Answer: 2

$$c_3 = ig|$$
 -2 $ig|$ Answer: -2

STANDARD NOTATION

Solution:

The likelihood function is

$$L= heta^{n/2}\prod_i X_i^{\sqrt{ heta}-1} \mathbf{1}\{0\leq X_i\leq 1\}.$$

The log-likelihood function is

$$l = rac{n}{2} {\ln heta} + (\sqrt{ heta} - 1) \sum_i {\ln X_i}.$$

Take the derivative with respect to θ and set it to 0:

$$rac{\partial l}{\partial heta} = rac{n}{2 heta} + rac{1}{2 heta^{1/2}} \sum_i \ln X_i = 0.$$

Then we get

$$\hat{ heta} = rac{n^2}{\left(\sum \ln X_i
ight)^2}.$$

Submit

You have used 2 of 3 attempts

1 Answers are displayed within the problem

(c)

4/4 points (graded)

Compute the likelihood function and the maximum likelihood estimator for $\, heta$

$$f_{ heta}\left(x
ight)= heta au x^{ au-1}\exp\{- heta x^{ au}\}\mathbf{1}\left(x\geq0
ight),\quad heta>0,$$

where $\tau > 0$ is a known constant.

You will find that the maximum likelihood estimator for $\, heta\,$ is of the form

$$\hat{ heta}^{
m MLE} \; = \; c_1 n^{c_2} \left(\sum_{i=1}^n X_i^{c_3}
ight)^{c_4}.$$

Enter the c_1, c_2, c_3, c_4 in terms of τ if applicable.

(Enter **tau** for τ .)

$$c_1 = egin{bmatrix} 1 & & & \\ 1 & & & \\ 1 & & & \end{bmatrix}$$
 \checkmark Answer: 1

$$c_2=egin{bmatrix} 1 & & & \\ \hline 1 & & & \\ \hline 1 & & & \\ \end{bmatrix}$$
 \checkmark Answer: 1

$$c_3=egin{pmatrix} auu & ag{Answer: tau} \ ag{Answer: tau} \ ag{C}_4= egin{pmatrix} -1 & ag{Answer: -1} \ ag{STANDARD NOTATION} \ ag{STANDARD NOTATION} \ ag{Answer: -1} \ ag{Answer$$

Solution:

The likelihood function is

$$L= heta^n au^n\prod_i X_i^{ au-1}\exp\{- heta\sum_i X_i^ au\}\mathbf{1}\{X_i\geq 0\}.$$

The log-likelihood function is

$$l = n \ln heta + n \ln au + (au - 1) \sum_i \ln X_i - heta \sum_i X_i^ au.$$

Take the derivative with respect to $\, heta\,$ and set it to $\,0\,$

$$rac{\partial l}{\partial heta} = rac{n}{ heta} - \sum_i X_i^{ au} = 0,$$

we get

$$\hat{ heta} = rac{n}{\sum_i X_i^ au}.$$

Submit

You have used 1 of 3 attempts

1 Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 3 Methods of Estimation: Homework 4: TV distance, KL-Divergence, and Introduction to MLE / 4. Maximum likelihood estimators

Add a Post

≺ All Posts Hint: you can't simply sum up the X_i's discussion posted 7 days ago by **yiuming h** I totally missed that we were working with random variables when doing the likelihood calculations, and kept getting the wrong answers. Thank you for this exercise, I think I understand likelihood a lot better now as a result This post is visible to everyone. 0 responses Preview