



You are taking "[Exam \(Timed, No Correctness Feedback\)](#)," as a timed exam. [Show more](#)

End My Exam

44:55:45



< Previous



## 9. Hikers in a canyon continued

🔖 Bookmark this page



Calculator



Hide Notes


Lecture due Aug 4, 2021 20:30 IST   Completed



Apply

Hikers in a canyon continued

To explain the previous question, let's consider the level curves of  $f(x,y) = x^2 + y^2$  shown below.




By starting at  $(-1, 1)$  and moving in the positive  $x$ -direction, we can see that Hiker 1 is moving from near the level curve  $x^2 + y^2 = 2$  towards the level curve  $x^2 + y^2 = 1$ . This indicates that the hiker is moving towards a lower elevation, and therefore, the hiker is moving downhill.

You can check the elevation on the level curves below. Use the cursor to hover over a point on the plot of level curves. You will see the read out of the x position, y position, and z value below the plot.

► Level Curves

Equation 1   ▾

$z = f(x,y) = x^2 + y^2$



Calculator

Hide Notes

PLEASE RATE THIS MATHLET  
(Use a one star to five star rating scale.)

RESULTS

<input type="radio"/> ☆	1%
<input type="radio"/> ☆ ☆	0%
<input type="radio"/> ☆ ☆ ☆	5%
<input type="radio"/> ☆ ☆ ☆ ☆	19%
<input checked="" type="radio"/> ☆ ☆ ☆ ☆ ☆	75%

Submit

Results gathered from 841 respondents.

FEEDBACK  
Your response has been recorded

Hiker 2

1/1 point (graded)  
Recall that Hiker 1 started at  $(-1, 1)$  and moved downhill when going in the positive  $x$ -direction. Now let's consider Hiker 2. This hiker starts at the point  $(-2, 0)$  and moves in the positive  $x$ -direction. In this case, Hiker 2 is also moving downhill. Between Hiker 1 and Hiker 2, which hiker is going downhill more steeply? (This question is graded.)

☐ Hiker 1

☒ Hiker 2

☐ The steepness is the same for both.



Solution:

Hiker 2 is moving downhill more steeply. If we consider the level curves, we can see that Hiker 2 has to move a shorter distance than Hiker 1 to go from one level curve to the next.

Submit    You have used 2 of 2 attempts

Answers are displayed within the problem

9. Hikers in a canyon continued

Hide Discussion

Topic: Unit 1: Functions of two variables / 9. Hikers in a canyon continued




Add a Post

Show all posts

by recent activity

[Slope in a paraboloid?](#)

Hello. I'm having so much fun learning these concepts. I would just like know if this question could be solve

<input checked="" type="checkbox"/> <a href="#">mathlet not loading...</a> <a href="#">mathlet above and also previous ones are not loading. anyone an help?</a>	2
 <a href="#">The plot of the level curves helps</a> <a href="#">you can see how z values change from hiker 2's position compared to hiker 1</a>	1
 <a href="#">Sliding down a parabola</a>	1
 <a href="#">steepest descent</a> <a href="#">Hello everyone. The problem of the steepest descent will be clear when gradient and directional derivative will be introduced. Anyw...</a>	4



## edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

## Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)

## Connect

- [Blog](#)
- [Contact Us](#)
- [Help Center](#)
- [Media Kit](#)
- [Donate](#)



© 2021 edX Inc. All rights reserved.

深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)