



[Lecture 13: Chi Squared Distribution](#).

[Course](#) > [Unit 4 Hypothesis testing](#) > [T-Test](#)

> 10. The Student's T Test (T Test)


Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

10. The Student's T Test (T Test)

The T Test - One Sample, Two-Sided



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)


[Download Text \(.txt\) file](#)

Video note: In the videos on this page and on Slides 16 and 17, the test statistic T_n is missing a scaling factor. The correct equation for T_n is

$$T_n = \frac{\sqrt{n}(\bar{X}_n - \mu_0)}{\sqrt{\tilde{S}_n}}.$$

The unfilled and typed slides have both been corrected.

One Sample, One-Sided T Test



Video[Download video file](#)**Transcripts**[Download SubRip \(.srt\) file](#)[Download Text \(.txt\) file](#)**Concept Check: Student's T Distribution**

3/3 points (graded)

Consider the statistic

$$T_n := \sqrt{n} \left(\frac{\bar{X}_n - \mu}{\sqrt{\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2}} \right).$$

For all $n \geq 2$, the distribution of T_n is a standard Gaussian $\mathcal{N}(0, 1)$.☐ True☒ FalseAs $n \rightarrow \infty$, what does

$$\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2$$

converge to...

☐ The number μ (weakly)

☒ The number σ^2 (weakly)

☐ The distribution $\mathcal{N}(0, 1)$

☐ The distribution χ_{n-1}^2



As $n \rightarrow \infty$, the statistic T_n converges in distribution to

☒ $\mathcal{N}(0, 1)$

☐ $\mathcal{N}(\mu, \sigma^2)$

☐ χ_{n-1}^2

☐ χ_n^2



Solution:

The definition of the student's T distribution with $n - 1$ degrees of freedom is that it is given by the distribution of $\frac{Z}{\sqrt{V/(n-1)}}$ where $Z \sim \mathcal{N}(0, 1)$, $V \sim \chi_{n-1}^2$ and Z and V are independent. Since we are dividing by V , a χ^2 random variable, then T_n will not have the same distribution as $\mathcal{N}(0, 1)$ for all $n \geq 2$.

By the law of large numbers and Slutsky's lemma,

$$\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2 = \frac{n}{n-1} \left[\left(\frac{1}{n} \sum_{i=1}^n X_i^2 \right) - (\bar{X}_n)^2 \right] \rightarrow \sigma^2$$

in probability.

By the central limit theorem,

$$\sqrt{n} \left(\frac{\bar{X}_n - \mu}{\sigma} \right) \rightarrow \mathcal{N}(0, 1).$$

Hence, by the law of large numbers and Slutsky's theorem,

$$\sqrt{n} \left(\frac{\bar{X}_n - \mu}{\sqrt{\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2}} \right) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}(0, 1).$$

Submit

You have used 2 of 3 attempts

i Answers are displayed within the problem

Quantiles of the T Distribution:

The $(1 - \alpha)$ -quantile of the t_{n-1} (corresponding to a one-sided test with statistic T_n) can be computed using standard computational tools such as R. One can also find online tables for the quantiles via a simple Google search, which yields results such as [this](#), [this](#), and [this](#).

As a reminder, in this class the $(1 - \alpha)$ quantile of the distribution of a random variable T is the number q_α such that

$$P(T \leq q_\alpha) = 1 - \alpha.$$

Concept Check: T Test

1/1 point (graded)

Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu^*, \sigma^2)$ for some unknown $\mu^* \in \mathbb{R}$ and $\sigma^2 > 0$. You want to decide between the following null and alternative hypotheses on the mean of X_1, \dots, X_n :

$$H_0 : \mu^* = 0$$

$$H_1 : \mu^* \neq 0.$$

To do so, you define the student's T statistic

$$T_n = \sqrt{n} \frac{\bar{X}_n}{\sqrt{\tilde{S}_n}}$$

where

$$\tilde{S}_n = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2$$

is the unbiased sample variance.

The student's T test of level α is specified by

$$\psi_\alpha = \mathbf{1}(|T_n| > q_{\alpha/2})$$

where $q_{\alpha/2}$ is the unique number such that $P(T_n < q_{\alpha/2}) = 1 - \frac{\alpha}{2}$.

Which of the following are true about the student's T test? (Choose all that apply.)

☐ The statistic T_n is distributed as a standard Gaussian.

☒ The test requires the data X_1, \dots, X_n to be Gaussian.

☒ The distribution of T_n is pivotal, *i.e.*, its quantiles may be found in tables.

☒ The test is non-asymptotic. That is, for any fixed n , we can compute the level of our test rather than the *asymptotic* level.



Solution:


We examine the choices in order.

- The first choice is incorrect. Due to the fact that T_n has the sample variance \hat{S}_n in the denominator and not the *true* variance σ^2 , the statistic T_n will **not** be standard Gaussian.
- The second choice is correct. It is a key assumption that the data is Gaussian. Otherwise, the test statistic T_n will not necessarily follow the student's T distribution and, hence, may not even be pivotal.
- The third choice is correct. For any fixed n , we may find the quantiles of the student's T distribution in tables. Since the distribution does not depend on the value of the true parameter, the test statistic T_n is indeed pivotal.
- The last choice is also correct. As stated in the previous bullet, for any fixed n , the quantiles of the student's T distribution may be found in tables. Hence, we can find the non-asymptotic level of this test.

Remark: Assuming the data is Gaussian, the student's T test is useful in situations where the sample size is not very large, since the level may be precisely quantified even for small n .

Submit

You have used 1 of 2 attempts

 Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 4 Hypothesis testing:Lecture 13: Chi Squared Distribution, T-Test / 10. The Student's T Test (T Test)

Add a Post

Show all posts ▼

by recent activity ▼

There are no posts in this topic yet.

✕

Learn About Verified Certificates

© All Rights Reserved