



You are taking "[Exam \(Timed, No Correctness Feedback\)](#)," as a timed exam. [Show more](#)

End My Exam

26:25:08



< Previous



Next >

6. Types of critical points

Bookmark this page



Calculator




Hide Notes



Explore

Types of critical points

[Start of transcript. Skip to the end.](#)



[PROFESSOR]:: Let me point out to you immediately that there's more than maximum and minimum. So, remember, we saw the example of $x^2 + y^2$ squared that has a critical point, but the critical point is obviously a minimum. And of course it could be a local minimum because it could be that, if you have




Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)
[Download Text \(.txt\) file](#)


Just as we saw in single-variable calculus, the derivative equaling zero is a necessary but not sufficient condition for finding local extrema. This means that a critical point (where $f'(x) = 0$) might be a local max or min but it might not. For example, $f(x) = x^3$ has a critical point at $x = 0$. But the function is monotonically increasing everywhere and therefore $x = 0$ is not a local max or min.



Similar behavior can occur for functions of multiple variables. There are three types of critical points for a function of **2** variables.


Local minimum

A point whose value is smaller than any nearby point.



Local maximum

A point whose value is larger than any nearby point.




Calculator

Hide Notes

Saddle point

A point that is neither a minimum or a maximum, but can look like either depending on the direction you look at it from.



Example 6.1 Consider the function

$$f(x,y) = x^2 - y^2.$$

(4.31)

The partial derivatives are


$$f_x(x,y) = 2x \text{ and } f_y(x,y) = -2y$$

(4.32)


which means $(x,y) = (0,0)$ is a critical point of $f(x,y)$. However, if we inspect the graph of f (below), we see that $(0,0)$ is a **saddle point**.


In the x -direction along the curve $f(x,0) = x^2$, the function has a local minimum. (See the orange curve in the interactive image below.) However, in the y -direction along the curve $f(0,y) = -y^2$, the curve has a local maximum. (See the blue curve in the interactive image below). Thus $(0,0)$ is neither a local max nor a local min.

► SADDLE POINT



The gradient field for $f(x,y) = x^2 - y^2$ is shown below on the left, while the level curves are shown below on the right.





6. Types of critical points

Hide Discussion

Topic: Unit 3: Optimization / 6. Types of critical points

Add a Post

Show all posts ▾by recent activity ▾

[\[staff\] Possible error](#)

[In the example 6.1, I think the orange curve is for \$f\(x,0\)=x^2\$ and the blue curve is for \$f\(0,y\)=-y^2\$, which is contrary to what the text i...](#)

3 ▾

[Taxonomy of interesting points?](#)

[Community TA](#)

18 ▾



edX

[About](#)

[Affiliates](#)

[edX for Business](#)

Calculator

Hide Notes

[Open edX](#)
[Careers](#)
[News](#)

Legal

[Terms of Service & Honor Code](#)
[Privacy Policy](#)
[Accessibility Policy](#)
[Trademark Policy](#)
[Sitemap](#)

Connect

[Blog](#)
[Contact Us](#)
[Help Center](#)
[Media Kit](#)
[Donate](#)



© 2021 edX Inc. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)