



Course > Section... > 2.5 Ma... > 2.5.3 E...


2.5.3 Exploratory Quiz: Rates of Predation and Synthesis

🔖 Bookmark this page

Question 1

1/1 point (graded)

In the last video, Ethan explained how trophic levels work, with only about 10% of the energy from each level reaching the next. In particular, only about 10% of the energy at the level of the sardines (the primary consumer) reaches the marlin (the secondary consumer).



[View Larger Image](#)
[Image Description](#)



But Ethan misspoke in the last part when he said we might expect d , the synthesis rate for marlin, to be 10 times bigger than b , the predation rate on sardines. What might we expect instead about the relative sizes of b and d ? Why?

$$\frac{dS}{dt} = aS - bSM$$

$$\frac{dM}{dt} = -cM + dSM.$$

(Recall from earlier sections that b is the predation rate of marlin on the sardine, and the interaction term bSM represents roughly the amount of sardines marlin are eating per unit time. On the other hand, d is the synthesis rate of the marlin, with dSM representing roughly the increase in marlin population per unit time due to their consumption of sardine.)

- ☐ $b \approx d$
- ☐ d about 100 times greater than b
- ☐ b is about 100 times greater than d
- ☒ b is about 10 times greater than d ✓
- ☐ None of the above.

Explanation

Since only about 10% of the energy in the sardine reaches the marlin, this means if the marlin consume bSM sardines, only about 10% of this energy contributes to the growth of the marlin population. So we expect dSM to be approximately equal to 10% of bSM , which means that d is about 10% of b , or b is about 10 times greater than d .

Submit

i Answers are displayed within the problem

Learn About Verified Certificates

© All Rights Reserved



English ▼

© 2012–2018 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open edX logos are registered trademarks or trademarks of edX Inc. | 粤ICP备17044299号-2

POWERED BY
OPENedX®

