






< Previous







Next >

8.8.1 Forward Euler method

 Bookmark this page

M02.4

M02.7

We now consider our first numerical method for solving an IVP, the forward Euler method. The key concept is that from the governing differential equation, given some current state \underline{v} at time t we can always calculate the rate of change from the model differential equation $\underline{du}/\underline{dt} = \underline{f}(\underline{v}, t)$. Since we know the initial condition, we can then start from it, calculate the $\underline{du}/\underline{dt}(t_I) = \underline{f}(\underline{u}_I, t_I)$ and then use that slope to extrapolate the solution to time $t^1 = t_I + \Delta t$. This is shown graphically in Figure 8.13. Mathematically, this gives

$$\underline{v}^1 = \underline{u}(t_I) + \Delta t \underline{f}(\underline{u}(t_I), t_I).$$

(8.56)

© All Rights Reserved




Figure 8.13: Forward Euler method

Now, to find \underline{v}^2 , we can use the same idea to calculate the rate of change from $\underline{f}(\underline{v}^1, t^1)$ and extrapolate again,

$$\underline{v}^2 = \underline{v}^1 + \Delta t \underline{f}(\underline{v}^1, t^1)$$

(8.57)

Legal

[Terms of Service & Honor Code](#)
[Privacy Policy](#)
[Accessibility Policy](#)
[Trademark Policy](#)
[Sitemap](#)

$$\underline{v}^0 = \underline{u}_I$$

(8.58)

$$\underline{v}^{n+1} = \underline{v}^n + \Delta t \underline{f}(\underline{v}^n, t^n) \quad \text{for} \quad n \geq 0,$$

(8.59)

Connect

Using the Forward Euler method

[Blog](#)

[Contact Us](#)

[Help Center](#)

[Security](#)

[Media Kit](#)



INSTRUCTOR: So how do we approximate these values of v in our numerical method? We're going to look at probably what is the simplest

Discussions

All posts sorted by recent activity


© 2023 edX LLC. All rights reserved.

[Previous](#)

[Next](#)

深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)

Video

 [Download video file](#)

Transcripts