<u>Notes</u>

Discussion

<u>Course</u>

<u>Dates</u>

<u>Help</u>

sandipan_dey ~

Next >

<u>Calendar</u>

☆ Course / Unit 2: Geometry of Derivatives / Lecture 6: Gradients



You are taking "Exam (Timed, No Correctness Feedback)" as a timed exam. Show more

End My Exam

Previous

43:56:30





<u>Progress</u>

☐ Bookmark this page

Lecture due Aug 18, 2021 20:30 IST Completed




Synthesize

Guess the direction of the gradient

1/1 point (graded)

Consider the paraboloid $f\left(x,y
ight)=x^2+y^2$ whose level curves are concentric circles.



Which direction does abla f(-1,1) point?

diagonally up and right

diagonally up and left


diagonally down and right

diagonally down and left



Solution:

The point (-1,1) is in the second quadrant on a concentric circle corresponding to the level curve of height 2.



The gradient is normal to the level curve, therefore the two possible options are diagonally up and to the left, and diagonally down and to the right. Because the level curves are increasing as you move up and to the left, that is the direction of steepest increase.

Submit

You have used 1 of 2 attempts

1 Answers are displayed within the problem

Estimate the size of the gradient

1/1 point (graded) Estimate $|\nabla f(-1,1)|$.

pprox 1/3

pprox 1

 \bigcirc ≈ 3



Solution:

Note that $abla f(-1,1) = \langle -2,2
angle$, which has length $= \sqrt{8} pprox 3$.

Submit

You have used 1 of 2 attempts

1 Answers are displayed within the problem

Connection between gradient and 3D graph



0:00 / 0:00 66 ▶ 2.0x X

Start of transcript. Skip to the end.

PROFESSOR: So I have one more visual for you about--

let's get these out of the way.

OK, so let's come back to this picture.

So that's a picture of the gradient of the function

x squared plus y squared.

And we saw before that the gradient is always

Video


Download video file

Transcripts


Download SubRip (.srt) file Download Text (.txt) file

⊞ Calculator

Looking back at the level curves of $f(x,y)=x^2+y^2$ along with the gradient vectors, you can see that the gradient vectors are longer where the slope of the function is bigger. This means that f(x,y) is steeper at those places. The 3D plot of the surface is shown below for comparison.



► Paraboloid 💃



9. Putting it together

Topic: Unit 2: Geometry of Derivatives / 9. Putting it together

Hide Discussion

Add a Post

Show all posts by recent activity

[staff] number of attempts

Hello, For last question "Estimate the size of the gradient" there's 3 choices and 3 attempts. For previous question "Guess the directi....

2

Previous

Next >

© All Rights Reserved



edX

About

Affiliates

edX for Business

Open edX

Careers

News

Legal

Terms of Service & Honor Code

Privacy Policy

Accessibility Policy

Trademark Policy

<u>Sitemap</u>

Connect

Blog

Contact Us

Help Center

Media Kit

Donate













© 2021 edX Inc. All rights reserved.

深圳市恒宇博科技有限公司 <u>粤ICP备17044299号-2</u>