

Expectation time - General birth and death process

Asked 4 years, 1 month ago Active 4 years, 1 month ago Viewed 2k times



3



I study continuous time Markov chain and more specifically birth and death processes. I am trying to understand how to calculate the expectation time it takes to start from a state i to a state i+1. I am using the book "Introduction to probability models" by Sheldon Ross but I have trouble understanding it and I did not find alternative notes on my problem so I am asking here.



4)

We consider a general birth and death process with birth rate $\{\lambda_n\}$ and death rates $\{\mu_n\}$, where $\mu_0=0$ and we denote T_i as the time it takes starting from state i to enter state i+1. Since the times of death and births are exponential, we already know that $E[T_0]=\frac{1}{\lambda_0}$.

Then, the book follows this logic: For i > 0, we condition whether the first transition takes the process into state i - 1 or i + 1, i.e we let:

$$I_i = \left\{ egin{aligned} 0, & ext{if the first transition from } i & ext{is } i+1 \ 1, & ext{if the first transition from } i & ext{is } i-1 \end{aligned}
ight.$$

And therefore we note that:

$$E[T_i \mid I_i = 1] = \frac{1}{\lambda_i + \mu_i},$$

$$E[T_i \mid I_i = 0] = rac{1}{\lambda_i + \mu_i} + E[T_{i-1}] + E[T_i]$$

(Because independently of whether the first transition is a birth or a death, the time until it occurs is the minimum and thus $\sim \text{Exp}(\lambda_i + \mu_i)$.

And here is my problem:

To compute $E[T_i]$ I simply apply the following formula:

$$E[T_i] = E[T_i \mid I_i = 1] \cdot P(I_i = 1) + E[T_i \mid I_i = 0] \cdot P(I_i = 0)$$

and after working it out, I find:

$$E[T_i] = rac{1}{\lambda_i + \mu_i} + rac{\mu_i}{\lambda_i(\lambda_i + \mu_i)} + rac{\mu_i}{\lambda_i} E[T_{i-1}]$$

But the book gives:

"Since the probability that the first transition is a birth is $\frac{\lambda_i}{(\lambda_i + \mu_i)}$ we have :"

$$E[T_i] = rac{1}{\lambda_i + \mu_i} + rac{\mu_i}{(\lambda_i + \mu_i)} (E[T_{i-1}] + E[T_i])$$

$$\implies E[T_i] = \frac{1}{\lambda_i} + \frac{\mu_i}{\lambda_i} E[T_{i-1}]$$

Obviously the use of my formula is not appropriate here and I added useless terms, but I don't see which ones and the reasoning behind it. Understanding these kind of steps would help me a lot to better grasp this topic, so any help or idea is really appreciated, thanks!



Share Cite Edit Follow Flag

edited Jul 16 '17 at 12:54

asked Jul 16 '17 at 10:13 user386721

2 It happens that

$$rac{1}{\lambda_i + \mu_i} + rac{\mu_i}{\lambda_i (\lambda_i + \mu_i)} = rac{1}{\lambda_i}$$

hence the two recursions are equivalent. - Did Jul 16 '17 at 10:48



I feel a bit foolish I did not see this mini step.. Thanks for your help! – user386721 Jul 16 '17 at 12:53