



< Previous

✓

✓

✓


Next >

# 14.2.4 Example: Martian lander with drag and entry angle variability


 Bookmark this page

Now we consider the Martian lander problem with two parameters varying, specifically  $C_{DI}$  and  $\theta_e$ , using the uniform distributions over the ranges previously described. The implementation of the Monte Carlo method for this problem is shown in the code below. In this implementation the entire array of random values for  $C_{DI}$  and for  $\theta_e$  are done by calling `rng.uniform` with the size set to **Nsample**. The results for a typical simulation with sample size 100 are shown in Figures 14.9 and 14.10. For this simulation, 37 of the 100 instances had  $z_p < 9$  km, thus, the probability estimate would be  $P_{\text{low}} = 0.37$ .

```
rng = np.random.default_rng()
Nsample = 100
zps = np.zeros(Nsample)
CDIs = rng.uniform(1.5, 1.9, Nsample)
thetaes = rng.uniform(80., 86., Nsample)
for n in range(Nsample):
    lander_IVP.set_p('CD_l', CDIs[n])
    lander_IVP.set_p('theta_e', thetaes[n])
    zps[n] = lander_run_case(lander_IVP, dt,
IVP.step_RK4)
```




**Figure 14.9:** Scatter plot of  $(C_{DI}, \theta_e)$  values in a Monte Carlo simulation with a sample size of 100 drawn for uniform distributions of  $C_{DI}$  between 1.5 and 1.9 and  $\theta_e$  between  $80^\circ$  and  $86^\circ$ . Overlaid with contours of  $z_p$ .



Discussions

All posts sorted by recent activity




**Figure 14.10:** Scatter plot and histogram showing variability in  $z_p$  from a Monte Carlo simulation of sample size 100 with uniform distributions of  $C_{DI}$  between 1.5 and 1.9 and  $\theta_e$  between  $80^\circ$  and  $86^\circ$ . Similar to Figure 14.8, Figure 14.11 shows the values of  $C_{DI}$  in the Monte Carlo sample and the corresponding values of  $z_p$ . Because both  $C_{DI}$  and  $\theta_e$  are varying in this Monte Carlo simulation, the dependence of  $z_p$  on  $C_{DI}$  is not evident since variations in  $\theta_e$  create variations in  $z_p$  as well.

< Previous


Next >

© All Rights Reserved



**Figure 14.11:** Plot of  $z_p$  versus  $C_{DI}$  values from the Monte Carlo simulation results shown in Figure 14.10. The Python scripts used in the videos below (and several others) are available [here](#).

Video demonstrating Monte Carlo for Martian lander with CD and theta variation



[Start of transcript.](#)  
[Skip to the end.](#)

PROFESSOR: OK, now we're going to look at varying two parameters-- in this case, the drag coefficient, and theta, the entry

Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)



# edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

## Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)
- [Cookie Policy](#)
- [Your Privacy Choices](#)

## Connect

- [Idea Hub](#)
- [Contact Us](#)
- [Help Center](#)
- [Security](#)
- [Media Kit](#)



© 2023 edX LLC. All rights reserved.  
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)