

[Course](#)

[Progress](#)

[Dates](#)

[Discussion](#)

[Syllabus](#)

[Outline](#)

[laff routines](#)

[Community](#)

 [Course](#) / [Week 12: Eigenvalues and Eigenvectors](#) / [12.2 Getting Started](#)



< Previous	 ✓	 ✓	 ✓	 ✓		Next >
------------	---	---	---	---	---	--------

12.2.2 Simple Examples (continued)

 Bookmark this page

Week 12 due Dec 29, 2023 10:42 IST Completed

12.2.2 Simple Examples (continued)

Reading Assignment

0 points possible (ungraded)
Read Unit 12.2.2 of the notes. [\[LINK to Week12.pdf\]](#)

☒ Done/Skip

✓

Submit

Discussion

Topic: Week 12 / 12.2.2 Part 2

Hide Discussion

Add a Post

Show all posts ▼by recent activity ▼

There are no posts in this topic yet.

✕

Homework 12.2.2.4

1/1 point (graded)

Consider the diagonal matrix $\begin{pmatrix} \delta_0 & 0 & \cdots & 0 \\ 0 & \delta_1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \delta_{n-1} \end{pmatrix}$.

Eigenpairs for this matrix are given by $(\delta_0, e_0), (\delta_1, e_1), \dots, (\delta_{n-1}, e_{n-1})$, where e_j equals the j th unit basis vector.

Always ▼


✓ Answer: Always

We will simply say "by examination" this time and get back to a more elegant solution in Unit 12.3.3 .

Submit

ⓘ Answers are displayed within the problem

Video 12.2.2 Part 5



as is any multiple of that unit basis vector.

OK, so now let's make things a little bit more difficult.

Why don't you contemplate what the eigenvalues of an upper triangu...

Calculator



matrix are, and then find representative corresponding eigenvectors.
You will see me in the next video.

▶ 0:00 / 0:00

▶ 2.0x

🔊

🔍

CC

“

[End of transcript. Skip to the start.](#)

Video

📄 [Download video file](#)

Transcripts

📄 [Download SubRip \(.srt\) file](#)

📄 [Download Text \(.txt\) file](#)

Homework 12.2.2.5

10.0/10.0 points (graded)
Which of the following are eigenpairs (λ, x) of the 2×2 triangular matrix:

$$\begin{pmatrix} 3 & 1 \\ 0 & -1 \end{pmatrix} x = \lambda x,$$

where $x \neq 0$.

(Mark all correct answers.)

- ☒ $(-1, \begin{pmatrix} -1 \\ 4 \end{pmatrix}).$
- ☐ $(1/3, \begin{pmatrix} 1 \\ 0 \end{pmatrix}).$
- ☒ $(3, \begin{pmatrix} 1 \\ 0 \end{pmatrix}).$
- ☐ $(-1, \begin{pmatrix} 1 \\ 0 \end{pmatrix}).$
- ☒ $(3, \begin{pmatrix} -1 \\ 0 \end{pmatrix}).$
- ☐ $(-1, \begin{pmatrix} 3 \\ -1 \end{pmatrix}).$



Answers a), c), and e) are all correct, since for all $Ax = \lambda x$.

Answers b), d), and f) are not correct because they don't satisfy $Ax = \lambda x$.

Submit

Calculator

 Answers are displayed within the problem

Video 12.2.2 Part 6



 0:00 / 0:00

 2.0x












then you end up with the equation $4, 1; 0, 0$, times χ_0, χ_1 , equals $0, 0$.
Now this is in row echelon form.
Therefore, you pick the free variable to be equal to 1,
and you look for vectors of the form 1.
Something like that.
And if you then look at this right here, it's not hard to see that minus $1/4$ works.
And what you get are the eigenpairs.
Let's see, 3 comma 1, 0, and minus 1.
And the scalar multiple of that is also an eigenvector.
And therefore, why don't we instead pick minus 1, 4 for our eigenpair.
That's just a nicer vector, because it only involves integers.

Video
 [Download video file](#)

Transcripts
 [Download SubRip \(.srt\) file](#)
 [Download Text \(.txt\) file](#)

Homework 12.2.2.6

10.0/10.0 points (graded)

Consider the upper triangular matrix $U = \begin{pmatrix} v_{0,0} & v_{0,1} & \cdots & v_{0,n-1} \\ 0 & v_{1,1} & \cdots & v_{1,n-1} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & v_{n-1,n-1} \end{pmatrix}$.

The eigenvalues of this matrix are $v_{0,0}, v_{1,1}, \dots, v_{n-1,n-1}$.

Always 

 Answer: Always

The upper triangular matrix $U - v_{i,i}I$ has a zero on the diagonal. Hence it is singular. Because it is singular, $v_{i,i}$ is an eigenvalue. This is true for $i = 0, \dots, n - 1$.


Submit

 Answers are displayed within the problem

In the video, a few seconds in, the slide has an error (λ is subtracted from all elements above the diagonal).

Corrected slide:



 Calculator

$$\begin{pmatrix} 0 & v_{1,1} & \cdots & v_{1,n-1} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & v_{n-1,n-1} \end{pmatrix} x = \lambda x$$
$$\left[\begin{pmatrix} v_{0,0} & v_{0,1} & \cdots & v_{0,n-1} \\ 0 & v_{1,1} & \cdots & v_{1,n-1} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & v_{n-1,n-1} \end{pmatrix} - \lambda I \right] x = 0$$
$$\begin{pmatrix} v_{0,0} - \lambda & v_{0,1} & \cdots & v_{0,n-1} \\ 0 & v_{1,1} - \lambda & \cdots & v_{1,n-1} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & v_{n-1,n-1} - \lambda \end{pmatrix} x = 0$$

19 / 59

Video 12.2.2 Part 7

[Start of transcript. Skip to the end.](#)



▶ 0:00 / 0:00

▶ 2.0x

🔊

🔍

CC

“

Dr. Robert van de Geijn: What you have to do is subtract lambda from the diagonal, and you get this right here.

And then you ask yourself the question, under what circumstances is an upper triangular matrix singular?

Notice that if an upper triangular matrix has a 0 on the diagonal,

Video

📄 [Download video file](#)

Transcripts

- 📄 [Download SubRip \(.srt\) file](#)
- 📄 [Download Text \(.txt\) file](#)

Homework 12.2.2.7

4/4 points (graded)

Consider $A = \begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix}$

- The eigenvalue *largest in magnitude* is

✔ Answer: 4

- Which of the following are eigenvectors associated with this largest eigenvalue (in magnitude):

☐ $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$

☒ $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$

Calculator

☐ $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$

☒ $\begin{pmatrix} 2 \\ 2 \end{pmatrix}$

☐ $\begin{pmatrix} -1 \\ 2 \end{pmatrix}$



- The eigenvalue *smallest in magnitude* is

✔ Answer: -2

- Which of the following are eigenvectors associated with this smallest eigenvalue (in magnitude):

☒ $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$

☐ $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$

☐ $\begin{pmatrix} 2 \\ 2 \end{pmatrix}$

☐ $\begin{pmatrix} -1 \\ 2 \end{pmatrix}$



For the most part, you just take the solutions and plug them in to verify.

Submit

ⓘ Answers are displayed within the problem

Homework 12.2.2.8

10.0/10.0 points (graded)

Consider $A = \begin{pmatrix} -3 & -4 \\ 5 & 6 \end{pmatrix}$

- The eigenvalue *largest in magnitude* is

✔ Answer: 2

- The eigenvalue *smallest in magnitude* is

✔ Answer: 1

For the most part, you just take the solutions and plug them in to verify.

Submit

ⓘ Answers are displayed within the problem

Calculator

Video 12.2.2 Part 8

Start of transcript. Skip to the end.



Dr. Robert van de Geijn: Let's do another example.


Again it's a 2 by 2 matrix.

We should be able to go through the same procedure.

We subtract off lambda from the diagonal.

We recall that we need to look for the determinant of the matrix.

Video

 [Download video file](#)

Transcripts

 [Download SubRip \(.srt\) file](#)

 [Download Text \(.txt\) file](#)

Homework 12.2.2.9

1/1 point (graded)

Consider $\mathbf{A} = \begin{pmatrix} 2 & 2 \\ -1 & 4 \end{pmatrix}$. Which of the following are the eigenvalues of \mathbf{A} :

☐ 4 and 2.

☐ $3 + i$ and 2 .

● $3 + i$ and $3 - i$.

○ $2 + i$ and $2 - i$.



[← Previous](#)

Next >

© All Rights Reserved



edX

About

Affiliates

 Calculator

[edX for Business](#)
[Open edX](#)
[Careers](#)
[News](#)

Legal

[Terms of Service & Honor Code](#)
[Privacy Policy](#)
[Accessibility Policy](#)
[Trademark Policy](#)
[Sitemap](#)
[Cookie Policy](#)
[Your Privacy Choices](#)

Connect

[Idea Hub](#)
[Contact Us](#)
[Help Center](#)
[Security](#)
[Media Kit](#)



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)