Data Analysis: Statistical Modeling and Computation in Applications

<u>Help</u>

sandipan_dey >

<u>Course</u> <u>Progress</u>

Previous

gress Dates

Discussion

Resources






Next >



□ Bookmark this page

Exercises due Dec 1, 2021 17:29 IST Completed

The Effects of Measurement Noise



Start of transcript. Skip to the end.

Prof Jegelka: So finally, I would like to go through two other choices of modeling

that you can consider when you look at your data.

And these are measurement noise and non-stationarity

in your data.

So let's first remind ourselves about stationarity.

Video

Download video file

Transcripts

<u>Download SubRip (.srt) file</u> <u>Download Text (.txt) file</u>

Thus far, we have assumed that the kernel is the only contribution to the covariance matrix of the data. Although is was not stated explicitly, this means we took the data, or the observations, as exact.

However, what happens if our sensors are faulty or imprecise? How can one introduce in the presented framework the uncertainty in the observed values itself?

Let us return to the original example of temperature measurements. Imagine the sensor providing the measurements is not perfect, and each time it takes a measurement, it induces some additional noise due to its inherit limitations, such as thermal noise. One can characterize such noise, for example, as an additional variable ε that is also Normally distributed with mean zero and standard deviation τ . We can denote this as $\varepsilon \sim \mathcal{N}\left(0, \tau^2\right)$.

Now, using the same notation as before, instead of directly observing the realizations of the random variable \mathbf{X}_2 , we observe the realizations \mathbf{y}_2 of the random variable $\mathbf{Y}_2 = \mathbf{X}_2 + \varepsilon$. By definition, we will assume the random variable ε is independent of \mathbf{X}_2 .

Recall that the parameters of the conditional distribution of \mathbf{X}_1 given \mathbf{X}_2 are

$$\mu_{\mathbf{X}_1|\mathbf{X}_2} = \mu_1 + \Sigma_{12}\Sigma_{22}^{-1}\left(\mathbf{x}_2 - \mu_2
ight)$$

$$\Sigma_{\mathbf{X}_1|\mathbf{X}_2} = \Sigma_{11} - \Sigma_{12}\Sigma_{22}^{-1}\Sigma_{21}.$$

Adding arepsilon to $\mathbf{X_2}$ alters the covariance matrix:

$$oldsymbol{\Sigma} = egin{bmatrix} oldsymbol{\Sigma}_{11} & oldsymbol{\Sigma}_{12} \ oldsymbol{\Sigma}_{21} & oldsymbol{\Sigma}_{22} + au^2 I \end{bmatrix}$$


such that diagonal elements of magnitude au^2 are added to the $oldsymbol{\Sigma_{22}}$ component.

This changes the conditional equations to be

$$\mu_{\mathbf{X}_1 | \mathbf{Y}_2} = \mu_1 + \Sigma_{12} (\Sigma_{22} + au^2 I)^{-1} (\mathbf{y}_2 - \mu_2)$$

$$\Sigma_{\mathbf{X}_1|\mathbf{Y}_2} = \Sigma_{11} - \Sigma_{12} (\Sigma_{22} + au^2 I)^{-1} \Sigma_{21}.$$

The below figure shows a Guassian process for two different values of au, au=0.8 and au=0.1.



35: The effects of noisy measurements

We can note that as au is increases, so too does the variance on the estimate increase.

Observational Noise 1

1/1 point (graded)

What happens in the extreme case where $\tau \to \infty$? What happens to the mean of the estimates compared to the prior assumed mean?







Solution:

As $au o\infty$ we get

$$\mu_{\mathbf{X}_1|X_2} \ = \mu_1 + \Sigma_{12} (\Sigma_{22} + au^2 I)^{-1} \left(x_2 - \mu_2
ight)
ightarrow \mu_1,$$

and so the mean remains the same as the prior assumed mean of μ_1 .

Submit

You have used 1 of 2 attempts


1 Answers are displayed within the problem

Observational Noise 2

1/1 point (graded)

Consider, again, the extreme case where $\tau \to \infty$. If we increase the number of observations in this limit, do the additional observations reduce the variance on the estimate?

Variance decreases





Solution:

As $au o\infty$ we get

$$\Sigma_{\mathbf{X}_1|X_2} \ = \Sigma_{11} - \Sigma_{12} (\Sigma_{22} + au^2 I)^{-1} \Sigma_{21}
ightarrow \Sigma_{11},$$

and so the variance does not decrease with additional observations.

Submit

You have used 1 of 1 attempt

Answers are displayed within the problem

Discussion Hide Discussion

Topic: Module 5: Environmental Data and Gaussian Processes:Spatial Prediction / 6.

The Effects of Measurement Noise

Add a Post

Show all posts	∨	by recent activity 🗸
There are no posts in this topic yet.		
×		

© All Rights Reserved



edX

<u>About</u>

<u>Affiliates</u>

edX for Business

Open edX

Careers

<u>News</u>

Legal

Terms of Service & Honor Code

Privacy Policy

Accessibility Policy

<u>Trademark Policy</u>

<u>Sitemap</u>

Connect

<u>Blog</u>

Contact Us

Help Center

Media Kit

Donate















© 2021 edX Inc. All rights reserved.

深圳市恒宇博科技有限公司 <u>粤ICP备17044299号-2</u>