

MITx: 14.310x Data Analysis for Social Scientists

Heli

Bookmarks

- Module 1: The Basics of R and Introduction to the Course
- Entrance Survey
- Module 2:

 Fundamentals of
 Probability, Random

 Variables, Distributions, and Joint Distributions
- Module 3: Gathering and Collecting Data, Ethics, and Kernel Density Estimates
- Module 4: Joint,
 Marginal, and
 Conditional
 Distributions &
 Functions of Random
 Variable

Module 9: Single and Multivariate Linear Models > The Linear Model > Assumptions of the Linear Model - Quiz

Assumptions of the Linear Model - Quiz


 \square Bookmark this page

Recall our linear model:

$$Y_i = eta_0 + eta_1 x_i + \epsilon ext{ for } i = 1, 2, \dots, n$$

Question 1

1/1 point (graded)



- Module 5: Moments of a Random Variable,
 Applications to Auctions,
 Intro to Regression
- Module 6: Special
 Distributions, the
 Sample Mean, the
 Central Limit Theorem,
 and Estimation
- Module 7: Assessing and Deriving Estimators -Confidence Intervals, and Hypothesis Testing
- Module 8: Causality,
 Analyzing Randomized
 Experiments, &
 Nonparametric
 Regression
- Module 9: Single and Multivariate Linear Models

The Linear Model

due Nov 28, 2016 05:00 IST

Ø,

- \circ a. X_i, ϵ_i uncorrelated
- b. Identification
- lacksquare c. $E[\epsilon_i]=0$
- ullet d. Homoscedasticity. $E[\epsilon_i^2] = \sigma^2$ for all i.
- ullet e. No serial correlation. $E[\epsilon_i\epsilon_j]=0$ for i
 eq j.

Explanation

In the linear model, we assume that there is some variation in our regressor X. If X were always the exact same value regardless of the value of Y, then we would not be able to predict or learn anything about Y based on X.

Submit

You have used 1 of 2 attempts

Correct (1/1 point)

Question 2

<u>The Multivariate Linear</u> Model


due Nov 28, 2016 05:00 IST

Ø.

Module 9: Homework due Nov 21, 2016 05:00 IST

- Module 10: Practical
 Issues in Running
 Regressions, and
 Omitted Variable Bias
- Exit Survey

1/1 point (graded)



- lacksquare a. X_i, ϵ_i uncorrelated
- b. Identification
- \circ c. $E[\epsilon_i]=0$
- igcirc d. Homoscedasticity. $E[\epsilon_i^2] = \sigma^2$ for all i.
- ullet e. No serial correlation. $E[\epsilon_i\epsilon_j]=0$ for i
 eq j. ullet

Explanation

In the linear model, we assume that there are no areas where errors are mostly positive or other areas where errors are mostly negative.


Submit

You have used 1 of 2 attempts

Correct (1/1 point)

Question 3

1/1 point (graded)



- \circ a. X_i, ϵ_i uncorrelated
- b. Identification
- ullet c. $E[\epsilon_i]=0$ 🗸
- ullet d. Homoscedasticity. $E[\epsilon_i^2] = \sigma^2$ for all i.
- ullet e. No serial correlation. $E[\epsilon_i\epsilon_j]=0$ for i
 eq j.

Explanation

In the linear model, we assume that the expectation of the error is zero. There is no way for us to between a non-zero error mean and a different intercept β_0 , so we just assume the error is zero in expectation.


Submit

You have used 1 of 2 attempts

✓ Correct (1/1 point)

Question 4

1/1 point (graded)



- \circ a. X_i, ϵ_i uncorrelated
- b. Identification
- \circ c. $E[\epsilon_i]=0$
- ullet d. Homoscedasticity. $E[\epsilon_i^2] = \sigma^2$ for all i.
- ullet e. No serial correlation. $E[\epsilon_i\epsilon_j]=0$ for i
 eq j.

Explanation

Explanation: We assume homoscedasticity in the linear model, meaning the error variance should be consistent across all values of x. In the case above, the variance of the error is much higher at lower values of x.

Submit

You have used 1 of 2 attempts

✓ Correct (1/1 point)

Discussion

Show Discussion

Topic: Module 9 / Assumptions of the Linear Model - Quiz

© All Rights Reserved



© 2016 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

















