



15.2.1 Probability Density Functions (PDFs) and Random Variables

 [Bookmark this page](#)

M02.1

M02.13

A variable whose value depends on the outcome of a random phenomenon is known as a *random variable*. In the projectile example, the input parameters V_0 and θ_0 are random variables because we have said they have randomness. And, as a result, the impact location x_f is also a random variable (since it depends on these random variables).

A typical notation used for a random variable is capital letters. For example, the impact location would be referred to as X_f and the meaning is that X_f does not have a precise value but rather may take on a range of values. We say that X_f has a distribution of values.

A common way to describe the distribution of a random variable is through its probability density function (or PDF for short). Let the PDF of a variable X be $f(x)$ then the probability that $x - dx/2 < X < x + dx/2$ is equal to $f(x) dx$, i.e.,

$$P(x - \frac{dx}{2} < X < x + \frac{dx}{2}) = f(x) \, dx$$

(15.2)

Thus, the PDF is a measure of how likely the value of a random variable is. From this definition of the PDF, the probability that $X \leq x$ is related to $f(x)$ through the following integral,



edX

[About](#)
[Affiliates](#)
[edX for Business](#)
[OpenX](#)
[When X](#)
[Capreba](#)
[Nemo](#)

$$P(X \leq x) = \int_{-\infty}^x f(\xi) \, d\xi$$

(15.3)

When we consider all possible values, i.e. $x \rightarrow +\infty$, then the probability $P(X \leq +\infty)$ must be one. This means that the integral of $f(x)$ over the entire real numbers must be one,

Legal

[Terms of Service & Honor Code](#)
[Privacy Policy](#)
[Accessibility Policy](#)
[Trademark Policy](#)
[Sitemap](#)
[Cookie Policy](#)
[Your Privacy Choices](#)

$$\int_{-\infty}^{+\infty} f(\xi) \, d\xi = 1$$

(15.4)

Often, we are interested in the probability that X lies between x_a and x_b . That can be found as follows,

$$P(x_a \leq X \leq x_b) = \int_{x_a}^{x_b} f(\xi) \, d\xi$$

(15.5)

Connect

The following video discusses PDFs and how to use the matplotlib hist command to plot them from an array of data.

[Idea Hub](#)
[Contact Us](#)

Discussions

All posts sorted by recent activity

© All Rights Reserved

[Help Center](#)

[Security](#)

[Media Kit](#)

Previous

Next



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)



And so X would be a random variable.