

edX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



Lecture 3: Parametric Statistical

Course > Unit 2 Foundation of Inference > Models

> 5. Statistical model

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. **Upgrade now**

5. Statistical model

Statistical model: definition

[Start of transcript. Skip to the end.](#)





So let's talk about a statistical model.

Here I said, you know, I want to replace my PDF by a particular statistical model, which was

in this case a Poisson model.

A model just means something which is like slightly simpler than what reality actually

is, but hopefully captures most of it.

Well, that would be a good model.

And so I'm going to have a statistical



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

A Basic Statistical Model: Sample space

1/1 point (graded)

You have a coin that either lands heads, which you denote by 1 , or tails, which you denote by 0 . Let X be a random variable representing this coin flip, with an (unknown) distribution. You run a **statistical experiment** consisting of n iid tosses of the coin and record your data set as $X_1, X_2, X_3, \dots, X_n$.

(It makes sense to assume the coin tosses X_1, \dots, X_n as identically distributed, since we always toss the same coin; and as independent, since these tosses do not affect each other.)

We now construct a **statistical model** $(E, \{P_\theta\}_{\theta \in \Theta})$ associated with this experiment, where

- E is a sample space for X , i.e. a set that contains all possible outcomes of X ,
- $\{\mathbb{P}_\theta\}_{\theta \in \Theta}$ is a family of probability distributions on E ,
- Θ is a parameter set, i.e. a set consisting of some possible values of θ .

What is the **smallest sample space** for X ? We can use this as the sample space E in our statistical model.

(Below, $[0, 1]$ denotes the closed interval between 0 and 1 . In contrast, $\{0, 1\}$ denotes the set with two elements, 0 and 1 .)

☒ $\{0, 1\}$ ✓

☐ $[0, 1]$

☐ \mathbb{R}

☐ \mathbb{R}^2

Solution:

Here the coin is either heads (denoted by 1) or tails (denoted by 0), so $\{0, 1\}$ is the smallest sample space of X . The remaining choices are valid, but not the smallest, sample spaces of X .

You have used 1 of 2 attempts

i Answers are displayed within the problem

A Basic Statistical Model: Family of distributions and Parameter set

2/2 points (graded)

Continuing from the previous problem, which of the following is the smallest family of probability distributions that the distribution of X belongs to? We can use this family as $\{\mathbb{P}_\theta\}_{\theta \in \Theta}$ in our statistical model.

☒ Bernoulli ✓☐ Poisson☐ Binomial

The distribution of X is a member of the family with some unknown parameter θ . According to the information given about the experiment, which of the following represents the set of all possible values of the parameter θ ? We can use this set as the parameter set Θ in our statistical model.

☐ $\{0, 1\}$ ☐ $\{0, 1/2, 1\}$ ☒ $[0, 1]$ ✓☐ \mathbb{R} **Solution:**

1. Since the (smallest) sample space of X is $\{0, 1\}$, X follows a Bernoulli distribution.
2. The first and second choices, $\{0, 1\}$ and $\{0, 1/2, 1\}$, place too many restrictions on the distribution of X . Also, be sure to not confuse the space where the parameter θ lives with the sample space, where the random variable X lives! The fourth choice, \mathbb{R} , allows for values of θ that do not make sense according to modeling X as $\text{Ber}(\theta)$. For example, there is no such thing as $\text{Ber}(-1/2)$.
We are not given any assumptions on the distribution of the coin, so we need to allow θ to take all possible values that make sense according to our modeling assumption. Since θ represents the probability that $X = 1$, we must have $0 \leq \theta \leq 1$. Hence, the third choice, $[0, 1]$, is correct.

Using this problem and the previous one, we can construct the statistical model $(\{0, 1\}, \{\text{Ber}(\theta)\}_{\theta \in [0, 1]})$ for the distribution of the RV X representing the outcome of the coin flip.

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Discussion

[Hide Discussion](#)

Topic: Unit 2 Foundation of Inference:Lecture 3: Parametric Statistical Models / 5. Statistical model

[Add a Post](#)

Show all posts ▼

by recent activity ▼

? Set of outcomes of random variable X. What is it, the Domain or the Image of the function X?

3

As the definition of the random variable is the function I would like to clarify the definition of Sample Space. The set of the outcomes, is it the Domai...

[Learn About Verified Certificates](#)

© All Rights Reserved