



[Course](#) > [Midterm Exam 1](#) > [Midterm Exam 1](#) > 3.

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

3.

Setup:

We continue with the problem on the previous page.

Recall the lifetime (in months) of a battery is modeled by a random variable X that has pdf

$$f_{\theta}(x) = K\theta^x \mathbf{1}(x > 0) \quad \text{where } K = \ln(1/\theta)$$

for an unknown parameter $\theta \in (0, 1)$.

Assume instead that we cannot actually observe the lifetime of the batteries. Instead, we only observe if the battery is still working after τ months for some known τ to be chosen later (this is called **censored data**).

Let Y_1, \dots, Y_n be our observations where $Y_i = \mathbf{1}(X_i > \tau)$ indicates that the i th battery is still working after τ months. Our goal is to estimate $\theta \in (0, 1)$ (the parameter for the pdf of X) based on this new data.

Distribution of Y

2/2 points (graded)

What kind of distribution does $Y = \mathbf{1}(X > \tau)$ follow?

☐ Standard Gaussian

☐ Gaussian with mean $\mu \neq 0$

☒ Bernoulli

☐ Cannot be determined



What is $\mathbb{E}[Y_i]$?

$\mathbb{E}[Y_i] =$

theta^tau

✓ Answer: theta^tau

θ^τ

STANDARD NOTATION

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Statistical Model

1/1 point (graded)

What is the statistical model $(E, \mathbf{P}_{\theta \in \Theta})$ associated to this experiment?

☐ $(\{0, 1\}, \text{Ber}(\theta^\tau)_{\tau \in (0,1)})$

☒ $(\{0, 1\}, \text{Ber}(\theta^\tau)_{\theta \in (0,1)})$

☐ $((-\infty, \infty), \mathcal{N}(0, \theta^\tau)_{\theta \in (0,1)})$

☐ $((-\infty, \infty), \mathcal{N}(\mu, \theta^\tau)_{\tau \in (0,1)})$



Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

MLE

2.0/2.0 points (graded)

Find the maximum likelihood estimator $\tilde{\theta}$ of θ **based on the censored data**.

(Enter **barY_n** for $\overline{Y_n}$ and **tau** for τ .)

$\tilde{\theta} =$

$(\text{barY_n})^{1/\text{tau}}$

✓ Answer: $\text{barY_n}^{1/\text{tau}}$

STANDARD NOTATION

Solution:

the maximum likelihood estimator $\tilde{\theta}$ of θ **based on the censored data** is given by $\hat{\theta} = (\overline{Y_n})^{\frac{1}{\tau}}$.

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Consistency

1/1 point (graded)

Is $\tilde{\theta}$ a consistent estimator? If yes, enter the limit in probability; that is, enter the constant C such that $\tilde{\theta} \xrightarrow[n \rightarrow \infty]{\mathbf{P}} C$ (where the convergence is in probability). If no, enter **-999**.

$C =$

theta

✓ Answer: theta

θ

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Asymptotic Variance

3.0/3.0 points (graded)

The quantity $\sqrt{n}(\tilde{\theta} - \theta)$ converges in distribution to $\mathcal{N}(0, \tilde{\sigma}^2)$. Find the asymptotic variance $\tilde{\sigma}^2$.

$\tilde{\sigma}^2 =$

(theta^(2-tau)-theta^2)/tau^2

✓ Answer: 1/tau^2*theta^(-tau+2)*(1-theta^tau)

$\frac{\theta^{2-\tau} - \theta^2}{\tau^2}$

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Different Tau's

1/1 point (graded)

Previous studies indicate that $\theta > 1/3$. Would you recommend using $\tau = 1$ or $\tau = 2$? Select the correct reasoning.

(Choose all that apply.)

☐ $\tau = 1$ is recommended.

☒ $\tau = 2$ is recommended.

☒ The recommended τ value should give a smaller asymptotic variance.

☐ The recommended τ value should give a larger asymptotic variance.



Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Error and Bug Reports/Technical Issues

Topic: Midterm Exam 1:Midterm Exam 1 / 3.

Hide Discussion

Add a Post

Show all posts ▼

by recent activity ▼

💬 Progress Bar Chart	2
Is it only me who can't see the progress bar chart? If not, what happened to it?	
💬 Resolution of the midterm exam using Julia	7
💬 [Staff] Problem with the answer of Part '5'	6
I think the following answer is not correct: $\tau^2 \theta^{-(\tau+2)} (1-\theta^\tau)$. It could be: $(1/\tau^2) \theta^{-(\tau+2)} (1-\theta^\tau)$. Though, I have got the above answer. I m...	
💬 [Staff] Replacing θ^t with a Variable to Find MLE	1
💬 [Staff] Midterm Solution	5
Dear TA, Can we get the solution of the test? It can really help us to learn deeper from our own mistakes. Thanks, Tony.	
💬 Grader on Asymptotic Variance part has been corrected	2
At least for me, the grader shows now the right answer.	
💬 Asymptotic	32
Will there be a detailed explanation somewhere? I did that part on paper multiple times and I arrived to a slightly different result and now I'm wondering what did I do wr...	
? [STAFF] Grader on Asymptotic Variance part	10
💬 Is the answer for the Asymptotic variance correct?	31
Per MLE question, θ is the function of \bar{Y}_n, or $(\bar{Y}_n)^{1/\tau}$. If we take the derivative and square it, it should be $(\bar{Y}_n)^{2/(\tau-2)} / \tau^2$. Then the asy. var. sho...	
💬 [Staff] There's something I noticed in this exam	9
But it will most likely break the rules if I say it here. Where can I write my concern? (an e-mail address or PM system)	
? Question about Grading	2
Hi, It seems that one of answers was not graded correctly (Part 3 Asymptotic Variance). It is the same result as the answer, but the order of the coefficients is different. Is it...	
💬 My Answer of Asymptotic Variance with Explanations	2
✓ Problem with displaying of mathematical expressions	3
I am experiencing a parsing problem where mathematical expressions are displayed as simple text. Usually when this happens (it happens a lot) I just refresh the page un...	
💬 [Staff] is it Deleted by MW-CTA	2
Deleted by MW-CTA	

Learn About Verified Certificates

© All Rights Reserved

