

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).




[Course](#) > [Unit 3 Methods of Estimation](#) > [Likelihood Estimation](#) > [Lecture 9: Introduction to Maximum Likelihood Estimation](#) > 9. Worked Example: Concavity and Composition of Functions

Currently enrolled in **Audit Track** (expires December 25, 2019) [Upgrade \(\\$300\)](#)

9. Worked Example: Concavity and Composition of Functions

Worked Example: Hessian and Concavity



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Combination of Convex functions

3/3 points (graded)

Let f_1, f_2 be convex functions on \mathbb{R} .

Determine if the following functions are necessarily convex or concave.

Hint: Recall a function $g : I \rightarrow \mathbb{R}$ is convex in the interval I is an interval, if for all pairs of real numbers $x_1 < x_2 \in I$

$$g(tx_1 + (1-t)x_2) \leq tg(x_1) + (1-t)g(x_2) \quad \text{for all } 0 \leq t \leq 1.$$

- $3f_1 + 2f_2$:

☒ Convex

☐ Concave

☐ Cannot be determined without more information



- $-10f_1$:

☐ Convex

☒ Concave

☐ Cannot be determined without more information



- f_2f_1 :

☐ Convex

☐ Concave

☒ Cannot be determined without more information

Solution:

Given f_1, f_2 are convex, we have

$$f_1(tx_1 + (1-t)x_2) \leq tf_1(x_1) + (1-t)f_1(x_2) \quad \text{for all } 0 \leq t \leq 1$$

and the same holds for f_2 .

- The same inequality holds for $g = 3f_1 + 2f_2$:

$$\begin{aligned} g(tx_1 + (1-t)x_2) &= 3f_1(tx_1 + (1-t)x_2) + 2f_2(tx_1 + (1-t)x_2) \\ &\leq 3(tf_1(x_1) + (1-t)f_1(x_2)) + 2(tf_2(x_1) + (1-t)f_2(x_2)) \\ &= tg(x_1) + (1-t)g(x_2). \end{aligned}$$

Hence $3f_1 + 2f_2$ is also convex.

Remark: In general, any function $c_1f_1 + c_2f_2$ where $c_1, c_2 > 0$ is convex if f_1, f_2 are.

- $-10f_1$ is concave, because it is negative of a convex function.
- f_1f_2 is not necessarily convex. For example, if $f_1(x) = x$, and $f_2 = x^2$, then $(f_1f_2)(x) = x^3$ which is neither convex nor concave. Other examples of f_1 and f_2 , e.g. $f_1 = f_2 = x^2$ will lead to f_1f_2 being convex.

You have used 2 of 2 attempts

i Answers are displayed within the problem

Discussion

[Hide Discussion](#)

Topic: Unit 3 Methods of Estimation:Lecture 9: Introduction to Maximum Likelihood Estimation / 9. Worked Example: Concavity and Composition of Functions

[Add a Post](#)

Show all posts ▼

by recent activity ▼

? [Both concave and convex \(non-strictly\)?](#)

1

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

[Learn About Verified Certificates](#)

© All Rights Reserved