EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the Privacy Policy.





Homework 1: Estimation,

Confidence Interval, Modes of

2. Biased and unbiased estimation

Course > Unit 2 Foundation of Inference > Convergence

> for variance of Bernoulli variables

## **Audit Access Expires Dec 24, 2019**

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. **Upgrade now** 

# 2. Biased and unbiased estimation for variance of Bernoulli variables

(a)

2/2 points (graded)

Let  $X_1,\ldots,X_n$  be i.i.d. Bernoulli random variables, with unknown parameter  $p\in(0,1)$ . The aim of this exercise is to estimate the common variance of the  $X_i$  .

First, recall what  $Var(X_i)$  is for Bernoulli random variables.

Let  $\overline{X}_n$  be the sample average of the  $X_i$  ,

$$\overline{X}_n = rac{1}{n} \sum_{i=1}^n X_i.$$

We are interested in finding an estimator for  $Var(X_i)$ , and propose to use

$$\hat{V} = \overline{X}_n (1 - \overline{X}_n)$$
 .

Check the correct statement that applies to  $\,\hat{V}$  :

- $\hat{V}$  is not consistent because  $\mathsf{Var}(X_i)$  is not linear in p
- ullet  $\hat{V}$  is consistent because of the Law of Large Numbers
- $\hat{V}$  is consistent because of the Central Limit Theorem



STANDARD NOTATION

Submit

You have used 1 of 2 attempts

- ✓ Correct (2/2 points)
- (b)

2/2 points (graded)

Now, we are interested in the bias of  $\hat{V}$  . Compute:

2. Biased and unbiased estimation for variance of Bernoulli variables | Homework 1: Estimation, Confidence Interval, Modes of Convergence | 18.6501x Courseware | edX

$$\mathbb{E}\left[\hat{V}
ight] - \mathsf{Var}\left(X_i
ight) = egin{bmatrix} (\mathsf{p}^2-\mathsf{p})/\mathsf{n} \ \hline rac{p^2-p}{n} \end{bmatrix}$$

Using this, find an unbiased estimator  $\,\hat{V}^{'}\,$  for  $\,p\,(1-p)\,$  if  $\,n\geq 2$  .

Write barx\_n for  $\overline{X}_n$  .

$$\hat{V}' = \boxed{ \text{n/(n-1)*(barX_n*(1-barX_)}} \checkmark$$

### STANDARD NOTATION

Submit

You have used 3 of 3 attempts

✓ Correct (2/2 points)

## Discussion

**Hide Discussion** 

Topic: Unit 2 Foundation of Inference: Homework 1: Estimation, Confidence Interval, Modes of Convergence / 2. Biased and unbiased estimation for variance of Bernoulli variables

#### Add a Post

Show all posts by recent activity ▼ 3 Lalready checked other question posted here but I'm still unable to eliminate P from my exercise although the exercise seems to be straight forward I first calculated E[V], rep... Any hints for b) please? Particularly how to find E[V]? Hint for part (b) 3

2. Biased and unbiased estimation for variance of Bernoulli variables | Homework 1: Estimation, Confidence Interval, Modes of Convergence | 18.6501x Courseware | edX

STAFF - Add "Show Answer" button after correct submission or no more attempts left

Ljust could not figure out the answer to part b. I got the unbiased estimator right after a long time of researching the internet, but unfortunately I could not figure out the sol...

Learn About Verified Certificates

© All Rights Reserved