

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



Course > [Unit 3 Methods of Estimation](#) > [Likelihood Estimation](#) > [Lecture 9: Introduction to Maximum Likelihood Estimation](#) > 10. Concavity in higher dimensions and Eigenvalues

Currently enrolled in **Audit Track** (expires December 25, 2019) [Upgrade \(\\$300\)](#)

10. Concavity in higher dimensions and Eigenvalues

Concavity in 2 dimensions: Compute the Hessian

4/4 points (graded)

What is the Hessian $\mathbf{H}f$ of the function $f(x, y) = -2x^2 + \sqrt{2}xy - \frac{5}{2}y^2$? Fill in the values of the entries of $\mathbf{H}f$.

$(\mathbf{H}f)_{11} =$ ✓ Answer: -4 $(\mathbf{H}f)_{12} =$ ✓ Answer: sqrt(2)

$(\mathbf{H}f)_{21} =$ ✓ Answer: sqrt(2) $(\mathbf{H}f)_{22} =$ ✓ Answer: -5

Solution:

We compute that

$$\begin{aligned} (\mathbf{H}f)_{11} &= \frac{\partial^2 f}{\partial x^2} = -4, & (\mathbf{H}f)_{12} &= \frac{\partial^2 f}{\partial x \partial y} = \sqrt{2} \\ (\mathbf{H}f)_{21} &= \frac{\partial^2 f}{\partial y \partial x} = \sqrt{2}, & (\mathbf{H}f)_{22} &= \frac{\partial^2 f}{\partial y^2} = -5. \end{aligned}$$

So this implies that

$$\mathbf{H}f = \begin{pmatrix} -4 & \sqrt{2} \\ \sqrt{2} & -5 \end{pmatrix}.$$

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

(Optional) Concavity in 2 dimensions: Positive Definiteness and Eigenvalues

0 points possible (ungraded)

A symmetric (real-valued) $d \times d$ matrix \mathbf{A} is **positive semi-definite** (*resp.* **positive definite**) if and only if all of its eigenvalues are **non-negative** (*resp.* **positive**).

Analogously, it is **negative semi-definite** (*resp.* **negative definite**) if and only if all of its eigenvalues are **non-positive** (*resp.* **negative**).

As above, consider $f(x, y) = -2x^2 + \sqrt{2}xy - \frac{5}{2}y^2$.

What are the eigenvalues λ_1, λ_2 of $\mathbf{H}f$? Assume that $\lambda_1 < \lambda_2$.

$\lambda_1 =$ **✓ Answer: -6** $\lambda_2 =$ **✓ Answer: -3**

Based on your answer to the last question, f is ...

☐ Convex

☒ Concave

☐ None of the Above**Solution:**

Recall from the previous problem that the Hessian of f is

$$\mathbf{H}f = \begin{pmatrix} -4 & \sqrt{2} \\ \sqrt{2} & -5 \end{pmatrix}.$$

To find the eigenvalues, we need to solve for λ such that

$$\det(\mathbf{H}f - \lambda I) = \det \left(\begin{pmatrix} -4 - \lambda & \sqrt{2} \\ \sqrt{2} & -5 - \lambda \end{pmatrix} \right) = \lambda^2 + 9\lambda + 18 = 0.$$

Factoring the quadratic: $\lambda^2 + 9\lambda + 18 = (\lambda + 6)(\lambda + 3)$ shows that $\lambda_1 = -6$ and $\lambda_2 = -3$.

The function f is twice-differentiable, so it is concave if $x^T \mathbf{H}f x \leq 0$ for all $x \in \mathbb{R}^2$. By the remark in the problem statement, this is equivalent to all of the eigenvalues of $\mathbf{H}f$ being negative. Hence, f is concave (in fact it is *strictly* concave).

You have used 1 of 2 attempts

 Answers are displayed within the problem

Discussion

Topic: Unit 3 Methods of Estimation:Lecture 9: Introduction to Maximum Likelihood Estimation / 10. Concavity in higher dimensions and Eigenvalues

Add a Post

Show all posts ▼

by recent activity ▼

💬 [Typo in solution to the first problem](#)

1

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

[Learn About Verified Certificates](#)

© All Rights Reserved