

Search MathWorld

0,

Algebra

Applied Mathematics

Calculus and Analysis

Discrete Mathematics

Foundations of Mathematics

Geometry

History and Terminology

Number Theory

Probability and Statistics

Recreational Mathematics

Topology

Alphabetical Index

Interactive Entries

Random Entry

New in MathWorld

MathWorld Classroom

About MathWorld

Contribute to MathWorld

Send a Message to the Team

MathWorld Book

Wolfram Web Resources »

13.690 entries Last updated: Tue Sep 17 2019

Created, developed, and nurtured by Eric Weisstein at Wolfram Research

Probability and Statistics > Statistical Distributions > Limit Theorems > Interactive Entries > Interactive Demonstrations >

Central Limit Theorem



Let $X_1, X_2, ..., X_N$ be a set of N independent random variates and each X_i have an arbitrary probability distribution $P(x_1, ..., x_N)$ with mean μ_i and a finite variance σ_i^2 . Then the normal form variate

$$X_{\text{norm}} \equiv \frac{\sum_{i=1}^{N} x_i - \sum_{i=1}^{N} \mu_i}{\sqrt{\sum_{i=1}^{N} \sigma_i^2}}$$
(1)

has a limiting cumulative distribution function which approaches a normal distribution.

 $=\sum_{n=0}^{\infty}\frac{(2\pi i f)^n}{n!}\langle X^n\rangle.$

Under additional conditions on the distribution of the addend, the probability density itself is also normal (Feller 1971) with mean $\mu = 0$ and variance $\sigma^2 = 1$. If conversion to normal form is not performed, then the variate

$$X = \frac{1}{N} \sum_{i=1}^{N} x_i \tag{2}$$

is normally distributed with $\mu_X = \mu_X$ and $\sigma_X = \sigma_X / \sqrt{N}$.

Kallenberg (1997) gives a six-line proof of the central limit theorem. For an elementary, but slightly more cumbersome proof of the central limit theorem, consider the inverse Fourier transform of $P_X(f)$.

$$\mathcal{F}_{f}^{-1}\left[P_{X}\left(f\right)\right]\left(x\right) \equiv \int_{-\infty}^{\infty} e^{2\pi i f X} P\left(X\right) dX \tag{3}$$

$$= \int_{-\infty}^{\infty} \sum_{n=0}^{\infty} \frac{(2\pi i f X)^{n}}{n!} P\left(X\right) dX \tag{4}$$

$$= \sum_{n=0}^{\infty} \frac{(2\pi i f)^{n}}{n!} \int_{-\infty}^{\infty} X^{n} P\left(X\right) dX \tag{5}$$

Now write

$$\langle X^{n} \rangle = \langle N^{-n} (x_{1} + x_{2} + \dots + x_{N})^{n} \rangle$$

$$= \int_{-\infty}^{\infty} N^{-n} (x_{1} + \dots + x_{N})^{n} P(x_{1}) \cdots P(x_{N}) dx_{1} \cdots dx_{N},$$
(7)

so we have

$$\mathcal{F}_{f}^{-1}[P_{X}(f)](x) =$$
 (8)

THINGS TO TRY: = binomial distribution = normal distribution = ack(2,ack(2,1))





(6)

Chris Boucher

$$\sum_{n=0}^{\infty} \frac{(2\pi i f)^n}{n!} \langle X^n \rangle$$

$$= \sum_{n=0}^{\infty} \frac{(2\pi i f)^n}{n!} \int_{-\infty}^{\infty} N^{-n} (x_1 + \dots + x_N)^n \times P(x_1) \cdots P(x_N) dx_1 \cdots dx_N$$

$$= \int_{-\infty}^{\infty} \sum_{n=0}^{\infty} \left[\frac{2\pi i f (x_1 + \dots + x_N)}{N} \right]^n \frac{1}{n!} P(x_1) \cdots P(x_N) dx_1 \cdots dx_N$$

$$= \int_{-\infty}^{\infty} e^{2\pi i f (x_1 + \dots + x_N)/N} P(x_1) \cdots P(x_N) dx_1 \cdots dx_N$$

$$= \left[\int_{-\infty}^{\infty} e^{2\pi i f x_1/N} P(x_1) dx_1 \right] \times \cdots \times \left[\int_{-\infty}^{\infty} e^{2\pi i f x_N/N} P(x_N) dx_N \right]$$

$$= \left[\int_{-\infty}^{\infty} e^{2\pi i f x/N} P(x) dx \right]^N$$

$$= \left[1 + \left(\frac{2\pi i f}{N} \right) x + \frac{1}{2} \left(\frac{2\pi i f}{N} \right)^2 x^2 + \dots \right] P(x) dx \right]^N$$

$$= \left[1 + \frac{2\pi i f}{N} \langle x \rangle - \frac{(2\pi f)^2}{2N^2} \langle x^2 \rangle + O(N^{-3}) \right]^N$$

$$= \exp \left\{ N \ln \left[1 + \frac{2\pi i f}{N} \langle x \rangle - \frac{(2\pi f)^2}{2N^2} \langle x^2 \rangle + O(N^{-3}) \right] \right\}.$$
(16)

Now expand

$$\ln(1+x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 + \dots, \tag{17}$$

so

$$\mathcal{F}_{f}^{-1} [P_{X} (f)] (x) \approx \exp \left\{ N \left[\frac{2 \pi i f}{N} \langle x \rangle - \frac{(2 \pi f)^{2}}{2 N^{2}} \langle x^{2} \rangle + \frac{1}{2} \frac{(2 \pi i f)^{2}}{N^{2}} \langle x \rangle^{2} + O(N^{-3}) \right] \right\}$$

$$= \exp \left[2 \pi i f \langle x \rangle - \frac{(2 \pi f)^{2} (\langle x^{2} \rangle - \langle x \rangle^{2})}{2 N} + O(N^{-2}) \right]$$

$$\approx \exp \left[2 \pi i f \mu_{x} - \frac{(2 \pi f)^{2} \sigma_{x}^{2}}{2 N} \right],$$

since

$$\mu_{x} \equiv \langle x \rangle$$

$$\sigma_{x}^{2} \equiv \langle x^{2} \rangle - \langle x \rangle^{2}.$$
(21)

Taking the Fourier transform,

$$P_X \equiv \int_{-\infty}^{\infty} e^{-2\pi i f x} \mathcal{F}^{-1} [P_X (f)] df$$

$$= \int_{-\infty}^{\infty} e^{2\pi i f (\mu_X - x) - (2\pi f)^2 \sigma_X^2 / 2N} df.$$
(23)

This is of the form

$$\int_{-\infty}^{\infty} e^{i a f - b f^2} df, \tag{25}$$

where $a \equiv 2\pi (\mu_x - x)$ and $b \equiv (2\pi\sigma_x)^2/2$ N. But this is a Fourier transform of a Gaussian function, so



Arithmetic

Integration

Limits

Derivatives

Equation Solving

Expression Expansion

Matrix Row Reduction

Partial Fractions

Polynomial Factoring

Extrema

Student pricing

(18)

(19)

(20)

$$\int_{-\infty}^{\infty} e^{i \, a \, f - b \, f^2} \, d \, f = e^{-a^2/4 \, b} \, \sqrt{\frac{\pi}{b}} \tag{26}$$

(e.g., Abramowitz and Stegun 1972, p. 302, equation 7.4.6). Therefore,

$$P_X = \sqrt{\frac{\pi}{\frac{(2\pi\sigma_x)^2}{2N}}} \exp\left\{\frac{-[2\pi(\mu_x - x)]^2}{4\frac{(2\pi\sigma_x)^2}{2N}}\right\}$$
 (27)

= (28)

(29)

But and , so

(30)

The "fuzzy" central limit theorem says that data which are influenced by many small and unrelated random effects are approximately normally distributed.

SEE ALSO:

Berry-Esséen Theorem, Fourier Transform--Gaussian, Lindeberg Condition, Lindeberg-Feller Central Limit Theorem, Lyapunov Condition

REFERENCES:

Abramowitz, M. and Stegun, I. A. (Eds.). Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, 9th printing. New York: Dover, 1972.

Feller, W. "The Fundamental Limit Theorems in Probability." Bull. Amer. Math. Soc. 51, 800-832, 1945.

Feller, W. An Introduction to Probability Theory and Its Applications, Vol. 1, 3rd ed. New York: Wiley, p. 229, 1968.

Feller, W. An Introduction to Probability Theory and Its Applications, Vol. 2, 3rd ed. New York: Wiley, 1971.

Kallenberg, O. Foundations of Modern Probability. New York: Springer-Verlag, 1997.

Lindeberg, J. W. "Eine neue Herleitung des Exponentialgesetzes in der Wahrscheinlichkeitsrechnung." Math. Z. 15, 211-225, 1922.

Spiegel, M. R. Theory and Problems of Probability and Statistics. New York: McGraw-Hill, pp. 112-113, 1992.

Trotter, H. F. "An Elementary Proof of the Central Limit Theorem." Arch. Math. 10, 226-234, 1959.

Zabell, S. L. "Alan Turing and the Central Limit Theorem." Amer. Math. Monthly 102, 483-494, 1995.

Referenced on Wolfram|Alpha: Central Limit Theorem

CITE THIS AS:

Weisstein, Eric W. "Central Limit Theorem." From MathWorld--A Wolfram Web Resource. http://mathworld.wolfram.com/CentralLimitTheorem.html

Wolfram Web Resources

Mathematica »

The #1 tool for creating Demonstrations and anything technical.

Wolfram|Alpha »

Explore anything with the first computational knowledge engine.

Wolfram Demonstrations Project »

Explore thousands of free applications across science, mathematics, engineering, technology, business, art, finance, social sciences, and more.

Computerbasedmath.org »

Join the initiative for modernizing math education. Solve integrals with Wolfram|Alpha.

Online Integral Calculator »

Step-by-step Solutions »

Walk through homework problems step-by-step from beginning to end. Hints help you try the next step on your own.

Wolfram Problem Generator »

Unlimited random practice problems and answers with built-in Step-by-step solutions. Practice online or make a printable study sheet.

Wolfram Education Portal »

Collection of teaching and learning tools built by Wolfram education experts: dynamic textbook, lesson plans, widgets, interactive Demonstrations, and more.

Wolfram Language »

Knowledge-based programming for everyone.

Contact the MathWorld Team

© 1999-2019 Wolfram Research, Inc. | Terms of Use