

EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Course](#) > [Unit 2 Foundation of Inference](#) > [Homework 3: Introduction to Hypothesis Testing](#)

7. Quiz: Composite Hypotheses for  
> Bernoulli models

Currently enrolled in **Audit Track** (expires December 25, 2019) [Upgrade \(\\$300\)](#)

## 7. Quiz: Composite Hypotheses for Bernoulli models

(a)

1/1 point (graded)

Let  $X_1, \dots, X_n$  be i.i.d. **Bernoulli** random variables with unknown parameter  $p \in (0, 1)$ .

Find a function  $T_{n,p}(\bar{X}_n)$ , which depends on  $\bar{X}_n, n$ , and  $p$ , such that

$$T_{n,p}(\bar{X}_n) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}(0, 1),$$

by

- using the Central Limit Theorem on  $\bar{X}_n$  and
- **substituting any occurrence of  $p$  in the variance by a plug-in estimator for  $p$ .**

**Note:** If  $T_{n,p} \xrightarrow{(d)} \mathcal{N}(0, 1)$ , then so does  $-T_{n,p}$ . For this problem and the next part, use the expression for  $T_{n,p}(\bar{X}_n)$  that is of the form

Generating Speech Output  $f\left(\frac{\bar{X}_n - p}{\sqrt{\frac{p(1-p)}{n}}}\right)$  where  $f(n, \bar{X}_n)$  is always **positive**. (Or very loosely speaking, use  $(\bar{X}_n - p)$  and not  $(p - \bar{X}_n)$  where applicable.)

(Enter **barX\_n** for  $\bar{X}_n$ ).

$$T_{n,p}(\bar{X}_n) = \text{sqrt}(n) * (\text{barX}_n - p) / \text{sqrt}(\text{barX}_n * (1 - \text{barX}_n))$$



STANDARD NOTATION

Submit

You have used 2 of 3 attempts

✓ Correct (1/1 point)

(b)

3/3 points (graded)

(This is a quiz, hence only 1 attempt.)

Select a test with asymptotic level  $\alpha$ , in terms of the function  $T_{n,p}(\bar{X}_n)$ , for each of the following pairs of hypotheses:

(Choose one for each column. Note the absolute values in the first 2 rows.)

$H_0 : p = 0.5$  vs  $H_1 : p \neq 0.5$  :  $H_0 : p \leq 0.5$  vs  $H_1 : p > 0.5$  :  $H_0 : p \geq 0.5$  vs  $H_1 : p < 0.5$  :

Generating Speech Output

☒  $\mathbf{1} \left( \left| T_{n,0.5} \left( \bar{X}_n \right) \right| > q_{\alpha/2} \right)$

☐  $\mathbf{1} \left( \left| T_{n,0.5} \left( \bar{X}_n \right) \right| > q_{\alpha/2} \right)$

☐  $\mathbf{1} \left( \left| T_{n,0.5} \left( \bar{X}_n \right) \right| > q_{\alpha/2} \right)$

☐  $\mathbf{1} \left( \left| T_{n,0.5} \left( \bar{X}_n \right) \right| > q_{\alpha} \right)$

☐  $\mathbf{1} \left( \left| T_{n,0.5} \left( \bar{X}_n \right) \right| > q_{\alpha} \right)$

☐  $\mathbf{1} \left( \left| T_{n,0.5} \left( \bar{X}_n \right) \right| > q_{\alpha} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) > q_{\alpha/2} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) > q_{\alpha/2} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) > q_{\alpha/2} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) > q_{\alpha} \right)$

☒  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) > q_{\alpha} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) > q_{\alpha} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) < -q_{\alpha/2} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) < -q_{\alpha/2} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) < -q_{\alpha/2} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) < -q_{\alpha} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) < -q_{\alpha} \right)$

☒  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) < -q_{\alpha} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) < q_{\alpha/2} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) < q_{\alpha/2} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) < q_{\alpha/2} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) < q_{\alpha} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) < q_{\alpha} \right)$

☐  $\mathbf{1} \left( T_{n,0.5} \left( \bar{X}_n \right) < q_{\alpha} \right)$




You have used 1 of 1 attempt

✓ Correct (3/3 points)

 Generating Speech Output ☐ On

**Topic:** Unit 2 Foundation of Inference: Homework 3: Introduction to Hypothesis Testing / 7. Quiz: Composite Hypotheses for Bernoulli models

[Add a Post](#)

Show all posts ▼

by recent activity ▼

? [Why make the variance dependent on the estimator?](#)

3

💬 [Not sure if I understand the correct answer format for part a  
does the expression involve '>s'?](#)

1

**Audit Access Expires Dec 24, 2019**

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

[Learn About Verified Certificates](#)

© All Rights Reserved

Generating Speech Output