



[Course](#) > [Unit 1: Fourier Series](#) > [1. Introduction to Fourier Series](#) > 9. Meaning of the constant term

Audit Access Expires Jun 24, 2020

You lose all access to this course, including your progress, on Jun 24, 2020.

Upgrade by Jun 7, 2020 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

9. Meaning of the constant term

Recall that the average value of a function $f(t)$ on an interval (a, b) is defined to be the area under the curve over that interval, divided by the length of that interval:

$$\text{ave}(f) := \frac{\int_a^b f(t) dt}{b - a}.$$

The constant term of the Fourier series of f is

$$\frac{a_0}{2} = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) dt = \frac{\int_{-\pi}^{\pi} f(t) dt}{\pi - (-\pi)},$$

which is the average value of f on the interval $(-\pi, \pi)$.

Example 9.1



Find the constant term of the 2π -periodic function

$$f(t) = \begin{cases} t, & 0 < t < \pi \\ 0, & -\pi < t < 0 \end{cases}.$$

Solution: The average value of this function over one period is given by the area under the curve, divided by the length of the interval, which in this case is 2π . By sketching the function over the interval $-\pi < t < \pi$,




Figure 4: The graph of $f(t)$ over the interval $-\pi < t < \pi$.

we can compute the area under the curve directly to find the average value, which is

$$\frac{(1/2)\pi^2}{2\pi} = \frac{\pi}{4}.$$

Find the constant term, 1

1/1 point (graded)

Find the constant term of Fourier series for the following 2π -periodic function defined by

$$f(t) = |t|, \quad -\pi < t < \pi.$$




(This function is called the triangle wave.)

$$\frac{a_0}{2} = \boxed{\pi/2} \quad \checkmark \text{ Answer: } \pi/2$$

$\frac{\pi}{2}$

Solution:

Sketch the function to determine the area under the graph over the interval $-\pi < t < \pi$.



The area under the curve is π^2 , and the length of the interval is 2π , so the average value of this function on this interval is $\pi/2$.

Submit

You have used 1 of 5 attempts

i Answers are displayed within the problem

Find the constant term, 2

1/1 point (graded)

Find the constant term of Fourier series for the following 2π periodic function defined by



$$f(t) = t, \quad -\pi < t < \pi.$$

(This function is called the sawtooth wave.)

$$\frac{a_0}{2} =$$


0

✔ Answer: 0

0

Solution:

First we sketch the function.



The area of this odd function over the interval is zero, therefore the average value is 0.

Submit

You have used 1 of 5 attempts



Find the constant term, 3

1/1 point (graded)


Find the constant term of Fourier series for the following 2π -periodic function.

$$f(t) = \sqrt{\pi^2 - t^2}, \quad -\pi < t < \pi.$$

$\frac{a_0}{2} =$ ✓ Answer: $\pi^2/4$

Solution:

The graph of this function is a semicircle of radius π centered at zero, and repeated periodically.



The area under this curve from $-\pi < t < \pi$ is the area of half a circle of radius π , which is $\pi^3/2$. Dividing by the length of the interval, which is 2π , we get the average value, which is $\pi^2/4$.

Submit

You have used 1 of 5 attempts



Answers are displayed within the problem

Find the constant term, 4

1/1 point (graded)

Find the constant term of Fourier series for the following 2π -periodic function.

$$f(t) = t^2, \quad -\pi < t < \pi.$$

$$\frac{a_0}{2} =$$

$\pi^2/3$

✓ Answer: $\pi^2/3$

Solution:

The graph of this function is seen below.



The area under this curve from $-\pi < t < \pi$ is

$$\int_{-\pi}^{\pi} t^2 \, dt = \left. \frac{t^3}{3} \right|_{-\pi}^{\pi} = \frac{2\pi^3}{3}.$$

Dividing by the length of the interval, which is 2π , we get the average value, which is $\pi^2/3$.

Submit

You have used 1 of 5 attempts

Answers are displayed within the problem



9. Meaning of the constant term

Topic: Unit 1: Fourier Series / 9. Meaning of the constant term

Hide Discussion

Add a Post

Show all posts ▼

by recent activity ▼

? [Example 9.1?](#)

if I integrate $f(t)$ from $[-\pi, \pi]$, the integral from $[-\pi, 0]$, is 0 since f is zero on that interval. The remaining integral from $[0, \pi]$ of t is $t^2/2$ giving $\pi^2/2$. Divide by π I get $\pi/2$, no...

2 ▼

Learn About Verified Certificates

© All Rights Reserved

