



[Course](#) > [Unit 1: Fourier Series](#) > [Period 2L](#) > [2. Properties of Fourier Series \(of](#) > [5. Convergence of a Fourier series](#)


Audit Access Expires Jun 24, 2020

You lose all access to this course, including your progress, on Jun 24, 2020.

Upgrade by Jun 7, 2020 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

5. Convergence of a Fourier series

Convergence at jump discontinuities



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Definition 5.1 A periodic function f of period $2L$ is called **piecewise differentiable** if

- at points where the derivative exists it is bounded, (that is, there is an M such that $|f'(t)| \leq M < \infty$ at all t for which $f'(t)$ exists),



- there are at most finitely many points in $[-L, L)$ where $f'(t)$ does not exist, and
- at each such point τ , the left limit $f(\tau^-) := \lim_{t \rightarrow \tau^-} f(t)$ and right limit $f(\tau^+) := \lim_{t \rightarrow \tau^+} f(t)$ exist (although they might be unequal, in which case we say that f has a **jump discontinuity** at τ).

Theorem 5.2 If f is a piecewise differentiable periodic function, then the Fourier series of f (with the a_n and b_n defined by the Fourier coefficient formulas)

- converges to $f(t)$ at values of t where f is continuous, and
- converges to $\frac{f(t^-) + f(t^+)}{2}$ where f has a jump discontinuity.

Example 5.3 The left limit $\text{Sq}(0^-) = -1$ and right limit $\text{Sq}(0^+) = 1$ average to 0. The Fourier series

$$\frac{4}{\pi} \left(\sin t + \frac{\sin 3t}{3} + \frac{\sin 5t}{5} + \dots \right)$$

evaluated at $t = 0$ converges to 0 too.

(Note that the examples here are for 2π -periodic functions, but this theorem on convergence holds for periodic functions of any period, which we begin to discuss on the next page.)

Practice 1

1/1 point (graded)

Consider the 2π -periodic function

$$f(t) = \begin{cases} t & 0 < t < \pi \\ 0 & -\pi < t < 0 \end{cases}.$$

Let $g(t)$ be the Fourier series for $f(t)$. What is $g(\pi)$?



$g(\pi) =$

pi/2

✓ Answer: pi/2

$\frac{\pi}{2}$

[FORMULA INPUT HELP](#)

Solution:

The function $f(t)$ has a jump discontinuity at π . Note that $f(\pi^-) = \pi$ and $f(\pi^+) = 0$, therefore

$$g(\pi) = \frac{f(\pi^-) + f(\pi^+)}{2} = \frac{\pi + 0}{2} = \frac{\pi}{2}.$$

Submit

You have used 1 of 5 attempts

i Answers are displayed within the problem

Practice 2

1/1 point (graded)

Let $U(t)$ be the Fourier series of the 2π -periodic triangle wave

$$T(t) = |t|, \quad -\pi < t < \pi.$$

Find the value of $U(\pi)$.

$U(\pi) =$

pi

✓ Answer: pi

π

[FORMULA INPUT HELP](#)

Solution:

This function is continuous, so the value of the Fourier series is the value of the function at these points, which is π .

Submit

You have used 1 of 5 attempts

 Answers are displayed within the problem

5. Convergence of a Fourier series

[Hide Discussion](#)

Topic: Unit 1: Fourier Series / 5. Convergence of a Fourier series

[Add a Post](#)

Show all posts ▼

by recent activity ▼



[g\(pi\)](#)

[Please, This is a Odd or Even function? In mt point is a Even function, so you have only the an coeficiente, ok? Thanks, Ricardo](#)

6 ▼



[How to enter the sigma notation?](#)

[In order to answer the question 1 and 2, we need to enter the series which contains sigma notation, how I am supposed to enter it?](#)

2 ▼



[Piecewise differentiability - the first point](#)

3 ▼

[Learn About Verified Certificates](#)

© All Rights Reserved

