



[Course](#) > [Unit 4 Hypothesis testing](#) > [T-Test](#) [Lecture 13: Chi Squared Distribution](#), 7. The Chi-Squared Distribution and its Properties

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

7. The Chi-Squared Distribution and its Properties

The Chi-Squared Distribution and its Expectation

3/3 points (graded)

Note: This problem introduces the chi-squared distribution and is intended as an exercise in probability that you are encouraged to attempt before watching the following video.

The χ_d^2 **distribution with d degrees of freedom** is given by the distribution of

$$Z_1^2 + Z_2^2 + \cdots + Z_d^2,$$

where $Z_1, \dots, Z_d \stackrel{iid}{\sim} \mathcal{N}(0, 1)$.

What is the smallest possible sample space of χ_d^2 ?

☐ $\mathbb{Z}_{\geq 0}$

☐ \mathbb{Z}

☒ $\mathbb{R}_{\geq 0}$

☐ \mathbb{R}



If $X \sim \chi_d^2$, what is $\mathbb{E}[X]$? Give your answer in terms of d .

d

✓ Answer: d

d

STANDARD NOTATION

Let $\mathbf{Z} \sim \mathcal{N}(0, I_{d \times d})$ denote a random vector whose components are standard Gaussians: $Z^{(1)}, \dots, Z^{(d)} \sim \mathcal{N}(0, 1)$. Which one of the following random variables has a chi-squared distribution with d degrees of freedom?

☐ $\max(Z^{(1)}, \dots, Z^{(d)})$

☐ $|Z^{(1)}| + |Z^{(2)}| + \dots + |Z^{(d)}|$

☐ $\|\mathbf{Z}\|_2$

☒ $\|\mathbf{Z}\|_2^2$



Solution:

The smallest sample space of a Gaussian random variable Z is \mathbb{R} . Hence, the smallest possible sample space of Z^2 is $\mathbb{R}_{\geq 0}$. And the same holds for the sum

$$Z_1^2 + Z_2^2 + \cdots + Z_d^2,$$

so the smallest possible sample space for χ_d^2 is $\mathbb{R}_{\geq 0}$.

Next, by linearity of expectation,

$$\mathbb{E}[X] = \mathbb{E}[Z_1^2 + Z_2^2 + \cdots + Z_d^2] = d \cdot 1 = d,$$

because $Z_1, \dots, Z_d \stackrel{iid}{\sim} \mathcal{N}(0, 1)$.

The ℓ_2 norm $\|\cdot\|_2$ measures the Euclidean distance from the origin. Hence, if $\mathbf{Z} \sim \mathcal{N}(0, I_{d \times d})$, then

$$\|\mathbf{Z}\|_2^2 = \left(Z^{(1)}\right)^2 + \left(Z^{(2)}\right)^2 + \cdots + \left(Z^{(d)}\right)^2 \sim \chi_d^2.$$

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Distribution of Sample Variance of Gaussian: The Chi-Squared Distribution



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Throwing Darts

1/1 point (graded)

You are playing darts on a dart-board that is represented by the entire plane, \mathbb{R}^2 . You get a 'bullseye' if the dart lands inside of the unit disc $D^1 := \{(x, y) : x^2 + y^2 \leq 1\}$. Your dart throws are modeled by a Gaussian random vector \mathbf{Z} , where $Z^{(1)}, Z^{(2)} \stackrel{iid}{\sim} \mathcal{N}(0, 1)$.

Let f_d represent the density of the χ_d^2 distribution.

Which of the following equals the probability of getting a bullseye?

☐ $\int_0^1 f_1(x) dx$

☒ $\int_0^1 f_2(x) dx$

☐ $\int_1^\infty f_2(x) dx$

☐ $\int \int_{D^1} f_2(x) dx dy$



Solution:

A bullseye is given by the event $(Z^{(1)})^2 + (Z^{(2)})^2 \leq 1$. Since $(Z^{(1)})^2 + (Z^{(2)})^2 \sim \chi_2^2$, it follows that

$$P(\text{bullseye}) = \int_0^1 f_2(x) dx.$$


Remark: The $d = 2$ case is special, because it turns out that $\chi_2^2 = \text{Exp}(1/2)$. This can be seen using the explicit formula for the density of a χ_2^2 , but it is not necessary for this course to know the density of a chi-squared random variable with d degrees of freedom by heart.

Submit

You have used 1 of 2 attempts

i Answers are displayed within the problem

Properties of the Chi-Squared Distribution



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

The Chi-Squared Distribution and the Sample Second Moment

2/2 points (graded)

Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(0, \sigma^2)$ and let

$$V_n = \frac{1}{n} \sum_{i=1}^n X_i^2$$

denote the sample second moment. For an appropriate expression A given in terms of n and σ^2 , we have that $AV_n \sim \chi^2$.

What is A ?

$A =$ ✓ Answer: n/sigma^2

How many degrees of freedom does the above χ -squared random variable have? (Give your answer in terms of n .)

✓ Answer: n

STANDARD NOTATION

Solution:

Observe that

$$\frac{n}{\sigma^2} V_n = \sum_{i=1}^n \frac{X_i^2}{\sigma^2} = \sum_{i=1}^n \left(\frac{X_i}{\sigma} \right)^2,$$

and $X_i/\sigma \sim \mathcal{N}(0, 1)$ because $X_i \sim \mathcal{N}(0, \sigma^2)$. Hence, $\frac{n}{\sigma^2}V_n$ is a χ_n^2 random variable.

Submit

You have used 2 of 3 attempts

i Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 4 Hypothesis testing:Lecture 13: Chi Squared Distribution, T-Test / 7. The Chi-Squared Distribution and its Properties

Add a Post

Show all posts ▼

by recent activity ▼



[Staff] Throwing darts problem

can you please add a reasoning in the solution, why the incorrect answer choices are incorrect?

5

Learn About Verified Certificates

© All Rights Reserved