







< Previous	 	 			Next >
------------	---	---	---	---	--------

12.3.3 Diagonalization, Again

 Bookmark this page

Week 12 due Dec 29, 2023 10:42 IST Completed

12.3.3 Diagonalization, Again

Video

Start of transcript. Skip to the end.

Dr. Robert van de Geijn: A square matrix A that's n by n ,
can only be diagonalized if A has n linearly independent eigenvectors.
Often when you try to prove an if and only if, you prove it in one direction first, and then in the other direction.
So what we're going to start with is, assume that A can be diagonalized

▶ 0:00 / 0:00

▶ 2.0x

🔊

🔍

CC

🗣

Video

📄 [Download video file](#)

Transcripts

📄 [Download SubRip \(.srt\) file](#)

📄 [Download Text \(.txt\) file](#)

Reading Assignment

0 points possible (ungraded)
Read Unit 12.3.3 of the notes. [\[LINK\]](#)

☒ Done

✓

Submit

✓ Correct

Discussion

Topic: Week 12 / 12.3.3

Hide Discussion

Add a Post

Show all posts ▼

by recent activity ▼

There are no posts in this topic yet.

✕

Calculator

Homework 12.3.3.1

10.0/10.0 points (graded)

Consider $A = \begin{pmatrix} 2 & 1 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 2 \end{pmatrix}$.

- The algebraic multiplicity of $\lambda = 2$ is

✔ Answer: 4

4.

- The geometric multiplicity of $\lambda = 2$ is

✔ Answer: 2

2.

- The following vectors are linearly independent eigenvectors associated with $\lambda = 2$:

$$\begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}.$$

TRUE

▼

✔ Answer: TRUE

Just multiply out.

Submit

i Answers are displayed within the problem

Homework 12.3.3.2

1/1 point (graded)

Let $A \in \mathbb{A}^{n \times n}$, $\lambda \in \Lambda(A)$, and S be the set of all vectors x such that $Ax = \lambda x$. Finally, let λ have algebraic multiplicity k (meaning that it is a root of multiplicity k of the characteristic polynomial).

The dimension of S is k ($\dim(S) = k$).

Sometimes

▼

✔ Answer: Sometimes

An example of where it is true:

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

This matrix has 1 as its only eigenvalue, and it has algebraic multiplicity two. But both $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ are corresponding eigenvectors and hence $S = \mathbb{R}^2$, which has dimension 2 .

An example of where it is false:

Calculator

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

This matrix has **1** as its only eigenvalue, and it has algebraic multiplicity two. Now, to find the eigenvectors we consider

$$\begin{pmatrix} 1 - 1 & 1 \\ 0 & 1 - 1 \end{pmatrix} = \begin{pmatrix} 0 & \boxed{1} \\ 0 & 0 \end{pmatrix}.$$

It is in row echelon form and has one pivot. Hence, the dimension of its null space is $2 - 1 = 1$. Since \mathcal{S} is this null space, its dimension equals one.

Submit

 Answers are displayed within the problem

< Previous

Next >

© All Rights Reserved



edX

[About](#)

[Affiliates](#)

[edX for Business](#)

[Open edX](#)

[Careers](#)

[News](#)

Legal

[Terms of Service & Honor Code](#)

[Privacy Policy](#)

[Accessibility Policy](#)

[Trademark Policy](#)

[Sitemap](#)

[Cookie Policy](#)

[Your Privacy Choices](#)

Connect

[Idea Hub](#)

[Contact Us](#)

[Help Center](#)

[Security](#)

[Media Kit](#)



 Calculator



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)