

Statistics How To

Statistics for the rest of us!



[HOME](#) [TABLES](#) [PROBABILITY AND STATISTICS](#) [CALCULATORS](#) [STATISTICS BLOG](#) [MATRICES](#) [EXPERIMENTAL DESIGN](#)

[PRACTICALLY CHEATING STATISTICS HANDBOOK](#)



Lilliefors Test for Normality & Exponential Distributions

[Normality](#) > Lilliefors Test

What is Lilliefors Test?

The Lilliefors test is a **test for normality**. It is an improvement on the [Kolmogorov-Smirnov \(K-S\) test](#) — correcting the K-S for small values at the tails of probability distributions — and is therefore sometimes called the *K-S D test*. Many statistical packages (like SPSS) combine the two tests as a “Lilliefors corrected” K-S test.

Unlike the K-S test, Lilliefors can be used when you don’t know the [population mean](#) or [standard deviation](#). Essentially, the Lilliefors test is a K-S test that allows you to estimate these parameters from your sample.

Running the Test

- The null hypothesis (H_0) for the test is the data comes from a normal distribution.
- The alternate hypothesis (H_1) is that the data *doesn't* come from a normal distribution.

The test assumes that you have a random sample.

If the test statistic is significantly large, you can [reject the null hypothesis](#) and conclude that the data is not normal.

Calculation Steps

The formulas involved in the Lilliefors test are quite time consuming (in part due to the fact that you have to calculate [z-scores](#) for every single sample member!), so the test is usually performed with software:

- **Minitab**: At the time of writing, Minitab doesn’t have an option for this test.
- In **R**: Use `lillie.test` in the package `nortest`.
- In **SPSS** (21.0.0.1 and later only), run the K-S test, click “Options” and specify *normal*. If you don’t enter the population mean and standard deviation, SPSS will automatically run Lilliefors. A footnote will appear in the output indicating that the correction was used.

The general steps that the test follows are:


1. Calculate X_i using this formula:

$$Z_i = \frac{X_i - \bar{X}}{s}, \quad i = 1, 2, \dots, n$$

Where:

- Z_i = the individual z-scores for every member in your sample,

Find an article



Feel like "cheating" at Statistics? Check out the grade-increasing book that's recommended reading at top universities!

- X_i = individual member / data point,
- \bar{X} = sample mean,

2. Calculate the test statistic, which is the empirical distribution function (EDF) based on the Z_i s. The formula is

$$T_1 = \sup_x |F^*(x) - S(x)|$$

Where:

$F^*(x)$ = the standard normal distribution function.

$S(x)$ = the empirical distribution function of the z_i values.

3. Find the critical value for the test from [this table](#) and reject the null hypothesis if the test statistic T is greater than the critical value.

Cautions

Lilliefors test has a low [power](#). According to [R documentation](#), it is “known to perform worse” than either the Anderson-Darling or Cramer-von Mises tests.

Lilliefors test for Exponential Distribution

Lilliefors can also be used to test for a fit to an exponential distribution. The general steps are almost identical to the Lilliefors normality test, except with a couple of minor changes to the null hypothesis and formulas:

- The null hypothesis (H_0) for the test is the data comes from an exponential distribution.
- The alternate hypothesis (H_1) is that the data *doesn't* come from an exponential distribution.

The test assumes that you have a random sample.

The software commands are almost the same, except you would specify exponential instead of normal:

1. In **Matlab**, specific exp in the Lilliefors command: `h = lillietest([variable name], 'Distr', 'exp')`. If the test returns a 1, the data does not fit an exponential distribution.
2. In **SPSS**, click the “Options” button for the K-S test and choose *exponential*.

Reference:

IBM. K-S test of normality in NPAR TESTS and NPTESTS does not use Lilliefors correction prior to SPSS Statistics 21.0.0.1. Retrieved Nov 18 2016 from [here](#).

Lilliefors, H. W. (1967). On the Kolmogorov-Smirnov test for normality with mean and variance unknown, Journal of the American Statistical Association, 62, 399–402.

Need help with a homework or test question? With [Chegg Study](#), you can get step-by-step solutions to your questions from an expert in the field. Your first 30 minutes with a Chegg tutor is free!

Install our FREE Android app on Google Play !



Need help NOW with a homework problem? [Click here!](#)

Statistical concepts explained visually - Includes many concepts such as sample size, hypothesis tests, or logistic regression, explained by [Stephanie Glen](#), founder of StatisticsHowTo.

Comments? Need to post a correction? Please post a comment on our [Facebook page](#).

[Check out our updated Privacy policy and Cookie Policy](#)

By Stephanie | March 8, 2016 | Statistics How To |

[← Assumption of Normality / Normality Test](#)

[Finite Population Correction Factor FPC: Formula, Examples →](#)

Copyright ©2019 Statistics How To | Theme by: Theme Horse | Powered by: WordPress

We encourage you to view our updated policy on cookies and affiliates. [Find out more.](#)

Okay, thanks

